

Smoking and diabetes

Introduction

This fact sheet examines the association between smoking and diabetes including smoking as a risk factor, how smoking can lead to multiple complications of diabetes and the benefits of stopping smoking among people with diabetes.

What is diabetes?

Diabetes mellitus is a metabolic condition which causes increased glucose levels in the blood. Glucose is a sugar that the body produces primarily from the digestion of carbohydrates and levels are controlled by the hormone insulin. Insulin is made and stored in the pancreas and helps glucose to enter the cells where it is used as fuel by the body.¹

Types of diabetes

There are two types of diabetes:

Type 1 Diabetes (insulin dependent) develops when the insulin-producing cells in the pancreas have been destroyed, leaving the body unable to produce insulin. It is thought that the body's immune system destroys the cells but it's not known what causes this to happen. Type 1 Diabetes usually appears in children and young adults but can occur at any age. People with this form of the condition are treated with daily insulin injections and account for between 5 and 15 per cent of diabetes cases in the UK.¹

Type 2 Diabetes (insulin resistance) develops while the body can still produce insulin, but not enough, or when the insulin produced doesn't work properly. This form of the condition usually develops gradually after the age of 40. However, it is increasingly being diagnosed in younger people, including children. Both genetic and environmental factors contribute to the development of diabetes but the development of Type 2 diabetes is more likely if some or all of the following factors are also present: physical inactivity; being overweight; family history of Type 2 diabetes; previous diabetes in pregnancy. The condition is also more common in people of Asian and African-Caribbean origin.^{1 2 3}

Prevalence and health consequences of diabetes

In the UK, 3 million people have been diagnosed with diabetes, equivalent to about 5% of the population.¹ It has been estimated that there are a further 850,000 people with diabetes who remain undiagnosed. In 2010 there were 6,626 diabetes-related deaths recorded in the United Kingdom.

People with diabetes are at greater risk of:

- high blood pressure
- heart disease
- stroke
- kidney disease
- nerve damage leading to limb amputation
- eye damage such as retinopathy (disorders of the retina)⁶

Smoking and the risk of developing diabetes

There is a growing body of evidence to show that smoking is a risk factor for Type 2 diabetes. Several hypotheses have been proposed to explain this link.^{7 8 9 10} Smoking has been identified as a possible risk factor for insulin resistance (see below), a precursor for diabetes. Smoking has also been shown to deteriorate glucose metabolism which may lead to the onset of type 2 diabetes.¹¹ There is also some evidence which suggests that smoking increases diabetes risk through a body mass index independent mechanism.^{12 13}

Smoking has been associated with a risk of chronic pancreatitis and pancreatic cancer, suggesting that tobacco smoke may be toxic to the pancreas.^{14 15}

A systematic review of 25 studies found that all but one revealed an association between active smoking and an increased risk of diabetes.¹⁶ On the basis of this review, it is estimated that 12% of all type 2 diabetes in the United States may be attributable to smoking.¹⁷ If the same proportion is applied to the UK, smoking may account for as many as 360,000 cases of diabetes.

Smoking-related risk of diabetes increases with the number of cigarettes smoked. The Cancer Prevention Study 1, a cohort study of 275 190 men and 434 637 women, found that women who smoke more than 40 cigarettes a day have a 74% increased risk of developing diabetes, with men's risk increasing by 45%.¹⁸

There is also some evidence, including a 2011 cohort study of over 10,000 people, to suggest that exposure to secondhand smoke may be a risk factor for the development of type 2 diabetes.^{19 20 21 22}

Smoking as an aggravating factor of diabetes

People with diabetes already have an increased risk of heart disease, which is further elevated if they smoke. Diabetes acts in several ways to damage the heart: high glucose levels affect the walls of the arteries making them more likely to develop fatty deposits which in turn makes it more difficult for the blood to circulate. People with diabetes are more likely to have high blood pressure and high levels of fats such as triglycerides. They are also more likely to have lower levels of the protective HDL cholesterol.

Smoking, diabetes and pregnancy

Women who smoke during pregnancy are at increased risk of developing gestational diabetes and also increase the risk of their offspring developing diabetes later in life.²³ Gestational diabetes mellitus is defined as any carbohydrate intolerance that begins during pregnancy. Women who develop diabetes during pregnancy have a seven-fold increased risk of subsequently developing type 2 diabetes compared with women who have normal levels of glucose in pregnancy.²⁴

Metabolic syndrome

Metabolic syndrome is a condition in which a person has a number of different medical problems, all related to the body's metabolism, which together increase their risk of developing coronary heart disease and diabetes. These medical problems include: Type 2 diabetes, high blood pressure, high blood triglyceride levels, and low levels of HDL cholesterol (the protective type of cholesterol) in the blood.

Most commonly, patients suffering from metabolic syndrome will be overweight, particularly around the waist, and have resistance to insulin. Recent evidence suggests a strong association between cigarette smoking, insulin resistance and metabolic syndrome.^{25 26}

Smoking, diabetes and premature death

Compared to non-smokers with diabetes, people with diabetes who smoke have twice the risk of premature death. Furthermore, the risk of complications associated with tobacco use and diabetes in combination is nearly 14 times higher than the risk of either smoking or diabetes alone.²⁷

A large prospective study of US nurses found that among those with diabetes the relative risks of mortality were 1.31 for past smokers, 1.43 for current smokers of 1-14 cigarettes per day, 1.64 for smokers of 15-34 cigarettes per day, and 2.19 for current smokers of 35 or more cigarettes per day.²⁸

The effect of smoking on complications of diabetes

Smoking is associated with multiple complications of diabetes. Nephropathy (kidney disease) has been shown to be common in Type 1 diabetic patients who smoke²⁹ and smoking increases the risk of albuminuria in both types of diabetes.^{30 31} (Albuminuria refers to the presence of protein in the urine and can indicate signs of kidney disease.) Another small study of 33 people with Type 2 diabetes with kidney disease found that smokers' kidney function declined more rapidly than that of non-smokers, despite drug treatment, suggesting that smoking cessation could slow the progression of kidney disease in people with diabetes who use ACE inhibitors.³²

The relationship between cigarette smoking and retinopathy (disorders of the retina) is less well defined than that of other microvascular complications of diabetes.³³ However, some studies have found an association between smoking and diabetic retinopathy.^{30 34}

Smoking is also a documented risk factor for both the development and progression of various types of neuropathy (damage to the peripheral nervous system). A retrospective case control study of type 1 and type 2 diabetic patients found that current or ex-smokers were significantly more likely to have neuropathy than individuals who never smoked (64.8% vs. 42.8%).³⁵ A more recent prospective study found that cigarette smoking was associated with a 2-fold increase in risk.³⁶

Benefits of stopping smoking

Stopping smoking reduces the risk of cardiovascular disease, lung disease, cancer and stroke.³⁷ (See also ASH fact sheet: Stopping smoking - the benefits and aids to quitting.) As diabetes also increases the risk for heart disease and stroke, smokers with diabetes are strongly advised to quit.³⁸ However it appears that many smokers with diabetes are not receiving this advice. As part of a regular survey of its members, Diabetes UK found that 42% of smokers had not received support or advice to quit.³⁹ Few studies have evaluated smoking cessation treatment specifically for people with diabetes but the limited research available suggests that smokers with diabetes may be less successful in quitting than smokers without diabetes and those intensive strategies should be considered to optimise successful cessation.³⁴

One possible explanation for the lower quitting rates among people with diabetes is the fact that stopping smoking is associated with weight gain and this is likely to be of concern in people who have diabetes and are already overweight. One US study found that concerns about weight gain among smokers with Type 1 diabetes were particularly prevalent among women, obese smokers, and those in poor metabolic control.⁴⁰ Fear of weight gain was cited by 49% of smokers.

A British prospective study of 7,735 men aged 40-59 years found that the benefit of giving up smoking was only apparent after 5 years of smoking cessation and risk reverted to that of never-smokers only after 20 years. Men who gave up smoking during the first 5 years of follow-up showed significant weight gain and subsequently higher risk of diabetes than continuing smokers. However, the authors concluded that in the long term, the benefits of giving up smoking outweigh the adverse effects of early weight gain.⁴¹ The US Cancer Prevention Study also provided evidence that stopping smoking for 10 years in men and five years in women could reduce the risk of diabetes to that of nonsmokers.¹⁹

Stopping smoking also reduces the risk of premature death. The US Nurses' Study found that women with Type 2 diabetes who had stopped smoking for 10 or more years had a mortality relative risk of 1.11 compared with diabetic women who were never smokers.²⁹

In the light of evidence demonstrating that smoking is an independent risk factor for diabetes and that it is also an aggravating factor for diabetes complications, smoking cessation advice should be a routine component of diabetic care. Concerns about weight gain should be addressed by health care providers whilst emphasising the fact that the health benefits of smoking cessation far outweigh post cessation weight gain, even in people who are focused on weight management.⁴²

References

- 1 Diabetes UK website. [“What is diabetes”](#).
- 2 Diabetes UK website. [“Diabetes and ethnicity”](#). [Accessed 27 Jan. 2012.]
- 3 Oldroyd J, Banerjee M, Heald A et al. [Diabetes and ethnic minorities](#). Postgraduate Medical Journal, 2005. 81: 486-490.
- 4 Estimate provided by Diabetes UK based on APHO Diabetes Prevalence Model and 2010 QOF data.
- 5 Office for National Statistics. [Mortality Statistics: Deaths registered in England and Wales \(Series DR\)](#), 2010. Scotland Public Health Observatory website. Health, Wellbeing and Disease. Diabetes. [“Diabetes Key Points”](#). Northern Ireland Statistics and Research Agency. [Statistical Bulletin: Deaths in Northern Ireland](#).
- 6 US National Library of Medicine website. [Diabetes](#). [Accessed 27 Jan. 2012.]
- 7 Hsin-Chieh Y, Bruce D, Schmidt MA et al. [Smoking, smoking cessation and risk for type 2 diabetes mellitus](#). Annals of Internal Medicine, 2010. 152. 1: 10-17.
- 8 Ko G & Cockram C. [Cause as well as effect: smoking and diabetes](#). Diabetes Voice: Smoking and diabetes special issue, 2005; 50: 19-22.
- 9 Rimm E, Chan J, Stampfer M et al. Prospective study of cigarette smoking, alcohol use, and the risk of diabetes in men, British Medical Journal, 1995 310: 555–559.
- 10 Radzeviciene L, Ostrauskas R. [Smoking habits and the risk of type 2 diabetes: a case control study](#). Diabetes and Metabolism, 2009. 35. 3:192-7.
- 11 Fagard RH, Nilsson PM. [Smoking and diabetes – the double health hazard](#). Primary Care Diabetes, 2009. 3. 4: 2005-9.
- 12 Cullen MW, Ebbert JO, Vierkant RA et al. [No interaction of body mass index and smoking on diabetes mellitus risk in elderly women](#). Preventative Medicine, 2009. 48.1: 74-78.
- 13 Nagaya T, Yoshida H, Takahashi H et al. [Heavy smoking raises risk for type 2 diabetes mellitus in obese men: but, light smoking reduces the risk in lean men: a follow up study in Japan](#). Annals of Epidemiology, 2008. 18. 2: 113-8.
- 14 Johns Hopkins Medicine website. The Sol Goldman Pancreatic Cancer Research Center (sic). [What are the risk factors for pancreatic cancer?](#) [Accessed 27 Jan. 2012.]
- 15 Lynch SM, Vrieling A, Lubin JH et al. Cigarette smoking and pancreatic cancer: a pooled analysis from the pancreatic cancer cohort consortium. American Journal of Epidemiology, 2009. 170. 4: 403-13.
- 16 Willi, C et al. [Active smoking and the risk of Type 2 diabetes](#). Journal of the American Medical Association. 2007; 298: 2654-2664.
- 17 Ding EL & Hu FB. [Smoking and Type 2 diabetes](#). Underrecognized risks and disease burden. JAMA, 2007; 298: 2675-76
- 18 Will JC, Galuska DA, Ford ES, Mokdad A, Calle EE. [Cigarette smoking and diabetes mellitus: evidence of a positive association from a large prospective cohort study](#). International Journal of Epidemiology, 2001; 30: 540-6.
- 19 Houston T. [Active and passive smoking and development of glucose intolerance among young adults in a prospective cohort: CARDIA study](#). British Medical Journal, 2006; 332: 1064-1069.
- 20 Hayashino Y, Fukuhara S, Okamura T et al. [A prospective study of passive smoking and risk of diabetes in a cohort of workers](#). Diabetes Care 2008; 31:732-734.
- 21 Kowall B, Rathmann W, Strassburger K et al. [Association of passive and active smoking with incident type 2 diabetes mellitus in the elderly population: the KORA S/4/F4 cohort study](#). European Journal of Epidemiology, 2010. 25. 6:393-402.
- 22 Kwang-Pil K, Haesook M, Younjhin A et al. [A prospective study investigating the association between environmental tobacco smoke exposure and the incidence of type 2 diabetes in never smokers](#). Annals of Epidemiology, 2011. 21. 1: 42-47.
- 23 Montgomery S. [A very bad start: smoking, pregnancy and diabetes](#). Diabetes Voice: Smoking and diabetes special issue, 2005; 50: 30-32.
- 24 Bellamy L, Casas J-P, Hingorani A et al. [Type 2 diabetes mellitus after gestational diabetes: a systematic review and meta-analysis](#). The Lancet, 2009. 373: 1773-1779.
- 25 Sherwood Forest Hospitals NHS Foundation Trust. [Co-existence of smoking and metabolic syndrome among middle-aged patients with diabetes in the UK: a cross-sectional analysis](#). Diabetes Care, 2007; 4 (3): 241-242.
- 26 Ako G & Cockram C. [Cause as well as effect: smoking and diabetes](#). Diabetes Voice: Smoking and diabetes special issue, 2005; 50: 19-22.
- 27 Haire-Joshu D & Thomas J. [Gambling with addiction: Dangerous beliefs about smoking and diabetes](#). Diabetes Voice Smoking and diabetes special issue, 2005. 50: 15-18.
- 28 Al-Delaimy, WK, Willet WC, Maon JE et al. [Smoking and mortality among women with type 2 diabetes: The Nurses’ Health Study cohort](#). Diabetes Care, 2001; 12: 2043-8.
- 29 Mulhauser I, Bender R, Bott U, et al. [Cigarette smoking and progression of retinopathy and nephropathy in type 1 diabetes mellitus](#). Diabetes Medicine, 1996; 13: 536-543.

- 30 Ritz E, Keller C, Bergis K. [Nephrothapy of type II diabetes mellitus](#). Nephrol DialTransplant, 1996; 11 *Suppl
9 38-44.
- 31 Chase HP, Garg SK, Marshall G et al. [Cigarette smoking increases the risk of albuminuria among subjects
with type 1 diabetes](#). Journal of the Amerian Medical Association, 1991; 265. 5: 614-617.
- 32 Chuahirun, T and Wesson, D.E. [Cigarette smoking predicts faster progression of type 2 established diabetic
nephropathy despite ACE inhibition](#). American Journal of Kidney Diseases, 2002; 39: 376-382.
- 33 Haire-Josu D, Glasgow R.E, Tibbs, T.L. [Smoking and diabetes. \(Technical Review\)](#) Diabetes Care, 1999; 22.
11: 1887-1898.
- 34 Reichard P. [Risk factors for progression of microvascular complications in the Stockholm Diabetes
Intervention Study](#). Diabetes Research and Clinical Practice, 1992; 16: 151-156.
- 35 Mitchell B, Hawthorne V and Vinik A. [Cigarette smoking and neuropathy in diabetic patients](#). Diabetes Care,
1990; 13: 434-447.
- 36 Sands M, Shetterly SM, Franklin GM et al. Incidence of distal symmetric (sensory) neuropathy in NIDDM: the
San Luis Diabetes Study. Diabetes Care, 1997; 20: 322-329.
- 37 US Department of Health and Human Services. [The health benefits of smoking cessation: A report of the
Surgeon General](#), 1990.
- 38 Fagard RH, Nilsson PM. [Smoking and diabetes – the double health hazard](#). Primary Care Diabetes, 2009. 3.
4: 2005-9.
- 39 Diabetes UK. [15 Healthcare Essentials Survey](#), Nov. 2011.
- 40 Haire-Joshu D, Heady S, Thomas L et al. [Beliefs about smoking and diabetes care](#). Diabetes Education,
1994; 20: 410-415.
- 41 Wannamethee SG, Shaper AG, Pery IJ. [Smoking as a modifiable risk factor for type 2 diabetes in middle-
aged men](#). Diabetes Care, 2001; 24: (9) 1590-1595.
- 42 American Diabetes Association. Smoking and diabetes. Diabetes Care, 2003; 26 (S1): S89-90.