



Geomicrobiology Journal

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/ugmb20>

Microbial Activity and $^{13}\text{C}/^{12}\text{C}$ Ratio as Evidence of N-Hexadecane and N-Hexadecanoic Acid Biodegradation in Agricultural and Forest Soils

Anatoly Zyakun^{a b}, Seth Nii-Annang^c, Gabriele Franke^c, Thomas Fischer^d, Franz Buegger^e & Oliver Dilly^{c f}

^a Skryabin Institute of Biochemistry and Physiology of Microorganisms, Russian Academy of Sciences, Pushchino, Russia

^b Pushchino State University, Pushchino, Russia

^c Chair of Soil Protection and Recultivation, Brandenburg University of Technology, Cottbus, Germany

^d Central Analytical Laboratory, Brandenburg University of Technology, Cottbus, Germany

^e Institute for Soil Ecology, Helmholtz Center, Munich German Research Center for Environmental Health, Neuherberg, Germany

^f School of Integrated Climate System Sciences, University of Hamburg, Hamburg, Germany

Version of record first published: 19 Aug 2011

To cite this article: Anatoly Zyakun, Seth Nii-Annang, Gabriele Franke, Thomas Fischer, Franz Buegger & Oliver Dilly (2011): Microbial Activity and $^{13}\text{C}/^{12}\text{C}$ Ratio as Evidence of N-Hexadecane and N-Hexadecanoic Acid Biodegradation in Agricultural and Forest Soils, *Geomicrobiology Journal*, 28:7, 632-647

To link to this article: <http://dx.doi.org/10.1080/01490451.2010.489922>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.tandfonline.com/page/terms-and-conditions>

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae, and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand, or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

Microbial Activity and $^{13}\text{C}/^{12}\text{C}$ Ratio as Evidence of N-Hexadecane and N-Hexadecanoic Acid Biodegradation in Agricultural and Forest Soils

Anatoly Zyakun,^{1,2} Seth Nii-Annang,³ Gabriele Franke,³ Thomas Fischer,⁴ Franz Buegger,⁵ and Oliver Dilly³

¹Skryabin Institute of Biochemistry and Physiology of Microorganisms, Russian Academy of Sciences, Pushchino, Russia

²Pushchino State University, Pushchino, Russia

³Chair of Soil Protection and Recultivation, Brandenburg University of Technology, Cottbus, Germany

⁴Central Analytical Laboratory, Brandenburg University of Technology, Cottbus, Germany

⁵Institute for Soil Ecology, Helmholtz Center, Munich German Research Center for Environmental Health, Neuherberg, Germany

The dynamics of microbial degradation of exogenous contaminants, n-hexadecane and its primary microbial oxidized metabolite, n-hexadecanoic (palmitic) acid, was studied for topsoils, under agricultural management and beech forest on the basis the changes in O_2 uptake, CO_2 evolution and its associated microbial and non-microbial carbon isotopic signature, the respiratory quotient (RQ) and the priming effect (PE) of substrates. Soil microbial communities in agricultural soil responded to the n-hexadecane addition more rapidly compared to those of forest soil, with lag-periods of about 23 ± 10 and 68 ± 13 hours, respectively. Insignificant difference in the lag-period duration was detected for agricultural ($t_{\text{lag}} = 30 \pm 13$ h) and forest ($t_{\text{lag}} = 30 \pm 14$ h) soils treated with n-hexadecanoic (palmitic) acid. These results demonstrate that the soil microbiota differed in metabolic activities for using n-hexadecane as a reductive hydrocarbon and n-hexadecanoic acid as a partly oxidized hydrocarbon. The corresponding $\delta^{13}\text{C}$ of respired CO_2 after the addition of the hydrocarbon contaminants to soils indicates a shift in microbial activity towards the consumption of exogenous substrates with a more complete degradation of n-hexadecane in the agricultural soil, for which some initial contents of hydrocarbons are inherent. It was reflected in the carbon isotope signature of microbial biomass. It is supposed that the observed deviation of RQ from theoretically calculated value under microbial substrate mineralization is determined by difference in the time (Δt_i) of registration of CO_2 production and O_2 consumption. Positive priming effect (PE) of n-hexadecane and negative PE of n-hexadecanoic (palmitic) acid were detected in agricultural

and forest soils. It is suggested that positive PE of n-hexadecane is conditioned by the induction of microbial enzymes that perform hydroxylation/oxygenation of stable SOM compounds mineralized by soil microbiota to CO_2 . The microbial metabolism coupled with oxidative decarboxylation of n-hexadecanoic acid is considered as one of the most probable causes of the revealed negative PE value.

Keywords hydrocarbon contamination, microbial activity, microbial biomass, priming effect, oxygen consumption and carbon dioxide evolution, respiratory quotient, respired CO_2 -C isotopic signature

INTRODUCTION

The complex petroleum hydrocarbon mixtures, which include crude oil, diesel fuel, and creosote comprising various concentrations of linear and branched alkanes, cycloalkanes, phenolics, aromatics, and polycyclic aromatic hydrocarbons, may continue to be major environmental pollutants, since they remain important inputs in industrial and domestic activities, and together with their by-products, may eventually end up in the different compartments of the biosphere. The soil, although is an important sink for a wide range of substances, pollutant load exceeding certain threshold has the potential of impacting negatively on the capacity of the soil to perform its ecosystem functions with repercussions on sustainability issues such as plant growth (Adam and Duncan 2002, 2003; Palmroth et al. 2005). Thus, it is becoming increasingly evident that the fate of anthropogenic hydrocarbons pollutants entering the soil system requires efficient monitoring and control.

Soil microbial communities are able to adjust to unfavourable conditions and to use a broad spectrum of substrates (Jobson et al. 1974; Nikitina et al. 2003). They have unique metabolic

Received 22 December 2009; accepted 2 April 2010.

Current affiliation for Oliver Dilly: School of Integrated Climate System Sciences, University of Hamburg, Hamburg, Germany

Address correspondence to Anatoly Zyakun, Skryabin Institute of Biochemistry and Physiology of Microorganisms RAS, 142290, Pushchino, Moscow Region, Russia. E-mail: zyakun@ibpm.pushchino.ru

systems that allow them to utilise both natural and anthropogenic substances as a source of energy and tissue constituents. These unique characteristics make the microbiota a useful tool in monitoring and remediation processes. Bioremediation of soil contaminated with petroleum hydrocarbons has been established as an efficient, economic, versatile, and environmentally sound treatment (van Hamme et al. 2003). Several reports have already focused on the composition of natural microbial populations contributing to biotransformation and biodegradation processes in different environments polluted with hydrocarbons (Juck et al. 2000; Hamamura et al. 2008; Marques et al. 2008).

The bioremediation potential of microbial communities in soil polluted with oil hydrocarbons depends on their ability to adapt to new environmental conditions (Mishra et al. 2001; Kaplan and Kitts 2004). Investigations into how bioremediation influences the response of a soil microbial community, in terms of activity, biomass, and diversity, are presented in a series of publications (Jobson et al. 1974; Margesin and Schinner 2001; Zucchi et al. 2003; Hamamura et al. 2006; Margesin et al. 2007).

The most widely used bioremediation procedure is biostimulation of indigenous microorganisms by the addition of nutrients, because the input of large quantities of carbon sources (i.e., hydrocarbon contamination) results in rapid depletion of the available pools of major inorganic nutrients such as nitrogen and phosphorus (Alexander 1999; Dilly 1999; Margesin et al. 2007). In this case, the methods of monitoring and characterization of hydrocarbon degrading activity of soil microbiota are of special interest (Margesin and Schinner 2005; Abbassi and Shquirat 2008; Pleshakova et al. 2008).

To determine the pool of soil microorganisms utilizing oil hydrocarbons and to estimate microbial activity, n-hexadecane ($C_{16}H_{34}$) labeled with ^{13}C - or ^{14}C -isotope was used as an analogue of aliphatic hydrocarbon constituents of the complex petroleum hydrocarbon mixtures including crude oil, diesel fuel, creosote, etc. (Hamamura et al. 2006; Marchant et al. 2006; Dashti et al. 2008; Hassan and Sorial 2008). The initial stage of aliphatic hydrocarbons oxidation is carried out by specific microbial enzymes (alkane hydroxylase or oxygenase), which occur only in some microorganisms (Smits et al. 1999; Marchant et al. 2006).

Oxygen limitation that may occur under soil conditions can induce changes in the cytochrome levels (Asperger et al. 1986) and microorganisms are able to produce semi-oxidized metabolites. The n-hexadecanoic (palmitic) acid ($C_{16}H_{32}O_2$) may be one of the primary metabolites produced during microbial degradation of n-hexadecane. In the case of n-hexadecanoic acid, the oxidized end of the molecule is used as the initial site for β -oxidation process, whereas the microbial oxidation of saturated hydrocarbons (n-hexadecane) can not be initiated as easily as it does for n-hexadecanoic acid. (Metzler 1977; Van Hamme et al. 2003).

Comparing hydrocarbon consumption by soil microbiota in pristine and oily desert soils showed that alkane and n-hexadecane are essentially used by bacteria and palmitic acid as

a semi-oxidized hydrocarbon is utilized by fungi as additional carbon and energy sources (Dashti et al. 2008). Furthermore, the total content of soil microbial biomass (SMB) in arable soils was lower compared to forest and grassland sites (Dyckmans et al. 2003; Ananyeva et al. 2008). It was suggested that SMB depletion in arable soil compared to forest soil might be due to the decrease of the fungal biomass fraction in soil biomass.

Few studies in this context demonstrate that fungal spores dominated the total fungal biomass in arable soils and the major fraction of SMB was bacterial biomass (Stahl et al. 1999; Baath and Anderson 2002; Susyan et al. 2005; Susyan et al. 2011). The initial degradation of n-hexadecane compared to n-hexadecanoic acid would be expectedly more active in arable soils than in forest soil. Thus, the kinetic studies of microbial consumption of n-hexadecane and n-hexadecanoic acid as test substrates may be of profound importance for investigation of microbial diversity and oil degrading potential in soils.

Hydrocarbon biodegradation and transformation in soils can be monitored by estimating the concentration of pollutant and the formation of respective metabolites. One of the most ubiquitous and universal metabolites is carbon dioxide (CO_2), since respiration is by far the prominent pathway of biologically processed carbon. Soil microbiota can be characterized by means of the substrate-induced respiration (SIR). One of the indices of its activity is the priming effect (PE) of introduced exogenous substrate, which is easily accessible for microorganisms (e.g., glucose, cellulose, amino acids, etc.) (Fontaine et al., 2004; Zyakun and Dilly 2005; Blagodatskaya and Kuzyakov 2008).

The role of exogenous substrate in soil is assessed by the extent of activation of soil microbiota for utilization of native soil organic matter (SOM) and may be represented by the following three indices: (a) positive PE shows that exogenous substrate introduction concurrent with its mineralization increases SOM mineralization to a rate exceeding the previous rate; (b) zero PE shows that CO_2 is produced additionally only as a result of microbial mineralization of introduced substrate without changing the existing rate of SOM mineralization; and (c) negative PE values show that exogenous substrate introduction decreases SOM mineralization rate and CO_2 production is determined mainly by mineralization of the substrate.

Obviously, imprecise determination of PE by CO_2 production calls for an exogenous substrate different from SOM in carbon isotopes (Zyakun et al. 2003; Fontaine et al., 2004; Dilly and Zyakun 2008). PE determination only by the difference of CO_2 production rate before and after substrate introduction into soil suffers from the known uncertainty of CO_2 sources and does not allow distinguishing between the so-called "real" and "apparent" PE. (Blagodatskaya et al. 2007; Blagodatskaya and Kuzyakov 2008). Since complete oxidation of hydrocarbons demands high amounts of oxygen, the respiratory quotient (RQ) determined by the ratio of molar quantities of CO_2 produced and O_2 consumed is used as an indicator of microbial activity in soil (Dilly 2001; 2003).

The objectives of this study were (i) to examine the microbial response to n-hexadecane and n-hexadecanoic acid added to soils, (ii) to assess the importance of soil type (agricultural and forest) as determinant for the specific activity of hydrocarbon-degrading microbial population, and (iii) to compare reduced (n-hexadecane) and semi-oxidized (n-hexadecanoic acid) hydrocarbons added to native soils and the kinetics of microbial mineralization of hydrocarbons. The results will supposedly be a basis for estimation of the hydrocarbon bioremediation potential of impacted soils.

The metabolic activities of soil microbial community with respect to substrates (n-hexadecane and n-hexadecanoic acid) was determined from CO₂ evolution rates, the respiratory quotient, and the ¹³C-CO₂ isotope signature. The level of influence of exogenous hydrocarbons on SOM microbial mineralization was assessed by PE value and direction (positive, negative, or zero value). Nitrogen was added to avoid its limitation (Volke-Sepúlveda et al. 2006). After monitoring the respiratory characteristics, residual hydrocarbons and the changes in the carbon isotope signature of microbial biomass were estimated.

MATERIALS AND METHODS

Soil characteristics and sampling. Two soils (A horizon) were sampled at the Bornhöved Lake District in the northern Germany (54°06'N, 10°14'E). The first classified as eutricambic Arenosol (IUSS Working Group WRB, 2006) is an agricultural soil under *Lolium* perennial grass since 2003, after regular crop rotation and has a pH [H₂O] of 6.4, organic C and total N contents of 14.4 mg C and 1.4 mg N g⁻¹ soil, respectively. The second, which is a dystic-cambic Arenosol according to IUSS Working Group WRB (2006), developed under beech forest under a matured stratified litter layer, and is characterized by pH [H₂O] of 4.1, an organic C of 33.8 mg C g⁻¹ soil, and a total N content of 2.4 mg N g⁻¹ soil.

Three field replicates, each composed of more than 10 subsamples bulked together were collected at a distance of approximately 20 m from each other at both the agricultural and forest sites. Plant materials were carefully removed, and the soils were sieved to 2 mm and stored at 4°C for no longer than one month before the analysis. The soil moisture content was adjusted before incubation experiments and it was varied in the range of 40 to 70% of the water-holding capacity during incubations.

The microbial activities were estimated using three replicates of agricultural and forest soils which were amended with n-hexadecane and n-hexadecanoic acid, respectively. There were twelve independent soil samples in experimental measurements.

Preparation of soil samples and measurements. The hydrophobic compound was applied as follows: n-hexadecane or n-hexadecanoic (palmitic) acid (200 µl of n-hexadecane corresponding to 156 mg and 177 mg of palmitic acid) were added to 1 g of dried and dispersed soil together with 20 mg NH₄Cl (5 mg N) and then 1 g of the soil was mixed with fresh moist soil equivalent to 100 g of dry material. The final substrate concentration was 1.3 mg C g⁻¹ soil with a C/N-ratio of 25. Ni-

trogen was added to favour microbial metabolism (Dilly 1999) and n-hexadecane degradation (Volke-Sepúlveda et al. 2006). Nitrogen was applied as ammonium chloride since nitrate may be denitrified (Chayabutra and Ju 2000).

Moist soil corresponding to 100 g dry weight was pre-incubated for 7 to 10 days (BAS, basal respiration) prior to the addition of n-hexadecane or n-hexadecanoic acid with carbon isotopic signatures different from that of SOM to determine CO₂ sources in the experiments on substrate-induced respiration (SIR). The oxygen uptake by soil microbiota and total volume of CO₂ produced during the experiments was determined by respiratory and titration, respectively (Zyakun and Dilly 2005; Dilly and Zyakun 2008).

Five grams of soil were extracted with 20 mL of 0.01 M CaCl₂ solution with and without previous chloroform fumigation. The fraction after fumigation includes also the fraction immobilized by the microbial biomass (Vance et al. 1987) and will hereafter be indicated as "microbial." The hydrocarbon substances present in the native soils were quantified as follows: 5 g of soil was extracted with 10 mL of acetone-hexane (1:1) mixture for 2 hours on a shaker; the flask was filled up with 50 mL of deionised water. n-Hexadecane was determined in the hexane phase by gas chromatography with a flame ionization detector (Column: BPX5 0.25 µm 60 m × 0.25 mm ID, Column flow: 2 mL He min⁻¹, temperature programme: 100°C, 2 min hold, 20 K/min to 240°C, 12 min hold, 30 K/min to 300°C, 17 min hold, Injection: 1 µL splitless 300°C, Detector: 300°C). The n-hexadecanoic (palmitic) acid in the soil was quantified as follows: 40 µL of the hexane phase and 40 µL of the internal standard (0.1 µg mL⁻¹ heptadecanoic acid in hexane), was evaporated to dryness and dissolved in 40 µL chloroform plus 40 µL TMSH solution (0.25 M in methanol, Fluka 92732) and analysed as described above. A FAME-mixture (Supelco 47885-U) was used as the external standard.

Carbon isotopic analysis. The ratios of ¹³C/¹²C abundances of microbial respired CO₂-C in BaCO₃ were determined using the Thermo Finnigan Delta V Advantage Isotope Ratio Mass Spectrometer (IRMS) coupled with the Element Analyzer Vario EL (Elementar Instruments). The temperatures in the oxidation and reduction oven were 1150 and 850°C, respectively. Approximately 0.09 mg C, corresponding to 1.8 mg of the dried BaCO₃, mixed with about 0.5 mg wolframite for effective combustion, was used for the analysis of the ¹³C/¹²C ratio. The ratio between the peak intensities at m/z 45 (¹³C¹⁶O₂) and 44 (¹²C¹⁶O₂) was a quantitative characteristic of the ¹³C and ¹²C contents. By convention, the carbon isotope characteristics are expressed by equation [1]:

$$\delta^{13}\text{C} = (\text{R}_{\text{sa}}/\text{R}_{\text{st}} - 1)1000\text{‰}, \quad [1]$$

where $\text{R}_{\text{sa}} = ([^{13}\text{C}]/[^{12}\text{C}])_{\text{sa}}$ and $\text{R}_{\text{st}} = ([^{13}\text{C}]/[^{12}\text{C}])_{\text{st}}$ are isotope abundance ratios of a sample and the international Pee Dee Belemnite (PDB) standard (Craig 1957), respectively.

Mass isotopic balance for n-hexadecane. The test-substrate of n-hexadecane with $\delta^{13}\text{C}$ value of $+47.8 \pm 0.2\text{‰}$ was

prepared by mixing 30 mg of n-hexadecane-1,2 $^{13}\text{C}_2$ (where 99 Atom% ^{13}C were in 1 and 2 sites) and the remainder of 14 C atoms have natural ^{13}C isotope content (i.e., $1.0960 (\pm 0.0002)$ Atom% ^{13}C) (Sigma 485799) with 5000 mg of natural n-hexadecane (HEX) ($\delta^{13}\text{C} = -28 \pm 0.2\text{‰}$ relative to the PDB standard or $R_{\text{HEX}} = 0.010922(\pm 2 \cdot 10^{-6})$ and $R_{\text{st}} = 0.0112372$ (Craig 1957). The carbon isotope abundance ratio of n-hexadecane-1,2 $^{13}\text{C}_2$, 99 Atom% (HEX*) was equal to $R_{\text{HEX}}^* = 0.153855 (\pm 5 \cdot 10^{-6})$ and was calculated as a ratio of $[^{13}\text{C}]_{\text{HEX}} = 14 \cdot 0.01096 (\pm 2 \cdot 10^{-5}) + 2 \cdot 0.99 = 2.13344 (\pm 3 \cdot 10^{-4})$ to $[^{12}\text{C}]_{\text{HEX}} = 14 \cdot 0.98904 (\pm 2 \cdot 10^{-5}) + 2 \cdot 0.01 = 13.86656 (\pm 3 \cdot 10^{-4})$ (note that $[^{13}\text{C}]^* + [^{12}\text{C}]^* = 1$). As is shown by mass isotopic balance, the mixture (MIX) of 30 mg of HEX* with $R_{\text{HEX}}^* = 0.15385 (\pm 2 \cdot 10^{-5})$ and 5000 mg of natural n-hexadecane with $R_{\text{HEX}} = 0.010922 (\pm 2 \cdot 10^{-6})$ is characterized by the average carbon isotope abundance ratio equal to $R_{\text{MIX}} (0.011774 (\pm 2 \cdot 10^{-6})$ or $\delta^{13}\text{C}_{\text{MIX}} = +47.8 \pm 0.2\text{‰}$ relative to the PDB standard.

When n-hexadecane was used as aliphatic hydrocarbon, it was assumed that the initial and subsequent stages of microbial mineralization of this substance to CO_2 are independent of the presence and position of ^{13}C atoms in the molecule. The point is that, the pathways of aliphatic hydrocarbon degradation are represented at the initial stages of this process as molecules containing alcohol – aldehyde – acid groups (Delasse and Nyns 1974; Van Hamme et al. 2003). Oxidation of the mono-alcohol group into an aldehyde group (Tauchert et al. 1978) and a carboxyl group is accompanied by appearance of the key metabolite: n-hexadecanoic (palmitic) acid.

Then, Acetyl CoA, which is formed as a result of step by step β -rupture of the C-COO $^-$ bond in n-hexadecanoic acid, is metabolized into CO_2 molecules (Kawamoto et al. 1978). In this case, the characteristics of carbon isotope content ($\delta^{13}\text{C}$) of CO_2 pool produced during microbial n-hexadecane mineralization will inherit the carbon isotopic characteristics of the n-hexadecane pool. It means that the $\delta^{13}\text{C}$ value of CO_2 pool produced during microbial degradation of n-hexadecane (the average value of $\delta^{13}\text{C}_{\text{HEX}} = +47.8 \pm 0.2\text{‰}$) is rather different from CO_2 pool resulting from soil organic matter (SOM) mineralization ($\delta^{13}\text{C}_{\text{SOM}}$ is equal to $-26.8 \pm 0.2\text{‰}$ and $-26.2 \pm 0.2\text{‰}$ for agricultural and forest soils, respectively). These values of $\delta^{13}\text{C}$ can be used as specific parameters for mass isotopic balance calculations and PE substrate estimations.

Thus, after addition of the n-hexadecane as a test substrate to soil, the mass isotopic balance for CO_2 evolved during microbial mineralization of SOM and the substrate (SUB) was calculated using equation [2]:

$$\delta^{13}\text{C}_{\text{SOM}} \cdot Q_{\text{SOM}} + \delta^{13}\text{C}_{\text{SUB}} \cdot Q_{\text{SUB}} = \delta^{13}\text{C}_{\text{MIX}} \cdot (Q_{\text{SOM}} + Q_{\text{SUB}}), \quad [2]$$

where $\delta^{13}\text{C}_{\text{SOM}}$ and $\delta^{13}\text{C}_{\text{MIX}}$ are isotopic characteristics of ^{13}C content in CO_2 pool before and after substrate addition to the

soil; $\delta^{13}\text{C}_{\text{SUB}}$ is the isotopic characteristic of ^{13}C content in CO_2 produced during microbial mineralization of the substrate; Q_{SOM} and Q_{SUB} are CO_2 quantities resulted from microbial mineralization of SOM and added test substrate in the soil samples, respectively. Using the mass isotopic balance, the CO_2 fractions of SOM (F) and of the test substrate (1-F) in the total quantity of CO_2 formed after exogenous substrate addition to the soil were estimated by equation [3]:

$$F_{\text{SOM}} = (\delta^{13}\text{C}_{\text{MIX}} - \delta^{13}\text{C}_{\text{SUB}}) / (\delta^{13}\text{C}_{\text{SOM}} - \delta^{13}\text{C}_{\text{SUB}}) \quad [3]$$

Here by definition $F_{\text{SOM}} = Q_{\text{SOM}} / (Q_{\text{SOM}} + Q_{\text{SUB}})$ and $1 - F_{\text{SOM}} = Q_{\text{SUB}} / (Q_{\text{SOM}} + Q_{\text{SUB}})$

Mass isotopic balance for n-hexadecanoic acid. The test substrate of n-hexadecanoic acid with $\delta^{13}\text{C}$ value of $+29.6 (\pm 0.2)\text{‰}$ was prepared as follows: the 18.5 mg of ^{13}C labeled n-hexadecanoic (palmitic) acid (*PA) containing one ^{13}C in the carboxylic group $^{13}\text{COOH}$ about 99 Atom% ^{13}C and the rest of 15 C atoms with the $^{13}\text{C}/^{12}\text{C}$ ratio of $R_{\text{C15}} = 0.010960 (\pm 2 \cdot 10^{-6})$ was mixed with 2000 mg of natural n-hexadecanoic acid (PA) ($\delta^{13}\text{C}_{\text{PA}} = -26.0 (\pm 0.2)\text{‰}$ relative to PDB standard or $R_{\text{PA}} = 0.010945 (\pm 2 \cdot 10^{-6})$). The average carbon isotope abundance ratio of n-hexadecanoic acid- $^{13}\text{C}_1$ labeled in ^{13}C was equal to $R_{\text{PA}}^* = 0.07776 (\pm 2 \cdot 10^{-5})$ and calculated as a ratio of $[^{13}\text{C}]^* = 0.99 + 15 \cdot 0.01096 (\pm 2 \cdot 10^{-6}) = 1.1544 (\pm 3 \cdot 10^{-5})$ to $[^{12}\text{C}]^* = 15 \cdot 0.98904 (\pm 2 \cdot 10^{-6}) + 0.01 = 14.8456 (\pm 3 \cdot 10^{-5})$ (note that $[^{13}\text{C}]^* + [^{12}\text{C}]^* = 1$). The test substrate prepared as a mixture (MIX) of 18.5 mg PA* with $R_{\text{PA}}^* = 0.07776 (\pm 2 \cdot 10^{-5})$ and 2000 mg of natural n-hexadecanoic acid with $R_{\text{PA}} = 0.01096 (\pm 2 \cdot 10^{-6})$ was characterized by the average carbon isotope abundance ratio equal to $R_{\text{PA(MIX)}} = 0.011570 (\pm 2 \cdot 10^{-6})$ or $\delta^{13}\text{C}_{\text{PA(MIX)}} = +29.6 (\pm 0.2)\text{‰}$ relative to the PDB standard.

It means that the $\delta^{13}\text{C}$ value of CO_2 pool produced during microbial degradation of the test substrate (n-hexadecanoic acid pool with the average value of $\delta^{13}\text{C}_{\text{HEX}} = +29.6 (\pm 0.2)\text{‰}$) is rather different from CO_2 pool resulting from soil organic matter (SOM) mineralization ($\delta^{13}\text{C}_{\text{SOM}}$ is equal to $-26.6 \pm 0.3\text{‰}$ and $-26.2 \pm 0.4\text{‰}$ for agricultural and forest soils, respectively). Thus, the values of $\delta^{13}\text{C}$ (SOM) and $\delta^{13}\text{C}$ (PA) can be used as specific parameters for mass isotopic balance calculations and PE estimations (Eqs. 2 and 3).

Furthermore, it should be remembered that the test substrate PA represents the pool of PA molecules with non-homogeneity ^{13}C distribution in the carbon sites of molecule. This is due to fact that the PA test substrate is a mixture of 18.5 mg of PA- $^{13}\text{C}_1$ (99% ^{13}C) and 2000 mg of PA- $^{13}\text{C}_1$ (1.095% ^{13}C) and the probability to detect the ^{13}C isotope in the C_1 -site (-COOH) will be higher than in the C_{2-16} -sites of the test substrate pool. According to the mass isotopic balance, we determined that the content of ^{13}C in the sate of C_1 (carboxyl group) of the test substrate pool will characterized by $R(\text{C}_1) = [^{13}\text{C}/^{12}\text{C}] = 0.01992$, while the rest part of carbon atoms of $\text{C}_2\text{-C}_{16}$ -sates will be $R(\text{C}_2\text{-C}_{16}) = 0.01095$.

Respiratory quotient (RQ). To test for microbial activity, the respiratory quotient (RQ), defined as a molar ratio of CO₂ evolution per O₂ consumption during certain time intervals (Δt_i) of microbial growth in soils, was expressed by $RQ_i = [\text{mol CO}_2]/[\text{mol O}_2]$ (Dilly 2001). The average value of RQ_{av} for microbial activity throughout the total Δt -period defined as $\Delta t = \Sigma(\Delta t_i)$ was calculated by the equation

$$RQ_{av} = \Sigma(RQ_i \cdot \Delta t_i) / \Sigma(\Delta t_i) \quad [4]$$

According to the schemes of chemical reactions, the theoretical estimation of RQ values for quantitative aerobic oxidation of n-hexadecane (C₁₆H₃₄) and n-hexadecanoic acid (C₁₆H₃₂O₂) to CO₂ are equivalent to 0.65 mol CO₂ mol⁻¹O₂ and 0.70 mol CO₂ mol⁻¹O₂, respectively.

The Kinetics of Substrate-Induced Respiration

Specific CO₂ evolution rates (μ) of soil microorganisms after n-hexadecane or n-hexadecanoic acid addition to soil were estimated from the kinetic analysis of substrate-induced respiration (CO₂(t)) by fitting the parameters of equation [5]:

$$\text{CO}_2(t) = K + r \cdot \exp(\mu \cdot t) \quad [5]$$

where K is the initial respiration rate uncoupled from ATP production, r is the initial rate of respiration by the growing fraction of the soil microbiota coupled with ATP generation and cell growth, and t is time (Panikov and Sizova 1996; Stenström et al. 1998; Blagodatsky et al. 2000). The lag period duration (t_{lag}) was determined as the time interval between substrate addition and the moment when the increasing rate of microbial growth-related respiration $r \cdot \exp(\mu \cdot t)$ became as high as the rate of respiration uncoupled from ATP generation. According to the theory of microbial growth kinetic (Panikov 1995; Blagodatskaya et al. 2009), the lag period was calculated by using the parameters of approximated curve of respiration rate of microorganisms with [6]:

$$t_{lag} = \ln(K/r) / \mu \quad [6]$$

Cumulative CO₂ resulted from hydrocarbon mineralization. Cumulative CO₂ produced during the microbial substrate oxidation was calculated as follows. The ΔQ_i quantity of CO₂ evolved during the Δt_i -time interval ($i = 1, 2, \dots, n$) was estimated as $\Delta Q_i = \Delta t_i \cdot v_i$, where the v_i -value is the rate of CO₂ evolved during the time interval Δt_i . Using $\delta^{13}\text{C}_{\text{soil}}$, $\delta^{13}\text{C}_{\text{Subst}}$ and $\delta^{13}\text{C}_{\text{CO}_2(\text{mix})(i)}$, the fraction of CO₂ resulting from the exogenous substrate (n-hexadecane or n-hexadecanoic acid) oxidation during Δt_i can be calculated as [7]:

$$\Delta Q_{\text{Subst}(i)} = (1 - F_i) \cdot \Delta Q_i \quad [7]$$

where F_i value can be estimated using equation [3]. The cumulative CO₂ quantity ($Q_{\text{Subst}(\text{CO}_2)}$) resulting from microbial

oxidation of the substrates in soils was presented by [8]:

$$Q_{\text{Subst}(\text{CO}_2)} = \Sigma \Delta Q_{\text{Subst}(i)} \quad [8]$$

where i varied from 1 to n.

Calculation of priming effects. The addition of exogenous test substrates to soil was accompanied by the change in soil microbiota activity: the rate of CO₂ production initially increased as a result of substrate and probably SOM mineralization and then, on depletion of the substrate, gradually decreased. The amount of CO₂ evolved was divided by means of mass isotopic balance into two fractions: from the substrates (n-hexadecane or n-hexadecanoic acid) and from SOM mineralization. Thus, the difference between CO₂ evolved from SOM mineralization in n-hexadecane and n-hexadecanoic acid amended soil (C_{SOM}) and in the control soil (C_{SOM}) relative to the control (in percentage) was used to estimate the magnitude of the priming effect (PE) induced by n-hexadecane or n-hexadecanoic acid (denoted as SUB). The PE value was determined in two notations as *kinetic* PE(Δt_i) [9] and *total* PE(*total*) calculated as a weighted average value for the whole period of observation using equation. [10].

$$\text{PE}(\Delta t_i)[\%] = 100 \times (C_{\text{SOM}(i)} - C_{\text{SOM}(i)}) / C_{\text{SOM}(i)}, \quad [9]$$

$$\text{PE}(\text{total})[\%] = 100 \times \Sigma(\text{PE}(\Delta t_i) \cdot \Delta t_i) / \Sigma(\Delta t_i) \quad [10]$$

where $C_{\text{SOM}(i)} = F_i \times C_{(\text{SUB}+\text{SOM})i}$; $C_{(\text{SUB}+\text{SOM})i}$ is the total C evolved as CO₂ in the amended soil during Δt_i -time; and F_i is the share of CO₂-C resulting from the SOM in n-hexadecane and n-hexadecanoic acid amended soil in Δt_i -time, which was calculated by equation [3].

RESULTS

Basal respiration and hydrocarbon contents. The native hydrocarbon contents in agricultural and forest soils used in the experiments were measured by gas chromatography (Table 1). There are significant differences in the basal amounts of n-hexadecane (HEX) between agricultural and forest soils: n-hexadecane makes up 2.57% of total hydrocarbons in the agricultural soil (equal to 1.5 $\mu\text{g C-HEX g}^{-1}$ soil) but is absent in the forest soil. At the same time, n-hexadecanoic (palmitic) acid (PA) contents are close for both soils: about 7.02% and 7.61% of total hydrocarbons for agricultural and forest soils, respectively (C16:0, Table 1).

At the beginning of experiments the CO₂ evolution rates were not significantly different in agricultural and forest soils (Fig. 1, A, curve 1; D, curve 5; and Fig. 2, A, curve 1; D, curve 5). At the same time, the values of respiratory quotients (RQ) were rather different: 0.73 ± 0.05 and 0.87 ± 0.05 mol CO₂ mol⁻¹ O₂ for agricultural and forest soils, respectively (Fig. 1, C and F). The $\delta^{13}\text{C-CO}_2$ value during basal metabolism was $-26.88 \pm 0.08\text{‰}$ and $-26.15 \pm 0.18\text{‰}$ for agricultural and forest soil (Table 2, Time = 0) and was not significantly different from the carbon isotope signature of soil organic matter (SOM) (data

TABLE 1
The relative quantities of hydrocarbons and fatty acids in soils (in% of sum of total peak areas)

Number	Retardation time [min]	Peak name	Agricultural soil Relative area [%]	Forest soil Relative area [%]
1	8.67	C 10:0	1.23	—**
2	9.91	—*	1.02	1.18
3	10.68	—*	2.36	2.86
4	11.03	n-hexadecane	2.57	—**
5	11.83	—*	1.09	—**
6	12.19	—*	2.38	2.4
7	13.10	—*	1.86	3.69
8	14.03	C 16:1	2.01	2.37
9	14.19	C 16:0	7.02	7.61
10	14.54	—*	3.62	2.53
11	16.74	—*	1.88	1.72
12	16.95	C 18:2n6/C18:1n9	3.95	5.42
13	17.34	C 18:0	1.64	2.68
14	19.59	—*	0.99	1.22
15	22.20	C 20:0	1.18	1.33
16	22.68	—*	1.18	1.53
17	24.61	C 22:2	3.29	11.12
18	25.40	C 22:0	2.66	6.21
19	26.90	C 23:0	1.51	2.51
20	28.43	C 24:0	4.51	4.52

*Components could not be identified.

**No experimental peak was registered.

not shown) except that the $\delta^{13}\text{C}$ -CO₂ of agricultural soil was depleted by 0.7‰ relative to the bulk soil during the first 3 days.

Microbial consumption of n-hexadecane. The addition of the test substrate to soils at a rate of 6.77 μM n-hexadecane

per g soil or 1300 $\mu\text{g C g}^{-1}$ soil stimulated the CO₂ evolution rate in agricultural soil within 24 hours (Fig. 1, A, curve 1). In the forest soil (Fig. 1, D, curve 5), no significant increase of CO₂ evolution rate was observed during the same period of time.

TABLE 2
Carbon isotope indicators of metabolic CO₂ ($\delta^{13}\text{C}_{\text{CO}_2}$,‰) before and after n-hexadecane addition to soils and CO₂ fractions (F_{HEX}) produced during mineralization of n-hexadecane

Time [days]	Agricultural soil		Forest soil	
	$\delta^{13}\text{C}_{\text{CO}_2}$ [‰]	*F _{HEX}	$\delta^{13}\text{C}_{\text{CO}_2}$ [‰]	*F _{HEX}
0.00	−26.88 ± 0.08	0.000	−26.15 ± 0.18	0.000
1.00	−13.60 ± 2.70	0.178 ± 0.036	−21.40 ± 0.79	0.064 ± 0.024
1.75	18.07 ± 1.03	0.604 ± 0.014	−18.00	0.110 ± 0.020
2.74	14.20 ± 1.29	0.552 ± 0.017	−2.95 ± 2.48	0.315 ± 0.034
3.71	17.51 ± 1.18	0.596 ± 0.016	16.65 ± 4.18	0.580 ± 0.057
4.47	19.87 ± 0.90	0.628 ± 0.012	19.87 ± 0.90	0.624 ± 0.012
6.78	16.12 ± 1.58	0.577 ± 0.021	13.82 ± 5.56	0.542 ± 0.075
8.74	14.90 ± 2.42	0.561 ± 0.033	11.33 ± 3.30	0.508 ± 0.045
10.7	18.90 ± 1.00	0.615 ± 0.013	14.70 ± 1.00	0.554 ± 0.014
13.7	16.70 ± 1.50	0.585 ± 0.008	10.60 ± 1.40	0.498 ± 0.020

*F_{HEX} = ($\delta^{13}\text{C}_{\text{CO}_2} - \delta^{13}\text{C}_{\text{SOM}}$)/($\delta^{13}\text{C}_{\text{HEX}} - \delta^{13}\text{C}_{\text{SOM}}$) is the fraction of CO₂ produced during microbial mineralization of n-hexadecane, where $\delta^{13}\text{C}_{\text{HEX}} = +47.8\text{‰}$ and $\delta^{13}\text{C}_{\text{SOM}}$ for CO₂ evolved from agricultural and forest soil were equal to −26.88 (0.08)‰ and −26.15 (0.18)‰, respectively.

Value ± standard deviation.




FIG. 1. Carbon dioxide evolution rate and respiratory quotient (RQ) in experiments with n-hexadecane added to agricultural and forest soils: A—(1) total CO_2 rate after n-hexadecane addition to the agricultural soil; (2) rate of CO_2 from n-hexadecane mineralization; B—(3) rate of CO_2 from SOM mineralization after n-hexadecane addition; (4) rate of CO_2 from SOM (control); D—(5) total CO_2 rate after n-hexadecane addition to the forest soil; (6) rate of CO_2 from n-hexadecane mineralization; E—(7) rate of CO_2 from SOM mineralization after n-hexadecane addition; (8) rate of CO_2 from SOM mineralization (control); C and F—respiratory quotient values in agricultural and forest soils; bars extend to confidence limits at $P < 0.05$.

Parameterization of Equation [5] showed the response of growing microorganisms to the test substrate addition: with n-hexadecane added to the soils, the discrepancy of specific CO_2 production rates (μ) obtained for agricultural vs. forest soil was within the experiment error of analysis, but fraction of growing microorganisms (parameter r) was substantially higher in agricultural vs. forest soil (Table 3). Besides, a significant change in lag-period duration was revealed between agricultural and forest soil treated with n-hexadecane (Table 3). In 10–30 and in 45–65 hours, n-hexadecane introduced into agricultural and forest soils, respectively, caused an exponential

increase in the CO_2 emission rate indicating microbial growth after the lag-phase (Fig. 3).

In agricultural soil (Fig. 1, A, curve 1), two CO_2 respiration peaks were registered at $2.74 \mu\text{g CO}_2\text{-C g}^{-1}\text{ soil h}^{-1}$ and $2.20 \mu\text{g CO}_2\text{-C g}^{-1}\text{ soil h}^{-1}$ in 43 and 165 hours (or 1.8 and 6.9 days) from the moment of n-hexadecane addition, respectively. In forest soil (Fig. 1, D, curve 5), only one peak of the CO_2 emission rate was detected at $4.14 \mu\text{g CO}_2\text{-C g}^{-1}\text{ soil h}^{-1}$ in 120 hours (or 5 days) from the moment of n-hexadecane addition. On the basis of mass isotopic balance CO_2 fractions, produced during microbial mineralization of n-hexadecane (F_{HEX}), in agricultural and forest soil were calculated (Table 2).




FIG. 2. Carbon dioxide evolution rate and respiratory quotient (RQ) in experiments with n-hexadecanoic acid added to agricultural and forest soils: A – (1) total CO_2 rate after n-hexadecanoic acid addition to the agricultural soil; (2) rate of CO_2 from n-hexadecanoic acid mineralization; B – (3) rate of CO_2 from SOM (control); (4) rate of CO_2 from SOM mineralization after n-hexadecanoic acid addition; D – (5) total CO_2 rate after n-hexadecanoic acid addition to the forest soil; (6) rate of CO_2 from n-hexadecanoic acid mineralization; E – (7) rate of CO_2 from SOM mineralization (control); (8) rate of CO_2 from SOM mineralization after n-hexadecanoic acid addition; C and F – respiratory quotient values in agricultural and forest soils; bars extend to confidence limits at $P < 0.05$.

Using F_{HEX} values, the rates of total CO_2 production after addition of n-hexadecane to agricultural and forest soil were divided into two parts: production due to mineralization of n-hexadecane (Fig. 1, A, curve 2; D, curve 6) and due to mineralization of SOM in experiments (Fig. 1, B, curve 3; E, curve 7). For comparison the rates of CO_2 production prior to addition of n-hexadecane to the same soil (control) are given (Fig. 1, B, curve 4; E, curve 8). Simultaneously with the change of rates of CO_2 production alterations of respiratory coefficient

which reflected the ratio of CO_2 produced by soil microbiota to consumed oxygen herewith were registered (Fig. 1, C and F).

At the end of 17-day (414-hour) experiments with n-hexadecane, the total metabolic quantities of CO_2 about $631 \pm 42 \mu\text{g C-CO}_2 \text{ g}^{-1} \text{ soil}$ and about $587 \pm 55 \mu\text{g C-CO}_2 \text{ g}^{-1} \text{ soil}$ were evolved from agricultural and forest soils, respectively (Table 4, Q_{CO_2}). Based on the mass isotopic balance [Eq. 2], the quantities of CO_2 resulting from microbial mineralization of n-hexadecane were estimated at about $349 (\pm 26) \mu\text{g C-CO}_2$




FIG. 3. Kinetics of substrate-induced respiratory responses of microbial communities during periods of slow mineralization of n-hexadecane (left) and n-hexadecanoic (palmitic) acid (right) in the agricultural (1) and forest (2) soils: rates of microbial CO₂ production before substrate addition to soils are labeled as “-” value of time. h. (Experimental data are shown as symbols and model simulation as curves according to Equation 1 with parameters presented in Table 2).

g⁻¹ soil and 291(±32) µg C-CO₂ g⁻¹ soil in the experiments with agricultural and forest soils (Table 4, Q_{CO₂(SUB)}). At the same time, the total quantities of CO₂ derived from SOM mineralization after the n-hexadecane addition to agricultural and forest soils were 284(±26) µg C-CO₂ g⁻¹ soil and 296(±32) µg C-CO₂ g⁻¹ soil, respectively (Table 4, Q_{CO₂(SOM)}).

After the 17-day period, CO₂ production by microbiota from soil without the addition of exogenous substrates (control) was estimated at about 180(±25) µg C-CO₂ g⁻¹ soil and 196(±24) µg C-CO₂ g⁻¹ soil for agricultural and forest soil, respectively (Table 4, Q_{CO₂(contr)}). The results of calculated average PE are presented in Table 4: the positive priming effect of n-hexadecane (PE_{agr} = +57(±29)% and PE_{forest} = +51(±29)%) were estimated for agricultural and forest soil, respectively. Concurrently to the stimulation of respiratory response by n-hexadecane, the

kinetics of the positive priming production of CO₂ from SOM mineralization relative to native process was determined for agricultural soil and for forest soil (Fig. 1, B, curves 3 relative to curve 4; E, curves 7 relative to curve 8).

Microbial consumption of n-hexadecanoic acid. Experimentally, the addition of n-hexadecanoic (palmitic) acid to soils at a rate of 6.77 µM n-hexadecanoic acid g⁻¹ soil or 1300 µg C g⁻¹ soil stimulated CO₂ evolution in agricultural soil within 24 hours, while in forest soil CO₂ evolution was delayed within this period (Fig. 2, A, curve 1; and D, curve 5). Besides, two CO₂ respiration peaks were registered in agricultural soil at 1.8 µg CO₂-C g⁻¹ soil h⁻¹ and 1.4 µg CO₂-C g⁻¹ soil h⁻¹ in 24 and 120 h (or 1.0 and 5.0 days) after n-hexadecanoic acid addition (Fig. 2, A, curve 1), respectively. In forest soil, only one peak of CO₂ emission rate was detected at 4 ± 1 µg CO₂-C g⁻¹

TABLE 3
Parameters of the equations CO₂(t) = K+r·exp(µ·t) and t_{lag} = ln(K/r)/µ characterized the rates of n-hexadecane and n-hexadecanoic acid-induced CO₂ respiration and microbial growing in the agricultural and forest soils

Treatment	K	r	μ	t_{lag}
	[$\mu\text{g CO}_2 \text{ g}^{-1} \text{ soil h}^{-1}$]			[hour]
<i>n-hexadecane</i>				
Agricultural soil	0.4263 ± 0.0513	0.0289 ± 0.0160	0.1055 ± 0.0126	23 ± 10
Forest soil	0.5910 ± 0.0282	0.0009 ± 0.0005	0.0915 ± 0.0069	68 ± 13
<i>n-hexadecanoic acid</i>				
Agricultural soil	0.4726 ± 0.0360	0.0205 ± 0.0126	0.0953 ± 0.0132	30 ± 13
Forest soil	0.3978 ± 0.0830	0.0975 ± 0.0257	0.0427 ± 0.0028	30 ± 14

Value ± standard deviation.

TABLE 4

Microbial mineralization of substrates (SUB) (n-hexadecane and n-hexadecanoic (palmitic) acid) added to the agricultural and forest soils and soil organic matter (SOM) and priming effects of substrates

Parameters	n-hexadecane		n-hexadecanoic acid	
	Agricultural soil	Forest soil	Agricultural soil	Forest soil
Time (t), [hour]	414	414	385	385
Q_{CO_2} , $\mu\text{gC-CO}_2 \text{ g}^{-1} \text{ soil}$	631 ± 42	587 ± 55	387 ± 30	548 ± 35
$\delta^{13}\text{C}_{\text{av}}$, ‰	$+14.4 \pm 0.5$	$+10.5 \pm 0.5$	$+12.7 \pm 0.3$	$+15.7 \pm 0.5$
$X_{\text{(SUB)}}$	0.553 ± 0.005	0.496 ± 0.008	0.697 ± 0.013	0.751 ± 0.016
$Q_{CO_2(\text{SUB})}$, $\mu\text{gC-CO}_2 \text{ g}^{-1} \text{ soil}$	349 ± 26	291 ± 32	270 ± 26	412 ± 35
$Q_{CO_2(\text{SOM})}$, $\mu\text{gC-CO}_2 \text{ g}^{-1} \text{ soil}$	284 ± 26	296 ± 32	117 ± 26	136 ± 35
$Q_{CO_2(\text{contr})}$, $\mu\text{gC-CO}_2 \text{ g}^{-1} \text{ soil}$	180 ± 25	196 ± 24	184 ± 25	200 ± 27
PE _{average} , %	$+57(\pm 29)$	$+51(\pm 29)$	$-36(\pm 28)$	$-32(\pm 30)$

$t = \Sigma(\Delta t_i)$, (hour) is the time interval of measurements.

$Q_{CO_2} = \Sigma(\Delta t_i \cdot v_i)$, ($\mu\text{g C-CO}_2 \text{ g}^{-1} \text{ soil}$) is a total CO_2 produced during the time t where v_i ($\mu\text{g C-CO}_2 \text{ g}^{-1} \text{ soil h}^{-1}$) is a rate of CO_2 evolved from soil in the time interval of Δt_i .

$\delta^{13}\text{C}_{\text{av}} = \Sigma(\delta^{13}\text{C}_i \cdot \Delta t_i \cdot v_i) / Q_{CO_2}$ (‰) is an average value.

$X_{\text{(SUB)}} = (\delta^{13}\text{C}_{\text{av}} - \delta^{13}\text{C}_{t=0}) / (\delta^{13}\text{C}_{\text{substr}} - \delta^{13}\text{C}_{t=0})$ is a carbon fraction derived from substrates (n-hexadecane or n-hexadecanoic acid) in the total CO_2 respired and $\delta^{13}\text{C}_{t=0}$ is the carbon isotopic characteristic of CO_2 before substrates addition to soil.

$Q_{CO_2(\text{SOM})} = (1 - X) \cdot Q_{CO_2}$, ($\mu\text{g C-CO}_2 \text{ g}^{-1} \text{ soil}$) is a quantity of CO_2 evolved from soil amended with n-hexadecane or n-hexadecanoic acid.

$Q_{CO_2(\text{contr})} = t \cdot v_{\text{contr}}$, ($\mu\text{g C-CO}_2 \text{ g}^{-1} \text{ soil}$) is a total CO_2 produced during time of experiment, where v_{contr} ($\mu\text{g C-CO}_2 \text{ g}^{-1} \text{ soil h}^{-1}$) is a rate of CO_2 evolved from the agricultural and forest soils without the substrate addition.

PE = $100 \cdot (Q_{CO_2(\text{SOM})} - Q_{CO_2(\text{contr})}) / Q_{CO_2(\text{contr})}$, (%) is the average priming effect of microbial mineralization of substrates (n-hexadecane or n-hexadecanoic acid).

Value \pm standard deviation.

soil h^{-1} in 84 h (or 3.5 days) after n-hexadecanoic acid addition (Fig. 2, D, curve 5).

By analogy with n-hexadecane, parameterization of Equation [5] revealed the response of growing microorganisms after the addition of n-hexadecanoic acid to agricultural and forest soils. The parameter (r) for CO_2 production by agricultural soil was four times lower than for the forest soil, but specific CO_2 production rates (μ) were twice higher for agricultural than for forest soil (Table 3). The durations of lag-periods for CO_2 emission in agricultural and forest soils after n-hexadecanoic acid treatment were close to each other within the accuracy of analysis and estimated at about 30 hours (Table 3).

At the end of the 16-day experiments with n-hexadecanoic acid, the total quantities of metabolic CO_2 about $387 \pm 30 \mu\text{g C-CO}_2 \text{ g}^{-1} \text{ soil}$ and about $548 \pm 35 \mu\text{g C-CO}_2 \text{ g}^{-1} \text{ soil}$ were evolved from agricultural and forest soils, respectively (Table 4, Q_{CO_2}).

The carbon isotope indicator ($\delta^{13}\text{C}$) of CO_2 evolved from soils after the test substrate addition to the soils was measured during Δt_i time intervals (Table 5). Using of the test substrate as a mixture of n-hexadecanoic (PA) acid fractions labeled and unlabeled in C_1 - ^{13}C , the substrate content of ^{13}C isotope is characterized by different probability of ^{13}C detection on separate carbon sates of the PA molecule in the pool

of molecules, i.e., there is intramolecular ^{13}C isotope heterogeneity at the level of molecular pool (see section of *Mass isotopic balance for n-hexadecanoic acid*). In the present view, the succession of inclusion of C_1 - C_{16} atoms of the PA substrate into the microbial cell reaction is strictly determined and isotopic characteristic of PA fraction in CO_2 pool should be depended on the number carbon atoms of substrate molecules mineralized.

It is easily comprehended that the change in carbon isotope characteristics of CO_2 ($\delta^{13}\text{C}_{CO_2}$) as a result of PA mineralization will take place only at the initial stage of PA uptake (lag period). It should be specially emphasized that these considerations are significant only for presentation of the possible kinetics of CO_2 production from PA and SOM and are not used for estimation of the total value of priming effect.

As already noted, after all 16 carbon atoms of the first portions of consumed PA molecules are oxidized to CO_2 ; the isotope characteristics of CO_2 produced from the PA fraction will stabilize at a level of the mean value of isotopic indices of the PA molecule pool. In our case, it corresponds to $R = 0.011572$ or $\delta^{13}\text{C}_{\text{PA}} = +29.6\text{‰}$. The $\delta^{13}\text{C}_{CO_2}$ value (Table 5), which exceeds the mean $\delta^{13}\text{C}_{\text{PA}}$ at the initial stage of PA mineralization, is evidence of the existing mechanism of microbial mineralization of PA, i.e., PA mineralization begins with CO_2 production as a result of breakage of the $-\text{COOH}$ group. We believe that at

TABLE 5

Carbon isotope indicators of metabolic CO₂ ($\delta^{13}\text{C}_{\text{CO}_2}$, ‰) before and after n-hexadecanoic (palmitic) acid addition to soils and CO₂ fractions (F_{PA}) produced during mineralization of n-hexadecanoic acid

Time [days]	Agricultural soil		Forest soil	
	$\delta^{13}\text{C}_{\text{CO}_2}$ [‰]	F _{PA}	$\delta^{13}\text{C}_{\text{CO}_2}$ [‰]	F _{PA}
0.00	-26.6 ± 0.3	0.00	-26.2 ± 0.4	0.00
0.92	-10.0 ± 1.8	0.30 ± 0.04	-20.3 ± 0.68	0.11 ± 0.02
1.84	36.2 ± 1.8	<1.0	-10.0 ± 0.70	0.29 ± 0.02
2.80	20.0 ± 1.8	0.83 ± 0.04	19.4 ± 2.98	0.82 ± 0.06
3.67	13.6 ± 1.3	0.72 ± 0.03	31.1 ± 3.05	<1.0
4.58	8.7 ± 1.3	0.63 ± 0.03	21.6 ± 1.08	0.86 ± 0.03
5.57	8.6 ± 0.7	0.63 ± 0.02	16.1 ± 4.92	0.76 ± 0.09
6.49	8.9 ± 1.1	0.63 ± 0.02	12.8 ± 3.74	0.70 ± 0.07
8.40	7.5 ± 2.1	0.61 ± 0.04	9.9 ± 2.97	0.65 ± 0.06
11.40	7.0 ± 1.8	0.60 ± 0.04	7.5 ± 2.00	0.60 ± 0.04
15.10	7.0 ± 1.9	0.60 ± 0.05	-6.0 ± 3.00	0.36 ± 0.06
18.70	7.0 ± 2.0	0.60 ± 0.05	-10.0 ± 3.50	0.29 ± 0.07

*F_{PA} = ($\delta^{13}\text{C}_{\text{CO}_2} - \delta^{13}\text{C}_{\text{SOM}}$) / ($\delta^{13}\text{C}_{\text{PA}} - \delta^{13}\text{C}_{\text{SOM}}$) is the fraction of CO₂ produced during microbial mineralization of n-hexadecanoic acid, where $\delta^{13}\text{C}_{\text{PA}} = +29.6 \pm 0.2$ ‰ and $\delta^{13}\text{C}_{\text{SOM}}$ for CO₂ evolved from agricultural and forest soil were equal to $-26.6 (\pm 0.3)$ ‰ and $-26.2 (\pm 0.4)$ ‰, respectively.

Value \pm standard deviation.

the end of the lag period the mean value of CO₂ formed during mineralization of the PA test substrate approaches the value of $\delta^{13}\text{C} = +29$ ‰.

According to the pattern of experiments, the $\delta^{13}\text{C}$ value of the test substrate (n-hexadecanoic acid) amounted to as average of $+29.6$ ‰ and was rather different from $\delta^{13}\text{C}$ values of the SOM of agricultural and forest soils ($\delta^{13}\text{C}$ -CO₂ values which were equal to -26.6 ± 0.3 ‰ and -26.2 ± 0.4 ‰ for agricultural and forest soil, respectively)

The weighted average values of $\delta^{13}\text{C}_{\text{av}}$ of the respired CO₂ pools for agricultural and forest soils were $+12.7 \pm 0.3$ and $+15.7 \pm 0.5$ ‰ (Table 4, $\delta^{13}\text{C}_{\text{av}}$), respectively, and considered as an evidence of simultaneous metabolism of the introduced test substrate and SOM. Taking into account the isotopic characteristic of CO₂ evolved from soil with $\delta^{13}\text{C}_{\text{av}}$ (PA+SOM) and substrate introduced into soil with $\delta^{13}\text{C} = +29.6 \pm 0.2$ ‰, the carbon fractions of n-hexadecanoic (palmitic) acid and SOM was calculated using Eq. [3].

The quantities of CO₂ resulting from microbial mineralization of n-hexadecanoic acid were estimated at about 270 (± 26) $\mu\text{g C-CO}_2 \text{ g}^{-1} \text{ soil}$ and 412 (± 35) $\mu\text{g C-CO}_2 \text{ g}^{-1} \text{ soil}$ in the experiments with agricultural and forest soil (Table 4, Q_{CO₂(SUB)}), respectively. The total quantities of CO₂ derived from SOM mineralization after the n-hexadecanoic acid addition to agricultural and forest soils were estimated about 117 (± 26) $\mu\text{g C-CO}_2 \text{ g}^{-1} \text{ soil}$ and 136 (± 35) $\mu\text{g C-CO}_2 \text{ g}^{-1} \text{ soil}$, respectively.

Comparing the quantities of CO₂ produced with and without test substrate addition to soils, the priming of SOM mineralization in experiments was evaluated. The results of average PE calculation are presented (Table 4, PE(average)): the negative priming effects of n-hexadecanoic acid (PE_{agr} = $-36(\pm 28)\%$ and PE_{forest} = $-32(\pm 30)\%$) were detected for agricultural and forest soil, respectively. Concurrently to the stimulation of respiratory response by n-hexadecanoic acid, the kinetics of the production of CO₂ from SOM mineralization in soils amended with n-hexadecanoic acid relative to native process was determined for agricultural soil and forest soil (Fig. 2, B, curves 4 relative to curve 3; E, curves 8 relative to curve 7).

Microbial O₂ consumption and respiratory quotient. The kinetics of RQ variation in the soils with n-hexadecane (Fig. 1, C and F) and n-hexadecanoic acid (Fig. 2, C and F) were determined within certain periods (Δt_i) of about 8–12 h at the initial stages and 20–24 h at the final stages of the experiment. Following substrates addition, the respiratory quotient (RQ_i) decreased to 0.4 mol CO₂ mol⁻¹ O₂, which was lower than the theoretically expected values of 0.65 and 0.70 mol CO₂ mol⁻¹ O₂ for the complete oxidation of n-hexadecane and n-hexadecanoic acid and indicated relatively high O₂ requirement. The rapid increase of respiration rates in agricultural and forest soils at the initial stage of the experiments was reflected in the earlier lowering of the RQ_i value (Fig. 1, C and F and Fig. 2, C and F) and the ¹³C-CO₂ enrichment (Table 2, $\delta^{13}\text{C}_{\text{CO}_2}$ and Table 5, $\delta^{13}\text{C}_{\text{CO}_2}$).

TABLE 6

The quantities of O₂ taken up and CO₂ produced by soil microbiota in agricultural and forest soils with the test amounts of *n*-hexadecane and *n*-hexadecanoic acid

Treatment	O ₂ consumed [μM O ₂ g ⁻¹ soil]	CO ₂ produced [μM CO ₂ g ⁻¹ soil]	RQ _{av} [mol CO ₂ mol ⁻¹ O ₂]
<i>n</i> -hexadecane			
Agricultural soil	90 ± 5	54 ± 3	0.60 (±0.03)
Forest soil	82 ± 5	50 ± 5	0.61 (±0.04)
<i>n</i> -hexadecanoic acid			
Agricultural soil	56 ± 3	32 ± 3	0.57 (±0.02)
Forest soil	81 ± 4	52 ± 3	0.64 (±0.03)

Value ± standard deviation.

Henceforth, RQ_i continuously increased in proportion to substrate depletion, approaching the value observed prior to substrate addition into soil. Table 6 gives the amounts of oxygen taken up by microorganisms and CO₂ produced during the experiment with agricultural and forest soils after the addition of *n*-hexadecane and *n*-hexadecanoic acid, respectively. It is obvious that the ratios of CO₂ and O₂ molar quantities observed during the whole experimental period represent the average value of respiratory quotient (RQ_{av}) in the tested agricultural and forest soils, which is about 0.60(±0.03) and 0.61(±0.04) mol CO₂ mol⁻¹ O₂ in the experiments with *n*-hexadecane and 0.57(±0.02) and 0.64(±0.03) mol CO₂ mol⁻¹ O₂ in the experiments with *n*-hexadecanoic acid.

The carbon isotope characteristics were significantly modified in both the fumigated (microbial biomass) and non fumigated extracts by the hexadecane and palmitic acid addition after 17 and 35 days respectively (Table 7). The initial δ¹³C values ranged at -28‰ and the maximal values at -6‰ for the hexadecane treatment in the fumigated soil after 17 days. Thus, the highest values occurred 17 days after the hexadecane addition in the microbial C pool. The carbon isotope signature of microbial biomass and soil extract returned to -20 and -26‰ respectively after 28 days of palmitic acid addition. The maximal difference between the extracellular and the intracellular δ¹³C values was 8‰.

DISCUSSION

Basal soil respiration. The presence of *n*-hexadecane in native agricultural soil and its absence in forest soil (Table 1) can be considered as a result of difference in the contents of bacterial and fungal biomasses in soils (Ladygina et al. 2006; Ananyeva et al. 2008) and the associated different microbial biosynthesis of hydrocarbons. The obtained soil characteristics (Table 1) explain the fact that the RQ value during microbial oxidation of soil organic matter (SOM) and hydrocarbons in native agricultural soil (i.e., RQ_{init} = 0.73±0.05 mol CO₂ mol⁻¹ O₂) is lower than the value during mineralization of SOM and

partially oxidized hydrocarbons (e.g., fatty acids) in forest soil (i.e., RQ_{init} = 0.87 ± 0.05 mol CO₂ mol⁻¹ O₂).

***n*-Hexadecane consumption.** The addition of *n*-hexadecane stimulated microbial respiration in both agricultural and forest soils (Fig. 1, A, curve 1 and D, curve 5). The detection of two peaks of microbial CO₂ production rate in agricultural soil and one peak in forest soil after *n*-hexadecane addition may be treated as an evidence of essential difference in the microbial pools of these soil systems. This fact suggests that some of the hydrocarbon-oxidizing microorganisms may be present in soils in the active state (K-strategists), while others are in the “dormant” state (r-strategists).

Obviously, the presence of native *n*-hexadecane in agricultural soil (Table 1) contributes to the maintenance of some part of hydrocarbon-oxidizing microbiota in the active state, whereas its absence in forest soil may have kept this group of microbiota in the “dormant” state. Exogenous *n*-hexadecane added to agricultural soil is consumed by the active part of microbiota (K-strategists) and then, after a certain period of time, the “dormant” part of microbiota (r-strategists) is activated. Soil microbial communities in agricultural soil responded to *n*-hexadecane addition more rapidly compared to those of forest soil, with the lag-periods of about 23 ± 10 and 68 ± 13 hours, respectively (Table 3). The initial fractions of active microbial biomass capable of immediate growth on applied *n*-hexadecane in agricultural soil (parameter *r*) were higher by the order of magnitude than in forest soil (Table 3). The CO₂ production rates (μ) obtained for agricultural soil compared to forest soil were close to each other within the error limit of analysis (Table 3).

Previously it has been demonstrated with the growth of bacteria *Burkholderia* sp. on *n*-hexadecane, the carbon quantities of metabolic CO₂ and cell biomass were estimated by the ratio of 1.0 to 1.2 (Zyakun et al. 2003). It means that about 50% of the *n*-hexadecane carbon consumed by microbiota can be contained in the cell biomass. Like the ratio between cell biomass and metabolic CO₂ was presented in experiments with the yeast *Candida tropicalis* growth on *n*-hexadecane (Gmünder et al. 1981). Using techniques presented by Potthoff et al. 2003, the δ¹³C signatures of soil organic extracted samples of fumigation

TABLE 7

Carbon isotope signature in bulk soil and in the non-fumigated (–Fum) and fumigated (+ Fum) soil extracts in the agricultural and forest soil after *n*-hexadecane and *n*-hexadecanoic (palmitic) acid mineralization

Treatment	Soil, $\delta^{13}\text{C}$ [‰]	–Fum, $\delta^{13}\text{C}$ [‰]	+Fum, $\delta^{13}\text{C}$ [‰]	Fractionation, biotic [‰]
Agricultural soil				
Control	–26.3	–28.3	–28.2	0.1
+Hexadecane	–24.3	–14.6	–6.3	8.2
+Palmitic acid	–25.3	–26.8	–19.9	7.0
Forest soil				
Control	–26.6	–28.6	–27.2	1.3
+Hexadecane	–25.4	–23.1	–19.2	3.9
+Palmitic acid	–26.0	–27.8	–21.7	6.1

(+Fum) and non-fumigated (–Fum) with and without addition of *n*-hexadecane to the agricultural and forest soils were used to demonstrate the inclusion of *n*-hexadecane carbon into the microbial cell biomass (Table 7).

Assuming that the carbon yield of microbial biomass during *n*-hexadecane utilization by microorganisms is not higher than the yield of CO_2 , the total amount of substrate utilized in agricultural and forest soils may be $698 \pm 52 \mu\text{g C g}^{-1}$ soil and $582 \pm 64 \mu\text{g C g}^{-1}$ soil or about 54% and 45% of the $1300 \mu\text{g C g}^{-1}$ soil of *n*-hexadecane introduced into soils, respectively. In accordance with this estimation, about 46% (agricultural soil) and about 55% (forest soil) of introduced *n*-hexadecane may be adsorbed on soils and/or present as partially oxidized exometabolites.

The specific index of possible state of *n*-hexadecane introduced into soils is respiratory quotient (RQ) representing the ratio of molar concentrations of CO_2 produced and oxygen consumed during microbial metabolism of the substrate. Table 6 (RQ_{av}) gives the value of the average respiration quotient RQ_{av} for the 17-day experiment, equal to $0.60(\pm 0.03)$ and $0.61(\pm 0.04) \text{ mol CO}_2 \text{ mol}^{-1} \text{ O}_2$ during *n*-hexadecane metabolism in agricultural and forest soils, respectively. Since some part of oxygen is utilized for cell biomass production and biosynthesis of exometabolites, the resultant value was a little lower than the theoretical value ($\text{RQ}_{\text{av}} = 0.65 \text{ mol CO}_2 \text{ mol}^{-1} \text{ O}_2$).

The changes in RQ value during *n*-hexadecane uptake by soil microbiota, reached $0.4 \text{ mol CO}_2 \text{ mol}^{-1} \text{ O}_2$ (Fig. 1, C and F). This demonstrates that during the time Δt_i some part of the consumed O_2 is utilized for the formation of semi-oxidized products, from which CO_2 is formed later on. This fact is confirmed by the $\delta^{13}\text{C}$ values indicating substantial contribution of the carbon of introduced substrate into CO_2 production at the final stage of the experiment (Table 2, $\delta^{13}\text{C}_{\text{CO}_2}$).

n-Hexadecanoic acid consumption. The addition of *n*-hexadecanoic acid to soils under study is accompanied by the increase of metabolic activity of soil microbiota (Fig. 2, A, curve

1 and D, curve 5). We believe that two peaks of CO_2 production for agricultural soil and one peak for forest soil demonstrate the existence of two states of microorganisms (active and dormant) relative to the *n*-hexadecanoic acid consumption. Microbial activity represented by the rate of CO_2 production in forest soil is almost 4-fold higher compared to agricultural soil.

It is considered as an evidence of the known prevalence of fungal microflora in forest soil, for which *n*-hexadecanoic acid is an easily metabolized substrate (Dashti et al. 2008; Ananyeva et al. 2008). As follows from the analysis of parameters reflecting the kinetics of microbial metabolism of *n*-hexadecanoic acid in soils (Table 3), the initial specific rates of substrate uptake and CO_2 production (μ) in agricultural soil are twice higher than in forest soil, i.e., agricultural soil contains microorganisms capable for higher growth on *n*-hexadecanoic acid compared to forest soil. At the same time, value r as a metabolic index of the fraction of growing microorganisms in forest soil is 4-fold higher compared to agricultural soil, demonstrating the higher metabolic potential of microbial population resident in the forest soil with reference to *n*-hexadecanoic acid as a substrate. It is supposed that the same lag periods of microflora activation after substrate introduction into agricultural ($t_{\text{lag}} = 30 \pm 13 \text{ h}$) and forest ($t_{\text{lag}} = 30 \pm 14 \text{ h}$) soils is considered as an evidence of its consumption by fungal microbiota present in these soils.

The quantity of CO_2 formed due to mineralization of exogenous *n*-hexadecanoic acid during 16 days of exposure was $270 \pm 26 \mu\text{g C g}^{-1}$ soil and $412 \pm 35 \mu\text{g C g}^{-1}$ soil in agricultural and forest soil (Table 4, $\text{Q}_{\text{CO}_2(\text{SUB})}$), respectively. Similarly to the *n*-hexadecane microbial consumption, the $\delta^{13}\text{C}$ signatures of soil organic extracted samples of fumigation (+Fum) and non-fumigated (–Fum) with and without addition of *n*-hexadecanoic acid to the agricultural and forest soils were used to detect the inclusion of *n*-hexadecanoic acid carbon into the microbial cell biomass (Table 7).

Assuming that the carbon yield of microbial biomass during *n*-hexadecanoic acid metabolism is not higher than the yield of CO_2 , the total amount of substrate utilized in agricultural

and forest soils may be $540 \pm 52 \mu\text{g C g}^{-1}$ soil and $824 \pm 70 \mu\text{g C g}^{-1}$ soil or 42% and 63% of the $1300 \mu\text{g C g}^{-1}$ soil of introduced n-hexadecanoic acid, respectively. In accordance with this estimation, about 48% (agricultural soil) and about 37% (forest soil) of introduced n-hexadecanoic acid may be adsorbed on soils and/or present as partially oxidized exometabolites.

During the microbial metabolism of n-hexadecanoic acid in the course of 16-day experiment (Table 6), the value of total respiratory quotient in agricultural soil ($\text{RQ}_{\text{av}} = 0.57(\pm 0.02) \text{ mol CO}_2 \text{ mol}^{-1} \text{ O}_2$) was noticeably less than in forest soil ($\text{RQ}_{\text{av}} = 0.64(\pm 0.03) \text{ mol CO}_2 \text{ mol}^{-1} \text{ O}_2$), i.e., in agricultural soil the consumed oxygen is utilized to a greater extent for production of semi-oxidized metabolites and to a lesser extent for mineralization of the substrate to CO_2 . Like in the case of n-hexadecane, in the initial time (2-3 days) after substrate introduction (Fig. 2, C and F) RQ reached the value of about $0.4 \text{ mol CO}_2 \text{ mol}^{-1} \text{ O}_2$ and then, in 7 days, approached the theoretical value, i.e., $\text{RQ} = 0.7 \text{ mol CO}_2 \text{ mol}^{-1} \text{ O}_2$, in the experiments with forest soil, but remains at a level of $0.65 \text{ mol CO}_2 \text{ mol}^{-1} \text{ O}_2$ in agricultural soil.

Thus, the addition of both n-hexadecane and n-hexadecanoic (palmitic) acid induced intensive oxygen requirement (higher than expected theoretically) at the initial stages. The RQ values of the two soils decreased to $0.4 \text{ mol CO}_2 \text{ mol}^{-1} \text{ O}_2$ and were even lower than observed for the *Aspergillus niger* strain (Volke-Sepúlveda et al. 2003). Similarly, excessive O_2 requirement was observed with single species cultures (Chayabutra and Ju, 2000). The RQ value was about $0.52 \text{ mol CO}_2 \text{ mol}^{-1} \text{ O}_2$ during the growth of the yeast *Candida tropicalis* in chemostat culture, but this value changed from about 0.5 to $0.65 \text{ mol CO}_2 \text{ mol}^{-1} \text{ O}_2$ depending on growth condition (O_2 concentration, C/N ratios, growth rate) (Gmünder et al. 1981). This phenomenon may be caused by different times of O_2 consumption for oxidation of H_2 -equivalents/hydroxylation of alkane substrates and CO_2 production as the result of substrate mineralization in the citric acid cycle reactions.

Priming effects of substrates. The initial stage of n-hexadecane utilization by soil microbiota is its enzymatic hydroxylation and/or oxygenation by means of molecular oxygen (Metzler 1977; Smits et al. 1999; Marchant et al. 2006). The enzyme systems induced during this process, along with hydrocarbon substrate modification, are capable of hydroxylation of some part of SOM molecules, which are then used in microbial metabolism (Fontaine et al. 2004). Positive PE in the case of n-hexadecane is conditioned by the induction of microbial enzymes that perform hydroxylation/oxygenation of stable SOM compounds mineralized by soil microbiota to CO_2 .

The situation is different during the microbial uptake of n-hexadecanoic acid in the same soils. In the experiments with 16-day observation, negative PE values of this substrate were recorded after its addition to soil, i.e., introduction of exogenous substrate decreases the rate of SOM mineralization, and conse-

quently, CO_2 production is mainly due to biodegradation of the added substrate.

The special feature of n-hexadecanoic acid as compared to n-hexadecane lies in the fact that the oxidized end of the acid molecule is used as the initial site for β -oxidation process (van Hamme et al. 2003) and the microbial oxidation of saturated hydrocarbons (n-hexadecane) can not be initiated as easily as it does for n-hexadecanoic acid (Metzler 1977). The increased amount of ^{13}C during CO_2 production at the initial stage of n-hexadecanoic acid utilization as compared with calculated average content is an experimental evidence of this mechanism. As follows from Table 5 ($\delta^{13}\text{C}_{\text{CO}_2}$), the maximal value of $\delta^{13}\text{C}$ reached $+(31-36)\text{‰}$, whereas the calculated mean value was only $+29.6(\pm 0.2)\text{‰}$. Since the ^{13}C atom was in the carboxyl group of n-hexadecanoic acid (Experimental part), the share of ^{13}C isotope in CO_2 production at the first stages of test substrate degradation will be higher than it has been calculated for the average of n-hexadecanoic acid pool.

Nevertheless, under complete mineralization of molecule of n-hexadecanoic acid (i.e., 16 stages of successive removal of the COOH group) the carbon isotope composition of the formed total CO_2 from PA mineralization will inherit the average content of ^{13}C in the test substrate. In the case of mineralization of more than 50% of substrates introduced (Table 4, $X_{(\text{SUB})}$) it is absolutely justified to believe that the characteristics of carbon isotope composition of the formed CO_2 from PA mineralization will correspond to the calculated average value of isotopic characteristics for the substrate used.

Consequently, the mass isotopic balance was a basis for determining the production of CO_2 formed as a result of SOM mineralization after the introduction of substrate into soil. Comparison of this CO_2 production in the experiments with n-hexadecane and n-hexadecanoic acid and in controls made it possible to determine PE values with a high degree of reliability. It can be assumed that the microbial metabolism coupled with oxidative decarboxylation (β -oxidation) of n-hexadecanoic acid provides energy for growth of the main part of microbial cells in soils and it is considered as one of the most probable causes of the revealed phenomenon (i.e., negative PE of n-hexadecanoic acid with microbial mineralization of SOM).

Thus, in line with the research tasks mentioned, the following geomicrobiological index of soils should be noted: a) when exogenous hydrocarbons (n-hexadecane and n-hexadecanoic acid) are added to agricultural and forest soil, the metabolic activity of native soil microbiota noticeably increases. The enhancement of the rate of CO_2 production, consumption of molecular oxygen and the change of respiratory coefficient are indicative of microbial mineralization of organic substances in soils. b) The ^{13}C isotope signatures of metabolic CO_2 are incontestable evidence of the microbial use of test hydrocarbons and native soil organic matter as substrates; c) Quantitative comparison of rates of mineralization of SOM prior to and after addition of test hydrocarbons to agricultural and forest soils showed that microbial degradation of n-hexadecane or n-hexadecanoic acid

occurs with positive and negative priming-production of CO₂ at the expense of SOM, respectively. d) In agricultural and forest soils the microbial consumption of n-hexadecanoic acid is different from n-hexadecane and does not contribute to the activation of microbial SOM mineralization as strong as might be seen for n-hexadecane.

ACKNOWLEDGMENTS

These studies were financially supported by DFG-RUS 17/41/106, the Ministry for Science, Research and Culture (MWFK) of the Brandenburg state, EFRE and SENSOR and the Ministry of Education and Science of Russian Federation by grant RNP No 2.1.1/4341. The completion of this work was supported through the Cluster of Excellence "Climate System Analysis and Prediction- CliSAP" (EXC177), University of Hamburg, funded through the German Science Foundation (DFG). The authors thank Maria Koon for carefully editing the linguistic style of this article.

REFERENCES

- Adam G, Duncan H. 2002. Influence of diesel fuel on seed germination. *Environ Poll* 120:363–370.
- Adam G, Duncan H. 2003. The effect of diesel fuel on common vetch (*Vicia sativa* L.) plants. *Environ Geochem Health* 25:123–130.
- Abbassi BE, Shquirat WD. 2008. Kinetics of indigenous isolated bacteria used for ex-situ bioremediation of petroleum contaminated soil. *Water Air Soil Poll* 192:221–226.
- Alexander M. 1999. *Biodegradation and Bioremediation*. 2nd edn. Academic Press, London.
- Ananyeva ND, Susyan EA, Chernova OV, Wirth S. 2008. Microbial respiration activities of soils from different climatic regions of European Russia. *Euro J Soil Biol* 44:147–157.
- Asperger O, Sharychev AA, Matyashovaa RN, Losinov AB, Kleber H-P. 1986. Effect of oxygen limitation on the content of n-hexadecane-inducible cytochrome P-450 in *Acinetobacter calcoaceticus* strain EB 104. *J Basic Microbiol* 96:571–576.
- Baath E, Anderson T-H. 2002. Comparison of soil fungal/bacterial biomass ratios in soils investigated for enhanced carbon sequestration. *Soil Biol Biochem* 34:997–1007.
- Blagodatsky SA, Heinemeier O, Richter J. 2000. Estimating the active and total soil microbial biomass by kinetic respiration analysis. *Biol Fert Soils* 32:73–81.
- Blagodatskaya EV, Blagodatsky SA, Anderson T-H, Kuzyakov Y. 2007. Priming effects in Chernozem induced by glucose and N in relation to microbial growth strategies. *Appl Soil Ecol* 37: 95–105.
- Blagodatskaya EV, Blagodatsky SA, Anderson T-H, Kuzyakov Y. 2009. Contrasting effects of glucose, living roots and maize straw on microbial growth kinetics and substrate availability in soil. *Euro J Soil Sci* 60:186–197.
- Blagodatskaya EV, Kuzyakov Y. 2008. Mechanism of real and apparent priming effects and their dependence on soil microbial biomass and community structure: Critical review. *Biol Fert Soils* 45:115–131.
- Chayabutra C, Ju L. 2000. Degradation of n-hexadecane and its metabolites by *Pseudomonas aeruginosa* under microaerobic and anaerobic denitrifying conditions. *Appl Environ Microbiol* 66:493–498.
- Craig H. 1957. Isotopic standards for carbon and oxygen and correction factors for mass-spectrometric analysis of carbon dioxide. *Geochim Cosmochim Acta* 12:133–140.
- Dashti N, Al-Amadhi H, Khanafer M, Abdelhany S, Radwan S. 2008. Potential of hexadecane-utilizing soil microorganisms for growth on hexadecanol, hexadecanal, and hexadecanoic acid as sole sources of carbon and energy. *Chemosphere* 70:475–479.
- Delasse JH, Nyns EJ. 1974. Detection of cytochrome P-450 in subcellular fraction of *Endomycopsis lipolytica* grown on n-hexadecane. *Arch Internat Physiol Biochem* 82:179.
- Dilly O. 1999. Nitrogen and phosphorus requirement of the microbiota in soils of the Bornhöved Lake district. *Plant Soil* 212:175–183.
- Dilly O. 2001. Microbial respiratory quotient during basal metabolism and after glucose amendment in soils and litter. *Soil Biol Biochem* 33:117–127.
- Dilly O. 2003. Regulation of the respiratory quotient of soil microbiota by availability of nutrients. *FEMS Microbiol Ecol* 43:375–381.
- Dilly O, Zyakun A. 2008. Priming effect and respiratory quotient in a forest soil amended with glucose. *Geomicrobiol J* 25:425–431.
- Dyckmans J, Chander K, Joergensen RG, Priess J, Raubach M, Sehý U. 2003. Adenilates as an estimate of microbial biomass C in different soil groups. *Soil Biol Biochem* 35:1485–1491.
- Fontaine S, Bardoux G, Benest D, Verdier B, Mariotti A, Abbadie L. 2004. Mechanisms of the priming effect in a Savannah soil amended with cellulose. *Soil Sci Am J* 68:125–131.
- Gmünder FK, Käppeli O, Fiechter A. 1981. Chemostat studies on the assimilation of hexadecane by yeast *Candida tropicalis*. *Euro J Appl Microb Biotechnol* 12:120–134.
- Hamamura N, Olson SH, Ward DM, Inskeep WP. 2006. Microbial population dynamic associated with crude-oil biodegradation in diverse soils. *Appl Environ Microbiol* 72(9):6316–6324.
- Hamamura N, Fukui M, Ward DM, Inskeep WP. 2008. Assessing soil microbial populations responding to crude-oil amendment at different temperatures using phylogenetic, functional gene (alkB) and physiological analyses. *Environ Sci Technol* 42(20):7580–7586.
- Hassan AA, Sorial GA. 2008. n-hexadecane biodegradation in trickle bed air biofilters. *Water Air Soil Pollut* 8:287–296.
- Jobson A, McLaughlin M, Cook FD, Westlake DWS. 1974. Effect of amendments on the microbial utilization of oil applied to soil. *Appl Microbiol* 27(1):166–171.
- Juck D, Charles T, White L, Greer C. 2000. Polyphasic microbial community analysis of petroleum hydrocarbon-contaminated soils from two northern Canadian communities. *FEMS Microbiol Ecol* 33:241–249.
- Kaplan CW, Kitts CL. 2004. Bacterial succession in petroleum land treatment unit. *Appl Environ Microbiol* 70:1777–1786.
- Kawamoto S, Nosaki Ch, Tanaka A, Fukui S. 1978. Fatty acid β -oxidation system in microbodies of n-alkane-grown *Candida tropicalis*. *Eur J Biochem* 83:609–613.
- Ladygina N, Dedyukhina EG, Vainshtein MB. 2006. A review on microbial synthesis of hydrocarbons. *Proc Biochem* 41:1001–1014.
- Marchant R, Sharkey FH, Banat IM, Rahman TJ, Perfumo A. 2006. The degradation of n-hexadecane in soil by thermophilic geobacilli. *FEMS Microbiol Ecol* 56:44–54.
- Margesin R, Schinner F. 2001. Bioremediation (natural attenuation and biostimulation) of diesel-oil-contaminated soil in an alpine glacier skiing area. *Appl Environ Microbiol* 67(7):3127–3133.
- Margesin R, Schinner F. (eds.). 2005. *Manual for soil analysis-monitoring and assessing soil bioremediation*. Soil Biology, Vol. 5, Springer Verlag, Berlin, 359 pp.
- Margesin R, Hämmerle M, Tschërko D. 2007. Microbial activity and community composition during bioremediation of diesel-oil-contaminated soil: effects of hydrocarbon concentration, fertilizers, and incubation time. *Microb Ecol* 53:259–269.
- Marques AV, dos Santos SCC, Casella RC, Vital RL, Sebastian GV, Seldin L. 2008. Bioremediation potential of a tropical soil contaminated with a mixture of crude oil and production water. *J Microbiol Biotechnol* 18(12):1966–1974.
- Metzler DE. 1977. *Biochemistry. Theoretical Reactions of Living Cells*. Vol. 2. Chp. 9. Academic Press Inc. New York, San Francisco, London. P. 308–311.

- Mishra S, Jyot J, Kuhad RC, Lal B. 2001. Evaluation of inoculum addition to stimulate in situ bioremediation of oily-sludge-contaminated soil. *Appl Environ Microbiol* 67:1675–1681.
- Nikitina EV, Yakusheva OI, Zaripov SA, Galiev RA, Garusov AV, Naumova RP. 2003. Distribution and physiological state of microorganisms in petrochemical oily sludge. *Microbiology* 72:621–627.
- Palmroth MR, Munster U, Pichtel J, Puhakka JA. 2005. Metabolic response of microbiota to diesel fuel addition in vegetated soil. *Biodegradation* 16:91–101.
- Panikov NS. 1995. *Microbial Growth Kinetics*. Chapman and Hall, London, Glasgow, 378 pp.
- Panikov NS, Sizova MV. 1996. A kinetic method for estimating the biomass of microbial functional groups in soil. *J Microbiol Meth* 24:219–230.
- Pleshakova EV, Dubrovskaya EV, Turkovskaya OV. 2008. Efficiencies of introduction of an oil-oxidizing *Dietzia maris* strain and stimulation of natural microbial communities in remediation of polluted soil. *Appl Biochem Microbiol* 44(4):389–395.
- Potthoff M, Loftfild N, Buegger F, Wick B, John B, Joergensen RG, Flessa H. 2003. The determination of $\delta^{13}\text{C}$ in soil microbial biomass using fumigation-extraction. *Soil Biol Biochem* 35:947–954.
- Smits THM., Röthlisbergen M., Witholt B., van Beilen J.B. 1999. Molecular screening of alkane hydroxylase genes in gram-negative and gram-positive strains. *Environ Microbiol* 1:307–317.
- Stahl EA, Parkin TB, Christensen M. 1999. Fungal presence in paired cultivated and uncultivated soils in central Iowa, USA. *Biol Fertile Soils* 29: 91–100.
- Stenström J, Stedberg B, Johanson M. 1998. Kinetic of substrate-induced respiration (SIR): Theory. *Ambio* 27(1):35–39.
- Susyan EA, Ananyeva ND, Blagodatskaya EV. 2005. The antibiotic-aided distinguishing of fungal and bacterial substrate-induced respiration in various soil ecosystems. *Microbiology* 74:336–342.
- Susyan EA, Wirth S, Ananyeva ND, Stolnikova EV. 2011. Forest succession on abandoned arable soils in European Russia. – Impacts on microbial biomass, fungal-bacterial ratio, and basal CO₂ respiration activity. *Euro J Soil Biol* 47:169–174.
- Tauchert H, Grunow M, Aurich H. 1978. Regulation and properties of particular acceptor-dependent alcohol dehydrogenase of *Pseudomonas putida* during growth on n-alkanes. *Z Allg Microbiol* 18:675–680.
- Vance ED, Brooks PC, Jenkinson DS. 1987. Microbial biomass measurement in forest soils: The use of chloroform fumigation method for strongly acid soils. *Soil Biol Biochem* 19:697–702.
- Van Hamme JD, Singh A, Ward OP. 2003. Recent advantages in petroleum microbiology. *Microbial Mol Biol Rev* 67(4):503–549.
- Volke-Sepúlveda T, Gutiérrez-Rojas M, Favela-Torres E. 2003. Biodegradation of hexadecane in liquid and solid-state fermentations by *Aspergillus niger*. *Bioresour Technol* 87:81–86.
- Volke-Sepúlveda T, Gutiérrez-Rojas M, Favela-Torres E. 2006. Biodegradation of high concentrations of hexadecane by *Aspergillus niger* in solid-state system: Kinetic analysis. *Bioresour Technol* 97:1583–1591.
- Zucchi M, Angiolini L, Borin S, Brusetti L, Dietrich N, Gigliotti C, Barbieri P, Sorlini C, Daffonchio D. 2003. Response of bacterial community during bioremediation of an oil-polluted soil. *J Appl Microbiol* 94:248–257.
- Zyakun A, Dilly O. 2005. Respiratory quotient and priming effect in an arable soil induced by glucose. *Appl Biochem Microbiol* 41:512–520.
- Zyakun AM, Kosheleva IA, Zakharchenko VN, Kudryavtseva AI, Peshenko VP, Filonov AE, Boronin AM. 2003. The use of the $^{13}\text{C}/^{12}\text{C}$ ratio for the assay of the microbial oxidation of hydrocarbons. *Microbiology* 72:592–596.