

## PACKAGE LEAFLET: INFORMATION FOR THE USER

**Losec 10 mg hard capsules**  
**Losec 20 mg hard capsules**  
**Losec 40 mg hard capsules**  
Omeprazole

### **Read all of this leaflet carefully before you start taking this medicine.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you. Do not pass it on to others. It may harm them, even if their symptoms are the same as yours.
- If any of the side effects gets serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

### **In this leaflet:**

1. What Losec is and what it is used for
2. Before you take Losec
3. How to take Losec
4. Possible side effects
5. How to store Losec
6. Further information

## **1. WHAT LOSEC IS AND WHAT IT IS USED FOR**

Losec contains the active substance omeprazole. It belongs to a group of medicines called ‘proton pump inhibitors’. They work by reducing the amount of acid that your stomach produces.

Losec is used to treat the following conditions:

In adults:

- ‘Gastro-oesophageal reflux disease’ (GORD). This is where acid from the stomach escapes into the gullet (the tube which connects your throat to your stomach) causing pain, inflammation and heartburn.
- Ulcers in the upper part of the intestine (duodenal ulcer) or stomach (gastric ulcer).
- Ulcers which are infected with bacteria called ‘*Helicobacter pylori*’. If you have this condition, your doctor may also prescribe antibiotics to treat the infection and allow the ulcer to heal.
- Ulcers caused by medicines called NSAIDs (Non-Steroidal Anti-Inflammatory Drugs). Losec can also be used to stop ulcers from forming if you are taking NSAIDs.
- Too much acid in the stomach caused by a growth in the pancreas (Zollinger-Ellison syndrome).

In children:

*Children over 1 year of age and  $\geq 10$  kg*

- ‘Gastro-oesophageal reflux disease’ (GORD). This is where acid from the stomach escapes into the gullet (the tube which connects your throat to your stomach) causing pain, inflammation and heartburn.  
In children, the symptoms of the condition can include the return of stomach contents into the mouth (regurgitation), being sick (vomiting) and poor weight gain.

*Children and adolescents over 4 years of age*

- Ulcers which are infected with bacteria called ‘*Helicobacter pylori*’. If your child has this condition, your doctor may also prescribe antibiotics to treat the infection and allow the ulcer to heal.

## 2. BEFORE YOU TAKE LOSEC

### Do not take Losec

- If you are allergic (hypersensitive) to omeprazole or any of the other ingredients of Losec.
- If you are allergic to medicines containing other proton pump inhibitors (eg pantoprazole, lansoprazole, rabeprazole, esomeprazole).
- If you are taking a medicine containing nelfinavir (used for HIV infection)

If you are not sure, talk to your doctor or pharmacist before taking Losec.

### Take special care with Losec

Losec may hide the symptoms of other diseases. Therefore, if any of the following happen to you before you start taking Losec or while you are taking it, talk to your doctor straight away:

- You lose a lot of weight for no reason and have problems swallowing.
- You get stomach pain or indigestion.
- You begin to vomit food or blood.
- You pass black stools (blood-stained faeces).
- You experience severe or persistent diarrhoea, as omeprazole has been associated with a small increase in infectious diarrhoea.
- You have severe liver problems.

If you take Losec on a long-term basis (longer than 1 year) your doctor will probably keep you under regular surveillance. You should report any new and exceptional symptoms and circumstances whenever you see your doctor.

### Taking other medicines

Please tell your doctor or pharmacist if you are taking or have recently taken any other medicines, including medicines obtained without a prescription. This is because Losec can affect the way some medicines work and some medicines can have an effect on Losec.

Do not take Losec if you are taking a medicine containing **nelfinavir** (used to treat HIV infection).

Tell your doctor or pharmacist if you are taking any of the following medicines:

- Ketoconazole, itraconazole or voriconazole (used to treat infections caused by a fungus)
- Digoxin (used to treat heart problems)
- Diazepam (used to treat anxiety, relax muscles or in epilepsy)
- Phenytoin (used in epilepsy). If you are taking phenytoin, your doctor will need to monitor you when you start or stop taking Losec
- Medicines that are used to thin your blood, such as warfarin or other vitamin K blockers. Your doctor may need to monitor you when you start or stop taking Losec
- Rifampicin (used to treat tuberculosis)
- Atazanavir (used to treat HIV infection)
- Tacrolimus (in cases of organ transplantation)
- St John's wort (*Hypericum perforatum*) (used to treat mild depression)
- Cilostazol (used to treat intermittent claudication)
- Saquinavir (used to treat HIV infection)
- Clopidogrel (used to prevent blood clots (thrombi))

If your doctor has prescribed the antibiotics amoxicillin and clarithromycin as well as Losec to treat ulcers caused by *Helicobacter pylori* infection, it is very important that you tell your doctor about any other medicines you are taking.

### **Taking Losec with food and drink**

You can take your capsules with food or on an empty stomach.

### **Pregnancy and breast-feeding**

Before taking Losec, tell your doctor if you are pregnant or trying to get pregnant. Your doctor will decide whether you can take Losec during this time.

Your doctor will decide whether you can take Losec if you are breast-feeding.

### **Driving and using machines**

Losec is not likely to affect your ability to drive or use any tools or machines. Side effects such as dizziness and visual disturbances may occur (see section 4). If affected, you should not drive or operate machinery.

### **Important information about some of the ingredients of Losec**

Losec capsules contain lactose. If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicinal product.

## **3. HOW TO TAKE LOSEC**

Always take Losec exactly as your doctor has told you. You should check with your doctor or pharmacist if you are not sure.

Your doctor will tell you how many capsules to take and how long to take them for. This will depend on your condition and how old you are.

The usual doses are given below.

#### Adults:

To treat symptoms of GORD such as **heartburn and acid regurgitation:**

- If your doctor has found that your food pipe (gullet) has been slightly damaged, the usual dose is 20 mg once a day for 4-8 weeks. Your doctor may tell you to take a dose of 40 mg for a further 8 weeks if your gullet has not yet healed.
- The usual dose once the gullet has healed is 10 mg once a day.
- If your gullet has not been damaged, the usual dose is 10 mg once a day.

To treat **ulcers in the upper part of the intestine** (duodenal ulcer):

- The usual dose is 20 mg once a day for 2 weeks. Your doctor may tell you to take the same dose for a further 2 weeks if your ulcer has not yet healed.
- If the ulcer do not fully heal, the dose can be increased to 40 mg once a day for 4 weeks.

To treat **ulcers in the stomach** (gastric ulcer):

- The usual dose is 20 mg once a day for 4 weeks. Your doctor may tell you to take the same dose for a further 4 weeks if your ulcer has not yet healed.
- If the ulcer do not fully heal, the dose can be increased to 40 mg once a day for 8 weeks.

To **prevent the duodenal and stomach ulcers** from coming back:

- The usual dose is 10 mg or 20 mg once a day. Your doctor may increase the dose to 40 mg once a day.

To treat duodenal and stomach **ulcers caused by NSAIDs** (Non-Steroidal Anti-Inflammatory Drugs):

- The usual dose is 20 mg once a day for 4–8 weeks.

To **prevent duodenal and stomach ulcers** if you are taking **NSAIDs**:

- The usual dose is 20 mg once a day.

To treat **ulcers caused by *Helicobacter pylori*** infection and to stop them coming back:

- The usual dose is 20 mg Losec twice a day for one week.
- Your doctor will also tell you to take two antibiotics among amoxicillin, clarithromycin and metronidazole.

To treat too much acid in the stomach caused by a **growth in the pancreas (Zollinger-Ellison syndrome)**:

- The usual dose is 60 mg daily.
- Your doctor will adjust the dose depending on your needs and will also decide how long you need to take the medicine for.

Children:

To treat symptoms of GORD such as **heartburn and acid regurgitation**:

- Children over 1 year of age and with a body weight of more than 10 kg may take Losec. The dose for children is based on the child's weight and the doctor will decide the correct dose.

To treat **ulcers caused by *Helicobacter pylori*** infection and to stop them coming back:

- Children aged over 4 years may take Losec. The dose for children is based on the child's weight and the doctor will decide the correct dose.
- Your doctor will also prescribe two antibiotics called amoxicillin and clarithromycin for your child.

**Taking this medicine**

- It is recommended that you take your capsules in the morning.
- You can take your capsules with food or on an empty stomach.
- Swallow your capsules whole with half a glass of water. Do not chew or crush the capsules. This is because the capsules contain coated pellets which stop the medicine from being broken down by the acid in your stomach. It is important not to damage the pellets.

**What to do if you or your child have trouble swallowing the capsules**

- If you or your child have trouble swallowing the capsules:
  - Open the capsules and swallow the contents directly with half a glass of water or put the contents into a glass of still (non-fizzy) water, any acidic fruit juice (e.g. apple, orange or pineapple) or apple sauce.
  - Always stir the mixture just before drinking it (the mixture will not be clear). Then drink the mixture straight away or within 30 minutes.
  - To make sure that you have drunk all of the medicine, rinse the glass very well with half a glass of water and drink it. The solid pieces contain the medicine - do not chew or crush them.

**If you take more Losec than you should**

If you take more Losec than prescribed by your doctor, talk to your doctor or pharmacist straight away.

**If you forget to take Losec**

If you forget to take a dose, take it as soon as you remember it. However, if it is almost time for your next dose, skip the missed dose. Do not take a double dose to make up for a forgotten dose.

#### 4. POSSIBLE SIDE EFFECTS

Like all medicines, Losec can cause side effects, although not everybody gets them.

**If you notice any of the following rare but serious side effects, stop taking Losec and contact a doctor immediately:**

- Sudden wheezing, swelling of your lips, tongue and throat or body, rash, fainting or difficulties in swallowing (severe allergic reaction).
- Reddening of the skin with blisters or peeling. There may also be severe blisters and bleeding in the lips, eyes, mouth, nose and genitals. This could be ‘Stevens-Johnson syndrome’ or ‘toxic epidermal necrolysis’.
- Yellow skin, dark urine and tiredness which can be symptoms of liver problems.

Side effects may occur with certain frequencies, which are defined as follows:

Very common:	affects more than 1 user in 10
Common:	affects 1 to 10 users in 100
Uncommon:	affects 1 to 10 users in 1,000
Rare:	affects 1 to 10 users in 10,000
Very rare:	affects less than 1 user in 10,000
Not known:	frequency cannot be estimated from the available data

Other side effects include:

##### **Common side effects**

- Headache.
- Effects on your stomach or gut: diarrhoea, stomach pain, constipation, wind (flatulence).
- Feeling sick (nausea) or being sick (vomiting).

##### **Uncommon side effects**

- Swelling of the feet and ankles.
- Disturbed sleep (insomnia).
- Dizziness, tingling feelings such as “pins and needles”, feeling sleepy.
- Spinning feeling (vertigo).
- Changes in blood tests that check how the liver is working.
- Skin rash, lumpy rash (hives) and itchy skin.
- Generally feeling unwell and lacking energy.

##### **Rare side effects**

- Blood problems such as a reduced number of white cells or platelets. This can cause weakness, bruising or make infections more likely.
- Allergic reactions, sometimes very severe, including swelling of the lips, tongue and throat, fever, wheezing.
- Low levels of sodium in the blood. This may cause weakness, being sick (vomiting) and cramps.
- Feeling agitated, confused or depressed.
- Taste changes.
- Eyesight problems such as blurred vision.
- Suddenly feeling wheezy or short of breath (bronchospasm).
- Dry mouth.
- An inflammation of the inside of the mouth.
- An infection called “thrush” which can affect the gut and is caused by a fungus.
- Liver problems, including jaundice which can cause yellow skin, dark urine, and tiredness.
- Hair loss (alopecia).
- Skin rash on exposure to sunshine.
- Joint pains (arthralgia) or muscle pains (myalgia).

- Severe kidney problems (interstitial nephritis).
- Increased sweating.

#### **Very rare side effects**

- Changes in blood count including agranulocytosis (lack of white blood cells).
- Aggression.
- Seeing, feeling or hearing things that are not there (hallucinations).
- Severe liver problems leading to liver failure and inflammation of the brain.
- Sudden onset of a severe rash or blistering or peeling skin. This may be associated with a high fever and joint pains (Erythema multiforme, Stevens-Johnson syndrome, toxic epidermal necrolysis).
- Muscle weakness.
- Enlarged breasts in men.
- Hypomagnesaemia.

Losec may in very rare cases affect the white blood cells leading to immune deficiency. If you have an infection with symptoms such as fever with a **severely** reduced general condition or fever with symptoms of a local infection such as pain in the neck, throat or mouth or difficulties in urinating, you must consult your doctor as soon as possible so that a lack of white blood cells (agranulocytosis) can be ruled out by a blood test. It is important for you to give information about your medicine at this time.

Do not be concerned by this list of possible side effects. You may not get any of them. If any of the side effects get serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

### **5. HOW TO STORE LOSEC**

- Keep out of the reach and sight of children.
- Do not use Losec after the expiry date which is stated on the pack after EXP. The expiry date refers to the last day of that month.
- Do not store above 30°C.
- Store this blister in the original package or keep the bottle tightly closed in order to protect from moisture.
- Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

### **6. FURTHER INFORMATION**

#### **What Losec contains**

- The active substance is omeprazole. Losec capsules contain 10 mg, 20 mg or 40 mg of omeprazole.
- The other ingredients are disodium hydrogen phosphate dihydrate, hydroxypropylcellulose, hydroxypropyl methylcellulose, lactose anhydrous, magnesium stearate, mannitol, methacrylic acid co-polymer, microcrystalline cellulose, macrogol (polyethylene glycol), sodium lauryl sulphate, iron oxide, titanium dioxide, gelatine, printing ink (containing shellac, ammonium hydroxide, potassium hydroxide and black iron oxide).

### What Losec looks like and contents of the pack

- Losec 10 mg capsules have a pink body, marked 10 and a pink cap marked A/OS.
- Losec 20 mg capsules have a pink body, marked 20 and a reddish-brown cap marked A/OM.
- Losec 40 mg capsules have a reddish-brown body marked 40 and a reddish-brown cap marked A/OL.

#### Pack sizes:

- 10 mg:
  - HDPE bottles of 5, 7, 10, 14, 15, 28, 30, 50, 56, 60 or 100 capsules; hospital packs of 140, 280 or 700 capsules.
  - Blisters of 7, 14, 15, 28, 30, 35, 50, 56 and 84 capsules.
- 20 mg:
  - HDPE bottles of 5, 7, 10, 14, 15, 28, 30, 50, 60 or 100 capsules; hospital packs of 140, 280, or 700 capsules.
  - Blisters of 7, 14, 15, 28, 30, 50, 60 or 84 capsules.
- 40 mg:
  - HDPE bottles of 5, 7, 14, 15, 28, 30 or 60 capsules; hospital packs of 140, 280, or 700 capsules.
  - Blisters of 7, 14, 15, 28 or 30 capsules.

Not all pack sizes may be marketed.

### Marketing Authorisation Holder and Manufacturer

The Marketing Authorisations for Losec capsules are held by AstraZeneca UK Ltd, 600 Capability Green, Luton, LU1 3LU, United Kingdom.

Losec capsules are manufactured by AstraZeneca AB, S-151 85, Södertälje, Sweden or AstraZeneca UK Ltd, Silk Road Business Park, Macclesfield, Cheshire, SK10 2NA, United Kingdom.

To listen to or request a copy of this leaflet in Braille, large print or audio please call, free of charge:

0800 198 5000 (UK only)

Please be ready to give the following information:

<b>Product name</b>	<b>Reference number</b>
Losec 10 mg Capsules	17901/0132
Losec 20 mg Capsules	17901/0133
Losec 40 mg Capsules	17901/0134

This is a service provided by the Royal National Institute of Blind People.

**This medicinal product is authorised in the Member States of the EEA under the following names, Losec, Mopral and Antra 10, 20 and 40 mg hard capsules.**

This leaflet was last updated in July 2010.

© AstraZeneca 2010

Losec is a trade mark of the AstraZeneca group of companies.

GI 10 0012