Care Quality Commission

Inspection Evidence Table

Ashington House Surgery (1-659227076)

Inspection date: 11 August 2021

Date of data download: 09 August 2021

Overall rating: Good

We inspected Ashington House Surgery in November 2019; the service was rated as Requires Improvement overall. A requirement was made to improve systems and processes to ensure they operated effectively to ensure good governance of service provision.

At this inspection we found that the necessary improvements had been made and the requirement notice was met.

Please note: Any Quality Outcomes Framework (QOF) data relates to 2019/20.

Safe

Rating: Good

When we inspected Ashington House Surgery in November 2019, the service was rated as Requires Improvement for the safe key question.

• A requirement was made to improve systems for managing medicines, including tracking of prescriptions and ensuring they were collected; ensuring necessary blood tests were carried out prior to a prescription being issued; and reviewing use of antibiotic prescribing as this was higher than clinical commissioning group and national averages.

At this inspection we found that a protocol had been introduced to ensure that prescriptions were destroyed securely if they had not been collected after two months. Checks were made with patients prior to a prescription being destroyed. The practice had also moved to electronic prescriptions, so paperbased prescriptions were rarely used. Necessary blood tests were carried out prior to a prescription being issued. The practice had carried out reviews on antibiotic prescribing and levels of prescribing were in line with clinical commissioning group and national averages.

Safety systems and processes

The practice had clear systems, practices and processes to keep people safe and safeguarded from abuse.

Safeguarding	Y/N/Partial
There was a lead member of staff for safeguarding processes and procedures.	Y
Safeguarding systems, processes and practices were developed, implemented and communicated to staff.	Y

Safeguarding	Y/N/Partial	
There were policies covering adult and child safeguarding which were accessible to all staff.	Y	
Policies and procedures were monitored, reviewed and updated.	Y	
Partners and staff were trained to appropriate levels for their role.	Y	
There was active and appropriate engagement in local safeguarding processes.	Y	
The Out of Hours service was informed of relevant safeguarding information.	Y	
There were systems to identify vulnerable patients on record.		
Disclosure and Barring Service (DBS) checks were undertaken where required.		
Staff who acted as chaperones were trained for their role.		
There were regular discussions between the practice and other health and social care professionals such as health visitors, school nurses, community midwives and social workers to support and protect adults and children at risk of significant harm.		

Recruitment systems	Y/N/Partial
Recruitment checks were carried out in accordance with regulations (including for agency staff and locums).	Y
Staff vaccination was maintained in line with current Public Health England (PHE) guidance if relevant to role.	Y
There were systems to ensure the registration of clinical staff (including nurses and pharmacists) was checked and regularly monitored.	Y

Safety systems and records	Y/N/Partial
There was a record of portable appliance testing or visual inspection by a competent person. Date of last inspection/test: April 2021	Y
There was a record of equipment calibration. Date of last calibration: April 2021	Y
There were risk assessments for any storage of hazardous substances for example, liquid nitrogen, storage of chemicals.	Y
There was a fire procedure.	Y
A fire risk assessment had been completed. Date of completion: 23/3/2021	Y
Actions from fire risk assessment were identified and completed.	Y

Health and safety	Y/N/Partial
Premises/security risk assessment had been carried out.	Y

Date of last assessment: 6/3/2021	
Health and safety risk assessments had been carried out and appropriate actions taken.	V
Date of last assessment: 6/3/2021	T

Infection prevention and control

Appropriate standards of cleanliness and hygiene were met.

	Y/N/Partial
There was an infection risk assessment and policy.	Y
Staff had received effective training on infection prevention and control.	Partial
Infection prevention and control audits were carried out. Date of last infection prevention and control audit: 28/4/2021	Y
The practice had acted on any issues identified in infection prevention and control audits.	Y
There was a system to notify Public Health England of suspected notifiable diseases.	Y
The arrangements for managing waste and clinical specimens kept people safe.	Y

Explanation of any answers and additional evidence:

Staff we spoke to confirmed they had received infection control training; however, the practice was unable to demonstrate this through records of training held.

Policies and procedures had been reviewed to include COVID-19 risks, such as increased cleaning by practice staff of 'high touch points', for example door handles and light switches.

Cleaning of the premises was carried out by an external contractor who undertook monthly audits with a member of the practice staff. The most recent cleaning audit in July 2021 showed 100% compliance with cleaning schedules.

Legionella monitoring checks showed that there were issues with water temperatures not being within safe limits to prevent Legionella, in one of the patient's toilet. (Legionella is a term for a bacterium which can contaminate water systems in buildings). Funding had been secured to convert another toilet next to this one into a sluice. Work was being carried in the month following our site visit and plans showed that the faulty tap would be replaced, as part of this work.

Risks to patients

There were adequate systems to assess, monitor and manage risks to patient safety.

	Y/N/Partial
There was an effective approach to managing staff absences and busy periods.	Y
There was an effective induction system for temporary staff tailored to their role.	Y
The practice was equipped to respond to medical emergencies (including suspected sepsis) and staff were suitably trained in emergency procedures.	Y
Receptionists were aware of actions to take if they encountered a deteriorating or acutely unwell patient and had been given guidance on identifying such patients.	Ý

Information to deliver safe care and treatment

Staff had the information they needed to deliver safe care and treatment.

	Y/N/Partial
Individual care records, including clinical data, were written and managed securely and in line with current guidance and relevant legislation.	Y
There was a system for processing information relating to new patients including the summarising of new patient notes.	Ŷ
There were systems for sharing information with staff and other agencies to enable them to deliver safe care and treatment.	Y
Referrals to specialist services were documented, contained the required information and there was a system to monitor delays in referrals.	Y
There was a documented approach to the management of test results and this was managed in a timely manner.	Y
There was appropriate clinical oversight of test results, including when reviewed by non- clinical staff.	Y

Appropriate and safe use of medicines

The practice had systems for the appropriate and safe use of medicines, including medicines optimisation

Indicator	Practice	CCG average	England average	England comparison
Number of antibacterial prescription items prescribed per Specific Therapeutic group Age-sex Related Prescribing Unit (STAR PU) (01/04/2020 to 31/03/2021) (NHS Business Service Authority - NHSBSA)	0.78	0.66	0.70	No statistical variation
The number of prescription items for co- amoxiclav, cephalosporins and quinolones as a percentage of the total number of prescription items for selected antibacterial drugs (BNF 5.1 sub-set). (01/04/2020 to 31/03/2021) (NHSBA)	12.1%	11.3%	10.2%	No statistical variation
Average daily quantity per item for Nitrofurantoin 50 mg tablets and capsules, Nitrofurantoin 100 mg m/r capsules, Pivmecillinam 200 mg tablets and Trimethoprim 200 mg tablets prescribed for uncomplicated urinary tract infection (01/10/2020 to 31/03/2021)	5.59	4.86	5.37	No statistical variation
(NHSBSA) Total items prescribed of Pregabalin or Gabapentin per 1,000 patients (01/10/2020 to 31/03/2021) (NHSBSA)	168.1‰	123.1‰	126.9‰	No statistical variation

Indicator	Practice	CCG average	England average	England comparison
Average daily quantity of Hypnotics prescribed per Specific Therapeutic group Age-sex Related Prescribing Unit (STAR PU) (01/04/2020 to 31/03/2021) (NHSBSA)	1 20	0.71	0.66	No statistical variation
Number of unique patients prescribed multiple psychotropics per 1,000 patients (01/07/2020 to 31/12/2020) (NHSBSA)		5.8‰	6.7‰	No statistical variation

Note: ‰ means *per 1,000* and it is **not** a percentage.

Medicines management	Y/N/Partia		
The practice ensured medicines were stored safely and securely with access restricted to authorised staff.	Y		
Blank prescriptions were kept securely, and their use monitored in line with national guidance.			
Staff had the appropriate authorisations to administer medicines (including Patient Group Directions or Patient Specific Directions).	Y		
The practice could demonstrate the prescribing competence of non-medical prescribers, and there was regular review of their prescribing practice supported by clinical supervision or peer review.	Y		
There was a process for the safe handling of requests for repeat medicines and evidence of structured medicines reviews for patients on repeat medicines.	Y		
The practice had a process and clear audit trail for the management of information about changes to a patient's medicines including changes made by other services.			
There was a process for monitoring patients' health in relation to the use of medicines including high risk medicines (for example, warfarin, methotrexate and lithium) with appropriate monitoring and clinical review prior to prescribing.	Y		
The practice monitored the prescribing of controlled drugs. (For example, investigation of unusual prescribing, quantities, dose, formulations and strength).	Y		
There were arrangements for raising concerns around controlled drugs with the NHS England Area Team Controlled Drugs Accountable Officer.			
If the practice had controlled drugs on the premises there were appropriate systems and written procedures for the safe ordering, receipt, storage, administration, balance checks and disposal of these medicines, which were in line with national guidance.			
The practice had taken steps to ensure appropriate antimicrobial use to optimise patient outcomes and reduce the risk of adverse events and antimicrobial resistance.	Y		
For remote or online prescribing there were effective protocols for verifying patient identity.	Y		
The practice held appropriate emergency medicines, risk assessments were in place to determine the range of medicines held, and a system was in place to monitor stock levels and expiry dates.	Y		
There was medical oxygen and a defibrillator on site and systems to ensure these were regularly checked and fit for use.	Y		

Medicines management	Y/N/Partial
Vaccines were appropriately stored, monitored and transported in line with PHE guidance to ensure they remained safe and effective.	Y

Explanation of any answers and additional evidence:

- We carried out searches of clinical records prior to the site visit, to see how the practice managed patients' care and treatment. Results from the searches on medicines management showed that required blood tests were not carried out prior to some medicines being prescribed, such as disease modifying medicines and heart medicines.
- We discussed this with the practice, and they provided evidence which showed that patients prescribed these medicines had their condition monitored by secondary care who carried out the necessary blood tests. This information was held on a separate system, known as ICE and was referred to when clinicians authorised a prescription.
- The ICE system did not automatically link to patients' main records on SystmOne, which the practice used to record care and treatment. The practice said that a 'manual' download of results from ICE could be undertaken and the information added to each patient record individually. The practice also identified a coding issue with their systems, as each time a repeat medicine was prescribed, a review of blood tests was carried out, but not coded as a medicine review. Action was taken immediately to correct this.

Track record on safety and lessons learned and improvements made

The practice learned and made improvements when things went wrong.

Significant events	Y/N/Partial
The practice monitored and reviewed safety using information from a variety of sources.	Y
Staff knew how to identify and report concerns, safety incidents and near misses.	Y
There was a system for recording and acting on significant events.	Y
Staff understood how to raise concerns and report incidents both internally and externally.	Y
There was evidence of learning and dissemination of information.	Y
Number of events recorded in last 12 months:	6
Number of events that required action:	6

Example(s) of significant events recorded and actions by the practice.

Event	Specific action taken
Another GP practice contacted Ashington House Surgery to request that a patient was added to the paramedic home visiting list. The home visiting paramedic on that day was not informed of the additional visit.	visit not taking place, they contacted the other GP practice to check whether it was clinically appropriate to visit the following day. As there were no GPs on site

 day. They considered the event to be a 'near miss' and therefore treated the incident as a significant event. A protocol for home visiting was put in place which contained details on ensuring a visiting clinician was informed of all visit requests by telephone.
 The member of staff was given appropriate support and training.

Safety alerts	Y/N/Partial
There was a system for recording and acting on safety alerts.	Y
Staff understood how to deal with alerts.	Y
Explanation of any answers and additional evidence:	·
We saw examples of actions taken on recent alerts for example changes to the brane	d of the tablet

thyroxine, used to correct thyroxine levels in the blood, and use of ibuprofen, an anti-inflammatory medicine, in patients with COVID-19.

Effective

Rating: Good

Effective needs assessment, care and treatment

Patients' needs were assessed, and care and treatment was delivered in line with current legislation, standards and evidence-based guidance supported by clear pathways and tools.

	Y/N/Partial
The practice had systems and processes to keep clinicians up to date with current evidence-based practice.	Y
Patients' immediate and ongoing needs were fully assessed. This included their clinical needs and their mental and physical wellbeing.	Y
Patients presenting with symptoms which could indicate serious illness were followed up in a timely and appropriate way.	Y
We saw no evidence of discrimination when staff made care and treatment decisions.	Y
Patients' treatment was regularly reviewed and updated.	Y
There were appropriate referral pathways to make sure that patients' needs were addressed.	Y
Patients were told when they needed to seek further help and what to do if their condition deteriorated.	Υ

Older people

Population group rating: Good

Findings

- The practice used a clinical tool to identify older patients who were living with moderate or severe frailty. Those identified received a full assessment of their physical, mental and social needs.
- The practice followed up on older patients discharged from hospital. It ensured that their care plans and prescriptions were updated to reflect any extra or changed needs.
- The practice carried out structured annual medicines reviews for older patients. The practice was in the process of arranging medicine and health reviews to happen in the month of birth of patients.
- Staff had appropriate knowledge of treating older people including their psychological, mental and communication needs.
- Health checks, including frailty assessments, were offered to patients over 75 years of age.
- Flu, shingles and pneumonia vaccinations were offered to relevant patients in this age group.

People with long-term conditions

Population group rating: Good

Findings

- Patients with long-term conditions were offered a structured annual review to check their health and medicines needs were being met. For patients with the most complex needs, the GP worked with other health and care professionals to deliver a coordinated package of care.
- Staff who were responsible for reviews of patients with long-term conditions had received specific training. A practice nurse and the pharmacist had received training on managing heart failure and were responsible for monitoring these patients.
- GPs followed up patients who had received treatment in hospital or through out of hours services for an acute exacerbation of asthma.
- The practice shared clear and accurate information with relevant professionals when deciding care delivery for patients with long-term conditions.
- The practice could demonstrate how they identified patients with commonly undiagnosed conditions, for example diabetes, chronic obstructive pulmonary disease (COPD), atrial fibrillation and hypertension.
- Adults with newly diagnosed cardio-vascular disease were offered statins.
- Patients with suspected hypertension were offered ambulatory blood pressure monitoring.
- Patients with atrial fibrillation were assessed for stroke risk and treated appropriately.
- Patients with COPD were offered rescue packs.
- Patients with asthma were offered an asthma management plan.

Long-term conditions	Practice	CCG average	England average	England comparison
The percentage of patients with asthma, on the register, who have had an asthma review in the preceding 12 months that includes an assessment of asthma control using the 3 RCP questions. (01/04/2019 to 31/03/2020)	69.3%	75.2%	76.6%	No statistical variation
PCA* rate (number of PCAs).	11.5% (93)	10.9%	12.3%	N/A
The percentage of patients with COPD who have had a review, undertaken by a healthcare professional, including an assessment of breathlessness using the Medical Research Council dyspnoea scale in the preceding 12 months (01/04/2019 to 31/03/2020) (QOF)	77.7%	86.1%	89.4%	Tending towards variation (negative)
PCA rate (number of PCAs).	5.1% (8)	13.0%	12.7%	N/A

*PCA:. Personalised Care Adjustments allow practices to remove a patient from the indicator for limited, specified reasons.

Long-term conditions	Practice	CCG average	England average	England comparison
The percentage of patients aged 79 years or under with coronary heart disease in whom the last blood pressure reading (measured in the preceding 12 months) is 140/90 mmHg or less (01/04/2019 to 31/03/2020) (QOF)	79.4%	80.2%	82.0%	No statistical variation
PCA* rate (number of PCAs).	1.8% (4)	5.6%	5.2%	N/A
The percentage of patients with diabetes, on the register, without moderate or severe frailty in whom the last IFCC-HbA1c is 58 mmol/mol or less in the preceding 12 months (01/04/2019 to 31/03/2020) (QOF)	66.8%	66.0%	66.9%	No statistical variation
PCA rate (number of PCAs).	8.3% (48)	17.1%	15.3%	N/A
The percentage of patients aged 79 years or under with hypertension in whom the last blood pressure reading (measured in the preceding 12 months) is 140/90 mmHg or less (01/04/2019 to 31/03/2020) (QOF)	69.7%	71.3%	72.4%	No statistical variation
PCA rate (number of PCAs).	2.8% (41)	7.4%	7.1%	N/A
In those patients with atrial fibrillation with a record of a CHA2DS2-VASc score of 2 or more, the percentage of patients who are currently treated with anti-coagulation drug therapy (01/04/2019 to 31/03/2020) (QOF)	84.9%	92.6%	91.8%	No statistical variation
PCA rate (number of PCAs).	7.8% (10)	4.5%	4.9%	N/A
The percentage of patients with diabetes, on the register, without moderate or severe frailty in whom the last blood pressure reading (measured in the preceding 12 months) is 140/80 mmHg or less (01/04/2019 to 31/03/2020) (QOF)	85.9%	74.0%	75.9%	Tending towards variation (positive)
PCA rate (number of PCAs).	6.2% (36)	11.5%	10.4%	N/A

*PCA:. Personalised Care Adjustments allow practices to remove a patient from the indicator for limited, specified reasons.

Any additional evidence or comments

The impact of the COVID-19 pandemic had limited the practice's ability to carry out all the required checks for COPD. They had not been able to carry out an assessment of breathlessness during the first wave of the virus due to it being an aerosol producing procedure. In addition, Quality Outcome Framework indicators were suspended, this meant they were unable to make improvements to COPD indicators.

Families, children and young people

Population group rating: Good

Findings

- The practice has met the minimum 90% for all of five childhood immunisation uptake indicators. The practice has met the WHO based national target of 95% (the recommended standard for achieving herd immunity) for three of five childhood immunisation uptake indicators.
- The practice contacted the parents or guardians of children due to have childhood immunisations.
- The practice had arrangements for following up failed attendance of children's appointments following an appointment in secondary care or for immunisation and would liaise with health visitors when necessary.
- The practice had arrangements to identify and review the treatment of newly pregnant women on long-term medicines. These patients were provided with advice and post-natal support in accordance with best practice guidance.
- Young people could access services for sexual health and contraception.
- Staff had the appropriate skills and training to carry out reviews for this population group.

Child Immunisation	Numerator	Denominator	Practice %	Comparison to WHO target of 95%
The percentage of children aged 1 who have completed a primary course of immunisation for Diphtheria, Tetanus, Polio, Pertussis, Haemophilus influenza type b (Hib), Hepatitis B (Hep B) ((i.e. three doses of DTaP/IPV/Hib/HepB) (01/04/2019 to 31/03/2020) (NHS England)	116	121	95.9%	Met 95% WHO based target
The percentage of children aged 2 who have received their booster immunisation for Pneumococcal infection (i.e. received Pneumococcal booster) (PCV booster) (01/04/2019 to 31/03/2020) (NHS England)	117	123	95.1%	Met 95% WHO based target
The percentage of children aged 2 who have received their immunisation for Haemophilus influenza type b (Hib) and Meningitis C (MenC) (i.e. received Hib/MenC booster) (01/04/2019 to 31/03/2020) (NHS England)	117	123	95.1%	Met 95% WHO based target
The percentage of children aged 2 who have received immunisation for measles, mumps and rubella (one dose of MMR) (01/04/2019 to 31/03/2020) (NHS England)	115	123	93.5%	Met 90% minimum
The percentage of children aged 5 who have received immunisation for measles, mumps and rubella (two doses of MMR) (01/04/2019 to 31/03/2020) (NHS England)	107	113	94.7%	Met 90% minimum

Note: Please refer to the CQC guidance on Childhood Immunisation data for more information: https://www.cqc.org.uk/guidance-providers/gps/how-we-monitor-gp-practices

Working age people (including those recently retired and students)

Population group rating: Good

Findings

- The practice had systems to inform eligible patients to have the meningitis vaccine, for example before attending university for the first time.
- Patients had access to appropriate health assessments and checks including NHS checks for patients aged 40 to 74. There was appropriate and timely follow-up on the outcome of health assessments and checks where abnormalities or risk factors were identified.
- Patients could book or cancel appointments online and order repeat medicines without the need to attend the surgery.

Cancer Indicators	Practice	CCG average	England average	England comparison
The percentage of women eligible for cervical cancer screening at a given point in time who were screened adequately within a specified period (within 3.5 years for women aged 25 to 49, and within 5.5 years for women aged 50 to 64). (Snapshot date: 31/03/2021) (Public Health England)	74.3%	N/A	80% Target	Below 80% target
Females, 50-70, screened for breast cancer in last 36 months (3 year coverage, %) (01/04/2019 to 31/03/2020) (PHE)	79.1%	74.4%	70.1%	N/A
Persons, 60-74, screened for bowel cancer in last 30 months (2.5 year coverage, %) (01/04/2019 to 31/03/2020) (PHE)	65.4%	67.0%	63.8%	N/A
The percentage of patients with cancer, diagnosed within the preceding 15 months, who have a patient review recorded as occurring within 6 months of the date of diagnosis (01/04/2019 to 31/03/2020) (QOF)	92.0%	89.3%	92.7%	N/A
Number of new cancer cases treated (Detection rate: % of which resulted from a two week wait (TWW) referral) (01/04/2019 to 31/03/2020) (PHE)	44.1%	56.7%	54.2%	No statistical variation

Any additional evidence or comments

The practice was part of a primary care network cervical screening programme, who worked together to offer appointments. Alerts were placed on patient records to show when screening was due. When possible staff took would opportunistically screen women. Staff said that if they took an opportunistic cervical screen, before the recommended time period then the sample was not processed.

People whose circumstances make them vulnerable

Population group rating: Good

Findings

- Same day appointments and longer appointments were offered when required.
- All patients with a learning disability were offered an annual health check.
- End of life care was delivered in a coordinated way which took into account the needs of those whose circumstances may make them vulnerable.
- The practice had a system for vaccinating patients with an underlying medical condition according to the recommended schedule.
- The practice demonstrated that they had a system to identify people who misused substances.

People experiencing poor mental health

Population group rating: Good

(including people with dementia)

Findings

- The practice assessed and monitored the physical health of people with mental illness, severe mental illness, and personality disorder by providing access to health checks, interventions for physical activity, obesity, diabetes, heart disease, cancer and access to 'stop smoking' services.
- Same day and longer appointments were offered when required.
- There was a system for following up patients who failed to attend for administration of long-term medicines.
- When patients were assessed to be at risk of suicide or self-harm the practice had arrangements in place to help them to remain safe.
- Patients at risk of dementia were identified and offered an assessment to detect possible signs of dementia. When dementia was suspected there was an appropriate referral for diagnosis.
- Patients with poor mental health, including dementia, were referred to appropriate services.

Mental Health Indicators	Practice	CCG average	England average	England comparison
The percentage of patients with schizophrenia, bipolar affective disorder and other psychoses who have a comprehensive, agreed care plan documented in the record, in the preceding 12 months (01/04/2019 to 31/03/2020) (QOF)	79.3%	78.7%	85.4%	No statistical variation
PCA* rate (number of PCAs).	13.7% (13)	20.2%	16.6%	N/A
The percentage of patients diagnosed with dementia whose care plan has been reviewed in a face-to-face review in the preceding 12 months (01/04/2019 to 31/03/2020) (QOF)	77.8%	81.5%	81.4%	No statistical variation
PCA rate (number of PCAs).	7.7% (3)	8.9%	8.0%	N/A

*PCA:. Personalised Care Adjustments allow practices to remove a patient from the indicator for limited, specified reasons.

Any additional evidence or comments

Patients who had personal care adjustments were removed from the indicator appropriately, for example, when patients declined to be involved in reviews of their care after the practice had contacted them at

least three times. The practice continued to engage with these patients and encouraged them to attend reviews.

Monitoring care and treatment

The practice had a comprehensive programme of quality improvement activity and routinely reviewed the effectiveness and appropriateness of the care provided.

Indicator	Practice	England average
Overall QOF score (out of maximum 559)	541	533.9
Overall QOF score (as a percentage of maximum)	96.8%	95.5%
Overall QOF PCA reporting (all domains)	5.9%	5.9%

	Y/N/Partial
Clinicians took part in national and local quality improvement initiatives.	Y
The practice had a programme of targeted quality improvement and used information about care and treatment to make improvements.	Y
The practice regularly reviewed unplanned admissions and readmissions and took appropriate action.	Y

Examples of improvements demonstrated because of clinical audits or other improvement activity in past two years

An audit was carried out between May 2020 and May 2021 on intrauterine contraceptive device (IUD) insertions.

- Areas audited included ensuring past medical history had been recorded in a patient's records; current contraceptive use; provision of information to patients; and evidence that consent had been obtained.
- A total of 16 procedures were carried out during this time and were audited.
- Results showed that all patients who had an IUD fitted had been assessed appropriately; had a
 record of their past medical history; and consent had been obtained.
- The practice nurse who carried out these audits planned to repeat it again after another 12 months.

Audits undertaken by the practice pharmacist included one on moving patients from warfarin (a medicine used to prevent blood clots) to other medicines which were considered to provide a more stable treatment option for patients.

Effective staffing

The practice was able to demonstrate that staff had the skills, knowledge and experience to carry out their roles.

	Y/N/Partial
Staff had the skills, knowledge and experience to deliver effective care, support and treatment.	Y
The practice had a programme of learning and development.	Y
Staff had protected time for learning and development.	Y
There was an induction programme for new staff.	Y
Staff had access to regular appraisals, one to ones, coaching and mentoring, clinical supervision and revalidation. They were supported to meet the requirements of professional revalidation.	Y
The practice could demonstrate how they assured the competence of staff employed in advanced clinical practice, for example, nurses, paramedics, pharmacists and physician associates.	Y
There was a clear and appropriate approach for supporting and managing staff when their performance was poor or variable.	Y
Explanation of any answers and additional evidence:	

• The practice had a programme of training and development. However, training records were not presented in a format which showed clearly when training had been undertaken, the recommended frequency of training or the date for refresher training. The assistant practice manager was aware of this and was in the process of reviewing records to include this information.

Coordinating care and treatment

Staff worked together and with other organisations to deliver effective care and treatment.

Indicator	Y/N/Partial
Care was delivered and reviewed in a coordinated way when different teams, services or organisations were involved.	Y
Patients received consistent, coordinated, person-centred care when they moved between services.	Y

Helping patients to live healthier lives

Staff were consistent and proactive in helping patients to live healthier lives.

	Y/N/Partial
The practice identified patients who may need extra support and directed them to relevant services. This included patients in the last 12 months of their lives, patients at risk of developing a long-term condition and carers.	

Staff encouraged and supported patients to be involved in monitoring and managing their own health.	Y
Patients had access to appropriate health assessments and checks.	Y
Staff discussed changes to care or treatment with patients and their carers as necessary.	Y
The practice supported national priorities and initiatives to improve the population's health, for example, stop smoking campaigns and tackling obesity.	Y

Any additional evidence or comments

Information was available for patients to encourage them to adopt healthier lifestyles. This included antibiotic use and when they are needed. Information on cervical screening was displayed in patient toilets.

Consent to care and treatment

The practice always obtained consent to care and treatment in line with legislation and guidance.

	Y/N/Partial
Clinicians understood the requirements of legislation and guidance when considering consent and decision making. We saw that consent was documented.	Y
Clinicians supported patients to make decisions. Where appropriate, they assessed and recorded a patient's mental capacity to make a decision.	Y
Do Not Attempt Cardio Pulmonary Resuscitation (DNACPR) decisions were made in line with relevant legislation and were appropriate.	Y

Caring

Rating: Good

Ashington House Surgery was rated Good for the provision of Caring services as a result of our inspection in November 2019. In accordance with Care Quality Commission methodology, the ratings from our previous inspection for this key questions (caring) have been carried through to contribute to the overall rating for the practice.

Responsive

Rating: Good

Ashington House Surgery was rated Good for the provision of Responsive services as a result of our inspection in November 2019. In accordance with Care Quality Commission methodology, the ratings from our previous inspection for this key questions (responsive) have been carried through to contribute to the overall rating for the practice.

Well-led

Rating: Good

When we inspected Ashington House Surgery in November 2019, the service was rated as Requires Improvement for the well-led key question.

A requirement was made to improve systems for assessing, monitoring and managing risks to patient safety; ensuring risk registers and action plans were fully completed; and succession planning for the practice was communicated to all staff.

At this inspection we found that the shortfalls had been addressed and the requirement notice had been met.

Leadership capacity and capability

There was compassionate, inclusive and effective leadership at all levels. Leaders could demonstrate that they had the capacity and skills to deliver high quality sustainable care.

	Y/N/Partial
Leaders demonstrated that they understood the challenges to quality and sustainability.	Y
They had identified the actions necessary to address these challenges.	Y
Staff reported that leaders were visible and approachable.	Y
There was a leadership development programme, including a succession plan.	Y

Explanation of any answers and additional evidence:

- The practice had redesigned the role of salaried GPs to aid recruitment, for example by allocating 15-minute appointments and had become a training practice.
- Staff were able to work remotely if needed.
- A manager had been employed to run the COVID-19 vaccination programme organised by the Primary Care Network, which the practice were part of. This meant that staff could remain working in the practice.
- Bank staff were available if needed.

Vision and strategy

The practice had a clear vision and credible strategy to provide high quality sustainable care.

	Y/N/Partial
The practice had a clear vision and set of values that prioritised quality and sustainability.	Y
There was a realistic strategy to achieve their priorities.	Y
The vision, values and strategy were developed in collaboration with staff, patients and external partners.	Y
Staff knew and understood the vision, values and strategy and their role in achieving them.	Y

Progress against delivery of the strategy was monitored.	Y
--	---

Culture

The practice had a culture which drove high quality sustainable care.

Y/N/Partial
Υ
Y
Y
Y
Y
Y
Y
Y
Y

Examples of feedback from staff or other evidence about working at the practice

Source	Feedback
	Staff said they had the autonomy to be responsible for their work and plan accordingly. They said the practice had a 'no blame' culture and promoted learning from significant events. Staff said they were supported by leaders and would not hesitate to approach them for guidance and advice on patient care.

Governance arrangements

There were clear responsibilities, roles and systems of accountability to support good governance and management.

	Y/N/Partial
There were governance structures and systems which were regularly reviewed.	Y
Staff were clear about their roles and responsibilities.	Y
There were appropriate governance arrangements with third parties.	
Explanation of any answers and additional evidence:	

• Work was progressing on developing protocols for all staff to use to promote consistency of practice. For example, protocols for flu vaccinations; taking and recording a patient's blood pressure and undertaking an ECG. This work was being carried out in collaboration with staff who have the expertise and skills in the various areas.

• Complaints were handled in accordance with practice policy, however, not all responses had details of the Parliamentary Health Service Ombudsman included, for patients who were not satisfied with the response.

Managing risks, issues and performance

There were clear and effective processes for managing risks, issues and performance.

	Y/N/Partial
There were comprehensive assurance systems which were regularly reviewed and improved.	Y
There were processes to manage performance.	
There was a quality improvement programme in place.	Y
There were effective arrangements for identifying, managing and mitigating risks.	
A major incident plan was in place.	Y
Staff were trained in preparation for major incidents.	Y
When considering service developments or changes, the impact on quality and sustainability was assessed.	Y

The practice had systems in place to continue to deliver services, respond to risk and meet patients' needs during the pandemic

	Y/N/Partial
The practice had adapted how it offered appointments to meet the needs of patients during the pandemic.	Y
The needs of vulnerable people (including those who might be digitally excluded) had been considered in relation to access.	
There were systems in place to identify and manage patients who needed a face-to-face appointment.	Y
The practice actively monitored the quality of access and made improvements in response to findings.	
There were recovery plans in place to manage backlogs of activity and delays to treatment.	
Changes had been made to infection control arrangements to protect staff and patients using the service.	
Staff were supported to work remotely where applicable.	Y

Appropriate and accurate information

There was a demonstrated commitment to using data and information proactively to drive and support decision making.

	Y/N/Partial
Staff used data to monitor and improve performance.	Y
Performance information was used to hold staff and management to account.	Y
There were effective arrangements for identifying, managing and mitigating risks.	Y
Staff whose responsibilities included making statutory notifications understood what this entails.	Y

Governance and oversight of remote services

	Y/N/Partial
The practice used digital services securely and effectively and conformed to relevant digital and information security standards.	Υ
The provider was registered as a data controller with the Information Commissioner's Office.	
Patient records were held in line with guidance and requirements.	Y
Patients were informed and consent obtained if interactions were recorded.	Y
The practice ensured patients were informed how their records were stored and managed.	Y
Patients were made aware of the information sharing protocol before online services were delivered.	
The practice had arrangements to make staff and patients aware of privacy settings on video and voice call services.	Y
Online consultations took place in appropriate environments to ensure confidentiality.	Y
The practice advised patients on how to protect their online information.	

Engagement with patients, the public, staff and external partners

The practice involved the public, staff and external partners to sustain high quality and sustainable care.

	Y/N/Partial
Patient views were acted on to improve services and culture.	Y
The practice had an active Patient Participation Group.	Y
Staff views were reflected in the planning and delivery of services.	Y
The practice worked with stakeholders to build a shared view of challenges and of the needs of the population.	Y
Explanation of any answers and additional evidence:	<u>.</u>

- The practice had increased the number of reception staff in response to patient complaints about poor access.
- In response to national patient survey results the practice have moved to a new telephone system to improve call quality and ensure there were not limitations to telephone lines. Other areas which the practice had improved included, providing active signposting training for reception staff to ensure patients were directed to the most appropriate person or service for care and treatment.
- In addition, work was continuing on promoting continuity of care, which had included increasing the number of salaried GPs available to reduce the number of appointments carried out by locum GPs.

Continuous improvement and innovation

There were systems and processes for learning, continuous improvement and innovation.

	Y/N/Partial
There was a strong focus on continuous learning and improvement.	Y
Learning was shared effectively and used to make improvements.	Y

Examples of continuous learning and improvement

The surgery is a member of the Brunel Healthcare Group, which consists of 12 GP practices and is part of a primary care network alongside two other practices.

The primary care network the practice was part of had jointly employed a paramedic to provide a home visiting service, a primary care pharmacist to improve the quality and safety of prescribing, and also a health and wellbeing advisor to coordinate social initiatives and advise individual patients on how they can improve their health through lifestyle changes.

The Brunel Healthcare Group worked closely together with a range of local providers to offer more personalised and coordinated health and social care as well as support for the wellbeing of local populations. This included provision of a dedicated 'hot hub' for potential COVID-19patient assessment. (A 'hot hub' is a designated setting to see patients who may be infectious and reduces the risk of cross infection).

Notes: CQC GP Insight

GP Insight assesses a practice's data against all the other practices in England. We assess relative performance for the majority of indicators using a "z-score" (this tells us the number of standard deviations from the mean the data point is), giving us a statistical measurement of a practice's performance in relation to the England average. We highlight practices which significantly vary from the England average (in either a positive or negative direction). We consider that z-scores which are higher than +2 or lower than -2 are at significant levels, warranting further enquiry. Using this technique we can be 95% confident that the practices performance is genuinely different from the average. It is important to note that a number of factors can affect the Z score for a practice, for example a small denominator or the distribution of the data. This means that there will be cases where a practice's data looks quite different to the average, but still shows as no statistical variation, as we do not have enough confidence that the difference is genuine. There may also be cases where a practice's data looks similar across two indicators, but they are in different variation bands.

The percentage of practices which show variation depends on the distribution of the data for each indicator, but is typically around 10-15% of practices. The practices which are not showing significant statistical variation are labelled as no statistical variation to other practices.

N.B. Not all indicators in the evidence table are part of the GP insight set and those that aren't will not have a variation band.

The following language is used for showing variation:

Variation Bands	Z-score threshold	
Significant variation (positive)	≤-3	
Variation (positive)	>-3 and ≤-2	
Tending towards variation (positive)	>-2 and ≤-1.5	
No statistical variation	<1.5 and >-1.5	
Tending towards variation (negative)	≥1.5 and <2	
Variation (negative)	≥2 and <3	
Significant variation (negative)	≥3	

Note: for the following indicators the variation bands are different:

- Child Immunisation indicators. These are scored against the World Health Organisation target of 95% rather than the England average. Note that
 practices that have "Met 90% minimum" have not met the WHO target of 95%.
- The percentage of respondents to the GP patient survey who responded positively to how easy it was to get through to someone at their GP practice
 on the phone uses a rules based approach for scoring, due to the distribution of the data. This indicator does not have a CCG average.
- The percentage of women eligible for cervical cancer screening at a given point in time who were screened adequately within a specified period (within 3.5 years for women aged 25 to 49, and within 5.5 years for women aged 50 to 64). This indicator does not have a CCG average and is scored against the national target of 80%.

It is important to note that z-scores are not a judgement in themselves, but will prompt further enquiry, as part of our ongoing monitoring of GP practices.

Guidance and Frequently Asked Questions on GP Insight can be found on the following link: <u>https://www.cqc.org.uk/guidance-providers/gps/how-we-monitor-gp-practices</u>

Note: The CQC GP Evidence Table uses the most recent validated and publicly available data. In some cases at the time of inspection this data may be relatively old. If during the inspection the practice has provided any more recent data, this can be considered by the inspector. However, it should be noted that any data provided by the practice will be unvalidated and is not directly comparable to the published data. This has been taken into account during the inspection process.

Glossary of terms used in the data.

- **COPD**: Chronic Obstructive Pulmonary Disease.
- PHE: Public Health England.
- QOF: Quality and Outcomes Framework.
- **STAR-PU**: Specific Therapeutic Group Age-sex weightings Related Prescribing Units. These weighting allow more accurate and meaningful comparisons within a specific therapeutic group by taking into account the types of people who will be receiving that treatment.
- *PCA: Personalised Care Adjustment. This replaces the QOF Exceptions previously used in the Evidence Table (see <u>GMS QOF Framework</u>). Personalised Care Adjustments allow practices to remove a patient from the indicator for limited, specified reasons.
- ٠
- ‰ = per thousand.