

Care Quality Commission

Inspection Evidence Table

The Practice Beacon (1-626549188)

Inspection date: 5 February 2019

Date of data download: 01 February 2019

Overall rating: Good

Please note: Any Quality Outcomes Framework (QOF) data relates to 2017/18.

Safe

Rating: Good

Risks to patients

There were adequate systems to assess, monitor and manage risks to patient safety.

	Y/N/Partial
There was an effective approach to managing staff absences and busy periods.	Yes
There was an effective induction system for temporary staff tailored to their role.	Yes
Comprehensive risk assessments were carried out for patients.	Yes
Risk management plans for patients were developed in line with national guidance.	Yes
Panic alarms were fitted and administrative staff understood how to respond to the alarm and the location of emergency equipment.	Yes
Clinicians knew how to identify and manage patients with severe infections including sepsis.	Yes
Receptionists were aware of actions to take if they encountered a deteriorating or acutely unwell patient and had been given guidance on identifying such patients.	Yes
There was a process in the practice for urgent clinical review of such patients.	Yes
There was equipment available to enable assessment of patients with presumed sepsis or other clinical emergency.	Yes
There were systems to enable the assessment of patients with presumed sepsis in line with National Institute for Health and Care Excellence (NICE) guidance.	Yes
When there were changes to services or staff the practice assessed and monitored the impact on safety.	Yes
Explanation of any answers and additional evidence: The practice could demonstrate continuous monitoring of its emergency equipment since November 2018. It had recently identified some equipment (for example, defibrillator pads) that required replacement and showed us evidence that these had been ordered.	

Information to deliver safe care and treatment

Staff had the information they needed to deliver safe care and treatment.

	Y/N/Partial
Individual care records, including clinical data, were written and managed securely and in line with current guidance and relevant legislation.	Yes
There was a system for processing information relating to new patients including the summarising of new patient notes.	Yes
There were systems for sharing information with staff and other agencies to enable them to deliver safe care and treatment.	Yes
Referral letters contained specific information to allow appropriate and timely referrals.	Yes
Referrals to specialist services were documented.	Yes
There was a documented approach to the management of test results and this was managed in a timely manner.	Yes
The practice demonstrated that when patients use multiple services, all the information needed for their ongoing care was shared appropriately and in line with relevant protocols.	Yes

Appropriate and safe use of medicines

The practice had systems for the appropriate and safe use of medicines, including medicines optimisation

Indicator	Practice	CCG average	England average	England comparison
Number of antibacterial prescription items prescribed per Specific Therapeutic group Age-sex Related Prescribing Unit (STAR PU) (01/10/2017 to 30/09/2018) (NHS Business Service Authority - NHSBSA)	0.45	0.55	0.94	Significant Variation (positive)
The number of prescription items for co-amoxiclav, cephalosporins and quinolones as a percentage of the total number of prescription items for selected antibacterial drugs (BNF 5.1 sub-set). (01/10/2017 to 30/09/2018) (NHSBSA)	8.3%	10.7%	8.7%	No statistical variation
Average daily quantity per item for Nitrofurantoin 50 mg tablets and capsules, Nitrofurantoin 100 mg m/r capsules, Pivmecillinam 200 mg tablets and Trimethoprim 200 mg tablets prescribed for uncomplicated urinary tract infection (01/04/2018 to 30/09/2018) (NHSBSA)	5.84	5.65	5.64	No statistical variation
Average daily quantity of oral NSAIDs prescribed per Specific Therapeutic Group Age-sex Related Prescribing Unit (STAR-PU) (01/04/2018 to 30/09/2018) (NHSBSA)	0.70	0.86	2.22	Significant Variation (positive)

Medicines management	Y/N/Partial
The practice ensured medicines were stored safely and securely with access restricted to authorised staff.	Yes
Blank prescriptions were kept securely and their use monitored in line with national guidance.	Yes
Staff had the appropriate authorisations to administer medicines (including Patient Group Directions or Patient Specific Directions).	Yes
The practice could demonstrate the prescribing competence of non-medical prescribers, and there was regular review of their prescribing practice supported by clinical supervision or peer review.	Yes
There was a process for the safe handling of requests for repeat medicines and evidence of structured medicines reviews for patients on repeat medicines.	Yes
The practice had a process and clear audit trail for the management of information about changes to a patient's medicines including changes made by other services.	Yes
There was a process for monitoring patients' health in relation to the use of medicines including high risk medicines (for example, warfarin, methotrexate and lithium) with appropriate monitoring and clinical review prior to prescribing.	Yes
The practice monitored the prescribing of controlled drugs. (For example, investigation of unusual prescribing, quantities, dose, formulations and strength).	Yes
There were arrangements for raising concerns around controlled drugs with the NHS England Area Team Controlled Drugs Accountable Officer.	Yes
The practice had taken steps to ensure appropriate antimicrobial use to optimise patient outcomes and reduce the risk of adverse events and antimicrobial resistance.	Yes
The practice held appropriate emergency medicines, risk assessments were in place to determine the range of medicines held, and a system was in place to monitor stock levels and expiry dates.	Partial
The practice had arrangements to monitor the stock levels and expiry dates of emergency medicines/medical gases.	Yes
There was medical oxygen and a defibrillator on site and systems to ensure these were regularly checked and fit for use.	Yes
Vaccines were appropriately stored, monitored and transported in line with PHE guidance to ensure they remained safe and effective.	Yes
<p>Explanation of any answers and additional evidence:</p> <p>The practice maintained a stock of emergency medicines. This did not include any emergency medicine for treating croup. We were told this was under current consideration and risk assessment. The practice had risk assessed whether there was a need to hold opioid painkillers and did not stock these on the premises. The practice could demonstrate continuous monitoring of its stock of emergency medicines since November 2018.</p>	

Effective

Rating: Good

Effective needs assessment, care and treatment

Patients' needs were assessed, and care and treatment was delivered in line with current legislation, standards and evidence-based guidance supported by clear pathways and tools.

	Y/N/Partial
The practice had systems and processes to keep clinicians up to date with current evidence-based practice.	Yes
Patients' immediate and ongoing needs were fully assessed. This included their clinical needs and their mental and physical wellbeing.	Yes
We saw no evidence of discrimination when staff made care and treatment decisions.	Yes
Patients' treatment was regularly reviewed and updated.	Yes
There were appropriate referral pathways were in place to make sure that patients' needs were addressed.	Yes
Patients were told when they needed to seek further help and what to do if their condition deteriorated.	Yes

Prescribing	Practice performance	CCG average	England average	England comparison
Average daily quantity of Hypnotics prescribed per Specific Therapeutic group Age-sex Related Prescribing Unit (STAR PU) (01/10/2017 to 30/09/2018) <small>(NHSBSA)</small>	0.59	0.89	0.81	No statistical variation

People with long-term conditions

Population group rating: Good

Findings

- Patients with long-term conditions had a structured annual review to check their health and medicines needs were being met. For patients with the most complex needs, the GP worked with other health and care professionals to deliver a coordinated package of care.
- Staff who were responsible for reviews of patients with long-term conditions had received specific training.
- The practice could demonstrate how they identified patients with commonly undiagnosed conditions, for example diabetes, chronic obstructive pulmonary disease (COPD), atrial fibrillation and hypertension.
- Adults with newly diagnosed cardio-vascular disease were offered statins.
- Patients with atrial fibrillation were assessed for stroke risk and treated appropriately.

Diabetes Indicators	Practice	CCG average	England average	England comparison
The percentage of patients with diabetes, on the register, in whom the last IFCC-HbA1c is 64 mmol/mol or less in the preceding 12 months (01/04/2017 to 31/03/2018) <small>(QOF)</small>	82.4%	78.6%	78.8%	No statistical variation
Exception rate (number of exceptions).	7.6% (7)	11.1%	13.2%	N/A
The percentage of patients with diabetes, on the register, in whom the last blood pressure reading (measured in the preceding 12 months) is 140/80 mmHg or less (01/04/2017 to 31/03/2018) <small>(QOF)</small>	70.7%	77.4%	77.7%	No statistical variation
Exception rate (number of exceptions).	10.9% (10)	9.4%	9.8%	N/A

	Practice	CCG average	England average	England comparison
The percentage of patients with diabetes, on the register, whose last measured total cholesterol (measured within the preceding 12 months) is 5 mmol/l or less (01/04/2017 to 31/03/2018) <small>(QOF)</small>	65.0%	80.1%	80.1%	Variation (negative)
Exception rate (number of exceptions).	13.0% (12)	11.0%	13.5%	N/A

Other long-term conditions	Practice	CCG average	England average	England comparison
The percentage of patients with asthma, on the register, who have had an asthma review in the preceding 12 months that includes an assessment of asthma control using the 3 RCP questions, NICE 2011 menu ID: NM23 (01/04/2017 to 31/03/2018) <small>(QOF)</small>	72.7%	79.0%	76.0%	No statistical variation
Exception rate (number of exceptions).	7.2% (6)	4.7%	7.7%	N/A
The percentage of patients with COPD who have had a review, undertaken by a healthcare professional, including an assessment of breathlessness using the Medical Research Council dyspnoea scale in the preceding 12 months (01/04/2017 to 31/03/2018) <small>(QOF)</small>	96.3%	89.3%	89.7%	No statistical variation
Exception rate (number of exceptions).	22.9% (8)	10.2%	11.5%	N/A

Indicator	Practice	CCG average	England average	England comparison
The percentage of patients with hypertension in whom the last blood pressure reading measured in the preceding 12 months is 150/90mmHg or less (01/04/2017 to 31/03/2018) <small>(QOF)</small>	75.4%	80.0%	82.6%	No statistical variation
Exception rate (number of exceptions).	4.5% (11)	4.4%	4.2%	N/A
In those patients with atrial fibrillation with a record of a CHA2DS2-VASc score of 2 or more, the percentage of patients who are currently treated with anti-coagulation drug therapy (01/04/2017 to 31/03/2018) <small>(QOF)</small>	100.0%	87.5%	90.0%	Variation (positive)
Exception rate (number of exceptions).	4.5% (1)	6.7%	6.7%	N/A

Any additional evidence or comments

The practice had recently assigned the advanced nurse practitioner as the lead for COPD who was carrying out patient reviews. The practice had a call/recall system for carrying out reviews and current data (unvalidated) suggested that the exception rate had fallen in 2018/19.

Families, children and young people

Population group rating: Good

Findings

- Childhood immunisation uptake rates had fallen below the World Health Organisation (WHO) targets since our previous inspection.
- The practice was aware of this and had identified improvement as a priority. The practice had a practice nurse vacancy but had recently recruited a locum nurse whose role included encouraging parents to ensure their child attended for immunisation.
- The practice had arrangements for following up failed attendance of children's appointments following an appointment in secondary care or for immunisation and liaised with health visitors when necessary.

Child Immunisation	Numerator	Denominator	Practice %	Comparison to WHO target
The percentage of children aged 1 who have completed a primary course of immunisation for Diphtheria, Tetanus, Polio, Pertussis, Haemophilus influenza type b (Hib)(i.e. three doses of DTaP/IPV/Hib) (01/04/2017 to 31/03/2018) (NHS England)	18	22	81.8%	Below 90% minimum (variation negative)
The percentage of children aged 2 who have received their booster immunisation for Pneumococcal infection (i.e. received Pneumococcal booster) (PCV booster) (01/04/2017 to 31/03/2018) (NHS England)	12	17	70.6%	Below 80% (Significant variation negative)
The percentage of children aged 2 who have received their immunisation for Haemophilus influenza type b (Hib) and Meningitis C (MenC) (i.e. received Hib/MenC booster) (01/04/2017 to 31/03/2018) (NHS England)	12	17	70.6%	Below 80% (Significant variation negative)
The percentage of children aged 2 who have received immunisation for measles, mumps and rubella (one dose of MMR) (01/04/2017 to 31/03/2018) (NHS England)	12	17	70.6%	Below 80% (Significant variation negative)

Monitoring care and treatment

The practice was developing a comprehensive programme of quality improvement activity. The practice routinely reviewed the effectiveness and appropriateness of the care provided.

Indicator	Practice	CCG average	England average
Overall QOF score (out of maximum 559)	517.6	532.5	537.5
Overall QOF exception reporting (all domains)	9.8%	6.7%	5.8%

	Y/N/Partial
Clinicians took part in national and local quality improvement initiatives.	Yes
The practice had a comprehensive programme of quality improvement and used information about care and treatment to make improvements.	Partial

Examples of improvements demonstrated because of clinical audits or other improvement activity in past two years

- The practice had been actively involved in providing support to patients affected by the Grenfell fire disaster over the previous 18 months.
- The practice could not demonstrate that it had carried out additional completed two-cycle clinical audits since our previous inspection. However, it was actively developing its quality improvement programme. The new lead GP had introduced regular clinical meetings following their appointment and had identified priorities for audit and monitoring.
- The clinical pharmacist told us they planned to re-audit the prescribing of high risk medicines as their first piece of work at the practice.

Effective staffing

The practice was able to demonstrate that staff had the skills, knowledge and experience to carry out their roles.

	Y/N/Partial
Staff had the skills, knowledge and experience to deliver effective care, support and treatment. This included specific training for nurses on immunisation and on sample taking for the cervical screening programme.	Yes
The learning and development needs of staff were assessed.	Yes
The practice had a programme of learning and development.	Yes
Staff had protected time for learning and development.	Yes
There was an induction programme for new staff.	Yes
Induction included completion of the Care Certificate for Health Care Assistants employed since April 2015.	Yes
Staff had access to regular appraisals, one to ones, coaching and mentoring, clinical supervision and revalidation. They were supported to meet the requirements of professional revalidation.	Yes
The practice could demonstrate how they assured the competence of staff employed in advanced clinical practice, for example, nurses, paramedics, pharmacists and physician associates.	Yes
There was a clear and appropriate approach for supporting and managing staff when their performance was poor or variable.	Yes

Notes: CQC GP Insight

GP Insight assesses a practice's data against all the other practices in England. We assess relative performance for the majority of indicators using a "z-score" (this tells us the number of standard deviations from the mean the data point is), giving us a statistical measurement of a practice's performance in relation to the England average. We highlight practices which significantly vary from the England average (in either a positive or negative direction). We consider that z-scores which are higher than +2 or lower than -2 are at significant levels, warranting further enquiry. Using this technique we can be 95% confident that the practice's performance is genuinely different from the average. It is important to note that a number of factors can affect the Z score for a practice, for example a small denominator or the distribution of the data. This means that there will be cases where a practice's data looks quite different to the average, but still shows as no statistical variation, as we do not have enough confidence that the difference is genuine. There may also be cases where a practice's data looks similar across two indicators, but they are in different variation bands.

The percentage of practices which show variation depends on the distribution of the data for each indicator, but is typically around 10-15% of practices. The practices which are not showing significant statistical variation are labelled as no statistical variation to other practices.

N.B. Not all indicators in the evidence table are part of the GP insight set and those that aren't will not have a variation band.

The following language is used for showing variation:

	Variation Band	Z-score threshold
1	Significant variation (positive)	$Z \leq -3$
2	Variation (positive)	$-3 < Z \leq -2$
3	No statistical variation	$-2 < Z < 2$
4	Variation (negative)	$2 \leq Z < 3$
5	Significant variation (negative)	$Z \geq 3$
6	No data	Null

Note: for the following indicators the variation bands are different:

- Child Immunisation indicators. These are scored against the World Health Organisation target of 95% rather than the England average.
- The percentage of respondents to the GP patient survey who responded positively to how easy it was to get through to someone at their GP practice on the phone uses a rules based approach for scoring, due to the distribution of the data. This indicator does not have a CCG average.

It is important to note that z-scores are not a judgement in themselves, but will prompt further enquiry, as part of our ongoing monitoring of GP practices.

Guidance and Frequently Asked Questions on GP Insight can be found on the following link: <https://www.cqc.org.uk/guidance-providers/gps/how-we-monitor-gp-practices>

Glossary of terms used in the data.

- **COPD:** Chronic Obstructive Pulmonary Disease
- **PHE:** Public Health England
- **QOF:** Quality and Outcomes Framework

STAR-PU: Specific Therapeutic Group Age-sex weightings Related Prescribing Units. These weighting allow more accurate and meaningful comparisons within a specific therapeutic group by taking into account the types of people who will be receiving that treat