ChemRxiv[™]

Selective N-Terminal Cysteine Protein Modification with Cyclopropenones

Alena Istrate, Claudio D. Navo, Bárbara B. Sousa, Marta C. Marques, Michael Deery, Andrew Bond, Francisco Corzana, Gonzalo Jiménez-Osés, Gonçalo Bernardes

Submitted date: 26/08/2020 • Posted date: 27/08/2020

Licence: CC BY-NC-ND 4.0

Citation information: Istrate, Alena; Navo, Claudio D.; Sousa, Bárbara B.; Marques, Marta C.; Deery, Michael; Bond, Andrew; et al. (2020): Selective N-Terminal Cysteine Protein Modification with Cyclopropenones. ChemRxiv. Preprint. https://doi.org/10.26434/chemrxiv.12866873.v1

Protein conjugates are valuable tools to create therapeutics, such as antibody-drug conjugates, or to study biological processes. Despite a number of protein conjugation strategies having been developed over recent years, the ability to modify one specific amino acid on a protein in the presence of other side chains with similar reactivity remains a challenge. We used the reaction between a monosubstituted cyclopropenone (CPO) probe and the 1,2-aminothiol of an N-terminal cysteine to give a stable 1,4-thiazepa-5-none linkage under mild, biocompatible conditions. The method relies on the ability of cyclopropenones to ring-open after sequential sulfhydryl and α-amine conjugation to be truly specific for N-terminal cysteine. We show that our CPO probes selectively label N-terminal cysteine containing peptides and proteins even in the presence of internal, solvent-exposed cysteines, which can be subsequently modified by using conventional cysteine modification reagents. The ability to distinguish and specifically target N-terminal cysteine residues on a protein will facilitate the construction of elaborate multi-labelled bioconjugates.

File list (3)		
Manuscript.pdf (1.77 MiB)	view on ChemRxiv • download file	
graphical abstract.png (60.59 KiB)	view on ChemRxiv • download file	
Supporting Information.pdf (21.10 MiB)	view on ChemRxiv • download file	

Selective N-terminal cysteine protein modification with cyclopropenones

Alena Istrate,¹ Claudio D. Navo,² Barbara B. Sousa,³ Marta C. Marques,³ Michael J. Deery,⁴ Andrew D. Bond,¹ Francisco Corzana,⁵ Gonzalo Jiménez-Osés,^{*,2} and Gonçalo J. L. Bernardes^{*,1,3}

¹Department of Chemistry, University of Cambridge, Lensfield Road, CB2 1EW Cambridge, United Kingdom.

² Center for Cooperative Research in Biosciences (CIC bioGUNE), Basque Research and Technology Alliance (BRTA), Bizkaia Technology Park, Building 801A, 48160 Derio, Spain

³Instituto de Medicina Molecular, Faculdade de Medicina da Universidade de Lisboa, Av. Prof. Egas Moniz, 1649-028 Lisboa, Portugal.

⁴Cambridge Centre for Proteomics, Cambridge Systems Biology Centre, Department of Biochemistry, University of Cambridge, Tennis Court Road, Cambridge CB2 1QW, United Kingdom.

⁵Departamento de Química, Universidad de La Rioja, Centro de Investigación en Síntesis Química, 26006 Logroño, Spain.

*e-mail: <u>gb453@cam.ac.uk</u> (G.J.L.B.) and <u>gjoses@cicbiogune.es</u> (G.J.-O.)

ABSTRACT

Protein conjugates are valuable tools to create therapeutics, such as antibody-drug conjugates, or to study biological processes. Despite a number of protein conjugation strategies having been developed over recent years, the ability to modify one specific amino acid on a protein in the presence of other side chains with similar reactivity remains a challenge. We used the reaction between a monosubstituted cyclopropenone (CPO) probe and the 1,2-aminothiol of an N-terminal cysteine to give a stable 1,4-thiazepa-5-none linkage under mild, biocompatible conditions. The method relies on the ability of cyclopropenones to ring-open after sequential sulfhydryl and α -amine conjugation to be truly specific for N-terminal cysteine. We show that our CPO probes selectively label N-terminal cysteine containing peptides and proteins

even in the presence of internal, solvent-exposed cysteines, which can be subsequently modified by using conventional cysteine modification reagents. The ability to distinguish and specifically target N-terminal cysteine residues on a protein will facilitate the construction of elaborate multi-labelled bioconjugates.

INTRODUCTION

Protein conjugates are important tools to create valuable therapeutics, such as antibody-drug conjugates (ADCs)¹⁻² and PEGylated proteins,³ build new functionalised materials,⁴ and study biological processes.⁵⁻⁶ Among various strategies for protein conjugation, the modification of naturally occurring amino acids remains the method of choice because it offers the advantage of straightforward accessibility without the need for sequence alterations by means of genetic methods.⁷ Ideally, conjugation reactions should proceed with complete chemo- and regioselectivity to generate welldefined protein constructs, which is a crucial requirement for many applications such as ADCs.¹⁻² Similarly, such transformations should occur rapidly in mild aqueous solutions at room temperature and physiological pH. Although many protein conjugation strategies have been developed over recent years, the ability to modify one specific amino acid on a protein in the presence of other side chains with similar reactivity remains a challenge.⁷ Protection of particularly reactive amino acids such as cysteine⁸ or extensive sequence engineering with the introduction of specific tags for enhanced reactivity⁹ is often required in order to achieve selectivity for the intended residue.

Among proteinogenic amino acids, lysine¹⁰⁻¹¹ and cysteine^{9, 12-13} are the two most commonly targeted for bioconjugation because they are highly nucleophilic under physiological conditions. Native lysine residues are very convenient targets, but they are abundant on protein surfaces, so it is difficult to achieve a high degree of selectivity.¹⁰ Conversely, cysteines are less abundant in proteins (<2%) and commonly less solvent-exposed, which makes them an excellent target for site-selective conjugation.¹⁴ However, when relying on cysteine for protein modification, there are several factors that must be taken into account. Specifically, cysteines often form disulfide bonds that are critical for a protein's structure, and modification of such residues can lead to loss of protein function. Moreover, many surface-exposed endogenous cysteines are directly involved in the catalytic activity of the protein, and

2

thus cannot be exploited for modification. Therefore, methods for targeted, selective cysteine modification could further advance the field of protein conjugation and enable the construction of functional and well-defined biomolecule conjugates without the need for genetic encoding of unnatural amino acids.

The most reliable strategy to differentiate one cysteine in the presence of other thiol groups is to target the N-terminal cysteine residue (NCys). Several methods for selective NCys modification have been developed, including reaction with thioesters [native chemical ligation (NCL)], condensation with aromatic aldehydes, and condensation with 2-cyanobenzothiazole derivatives (Figure 1).¹⁵ NCL provides a means to link protein or peptide fragments in a robust and chemoselective manner through trans-thioesterification and S-to-N acyl exchange (Figure 1a).¹⁶⁻¹⁷ However, this method is rarely used to produce protein conjugates because of difficulties in preparation and lack of stability of suitable thioester reagents. The reaction of 1,2aminothiols with aldehydes to form thiazolidine has also been explored as a strategy for NCys modification (Figure 1b).¹⁸ Neri and co-workers have successfully applied this approach for site-specific coupling of cytotoxic aldehydes to tumor-targeting antibodies, which produced homogenous conjugates that were then used for targeted delivery and slow release of the cytotoxic component.¹⁹ However, this reaction requires long incubation times (\approx 48 h), occurs under acidic conditions (pH 4–5) and is typically performed with a large excess of the aldehyde derivative. These limitations can be addressed by using formyl benzeno boronic acids (FBBA) that stabilize thiazolidine formation through $N \rightarrow B$ coordination (Figure 1c). Recently, FBBA reagents have been used to rapidly modify several model NCys-containing peptides at neutral pH.²⁰⁻²¹ This reaction is reversible and the product is not stable in an acidic environment (pH < 6) or in the presence of nucleophiles (e.g., free cysteine or benzyl hydroxylamine). It is however possible to use a thiazolidino boronate intermediate which undergoes an intramolecular acyl transfer to give more stable N-acyl thiazolidines.²²

Another N-terminal Cys labelling reaction was inspired by the final step of chemical synthesis of D-luciferin,²³ and is based on the condensation of free cysteine with 2-cyanobenzothiazol (CBT) (**Figure 1d**). After Rao and co-workers first demonstrated

the potential of this reaction for NCys modification,²⁴ the approach has been widely used in site-selective protein labelling and molecular imaging. This click reaction has major advantages for bioconjugation because of its efficiency, biocompatibility and stability of the resulting luciferin linkage.²⁵ However, 2-cyanobenzothiazol derivatives also react quickly (although reversibly) with simple thiols. As a result, when using excess CBT to ensure complete conjugation, other free-protein thiols must be protected. Therefore, alternative bioconjugation reagents for fast and selective labelling of N-terminal cysteines are still required to enable construction of complex protein conjugates of well-defined structures.

In this work, we explore the possibility of using cyclopropenone (CPO) for siteselective labelling of N-terminal cysteines on peptides and proteins. Cyclopropenone is a small unit with unique reactivity, which makes it an attractive component for several biological applications, for example as a key part of selective protease inhibitors²⁶ or reagents for bioorthogonal protein ligation.²⁷⁻²⁸ Indeed, the aromatic character of the ring²⁹ contributes to the cyclopropenone's stability, whereas significant angle strain and large dipole moment allow these moieties to participate in cycloaddition and ring-opening reactions.³⁰⁻³¹ As α , β -unsaturated ketones, cyclopropenones also act as electrophiles in 1,2- and 1,4-nucleophilic addition reactions. Several natural products contain cyclopropenone substituents, which highlights its biocompatibility and potential stability under physiological conditions.³²⁻ ³³ Previous work has demonstrated that CPOs react with thiols under basic conditions,²⁸ yet the reaction mechanism or products are not well established. Here, we describe the selective and efficient reaction of monosubstituted cyclopropenones with 1,2-aminothiol of the N-terminal cysteine and their use for site-selective bioconjugation of peptides and proteins (Figure 1e). Importantly, the use of cyclopropenones to efficiently modify an N-terminal cysteine in the presence of solvent-exposed cysteines either on the same sequence, or in a mixture of proteins is demonstrated for the first time.



Fig. 1 | Protein modification at N-terminal cysteine. a. Modification with thioesters through native chemical ligation; **b.** Condensation with aromatic aldehydes to form thiazolidines; **c.** Modification using formyl benzeno boronic acids; **d.** Condensation with 2-cyanobenzothiazole derivatives resulting in the formation of luciferin linkage; **e.** Reaction with monosubstituted cyclopropenones to generate 1,4-thiazepa-5-none linkage.

RESULTS & DISCUSSION

Monosubstituted cyclopropenones are efficient reagents for N-terminal cysteine labelling.

To start our investigation, we synthesized 2-phenethylcyclopropenone (1) as a model cyclopropenone and tested its stability in aqueous buffer solutions. Compound 1 was prepared from commercially available 4-phenyl-1-butyne by a simple, two-step procedure. Specifically, 4-phenyl-1-butyne was treated with difluorocarbene generated from TMSCF₃ to afford a *gem*-difluorocyclopropen derivative,³⁴ which was hydrolyzed on wet silica gel to afford desired cyclopropenone **1**.^{28, 35} Importantly, cyclopropenone 1 showed excellent stability after incubation in phosphate buffers (50 mM, pH 7–8) at 37 °C for 7 days (Figure S6 and S7). We next studied the reaction of cyclopropenone **1** with L-cysteine in the presence of base (Na₂CO₃) at 4 °C. The exceptionally high reactivity of cyclopropenone 1 required assaying the labelling reactions with small, highly accessible nucleophiles (i.e. amino acids, peptides, solvent-exposed sidechains of proteins) at lower temperatures than usual (4 °C). This allowed proper reaction monitoring and measuring labelling kinetics without interference of undesired side reactions. The reaction resulted in four isomeric compounds in 87% yield after for 30 min in aqueous solution. Liquid chromatography (LC)-mass spectrometry (MS) and NMR spectroscopic analysis showed that the products are two pairs of diastereomeric regioisomers of 1,4-thiazepa-5-none derivative, a stable 7-membered ring adduct (Figure 2a). Isomer 2a was isolated and characterised by X-ray crystallography, which revealed its stereochemistry and confirmed its structure (Figure 2b). To study the reaction further and confirm its outcome on a simpler, non-chiral model, we incubated cyclopropenone 1 with cysteamine (CA). As expected, the reaction yielded two pairs of enantiomeric regioisomers (Figure 2a) that were separated by column chromatography and characterized by NMR spectroscopy.

Cyclopropenone ring-opening after double nucleophilic attack makes the reaction with 1,2-aminothiols irreversible.

The whole reaction mechanism was inspected through quantum mechanical (QM) calculations using an abbreviated cyclopropenone model (1') and CA as reagents (**Figure 2c**). In the first reaction step, cysteamine thiolate undergoes conjugate addition to either the di-substituted (*pathway A*) or tri-substituted (*pathway B*) alkene

6

carbon atoms of the cyclopropenone. The initial activation barriers for these competitive pathways ultimately leading to the two observed regioisomeric products are quite similar and easily accessible at the reaction temperature despite the different steric hindrance at both positions ($\Delta G^{\ddagger}_{TS1 A}$ = 14.6 and $\Delta G^{\ddagger}_{TS1 B}$ = 15.6 kcal mol⁻¹, respectively), indicating that both pathways are feasible. Thio-enolate adducts are quite unstable (ΔG_{CP} ~11 kcal mol⁻¹), as commonly calculated for such intermediates in S-Michael-type reactions.³⁶ However, and opposing the common trend calculated for non-cyclopropenone electrophiles, protonation of the enolates does not stabilize the adducts ($\Delta G_{CP} = 7-8$ kcal mol⁻¹). As a consequence, this first step as well as the addition of simple thiolates (see below) are endergonic and reversible. Two racemic pairs of cyclopropanone diastereomers (**CP** A_{syn,anti}) are generated upon protonation of enolates in pathway A, and a racemic cyclopropanone regioisomer (CP B) in pathway B. Of note, transition states for spontaneous ring-opening leading to the ketene-ylide intermediates proposed to mediate in the bioorthogonal ligation of cyclopropenones assisted by triarylphosphines²⁷ could not be located from these thiocyclopropanone adducts. Conversely, intramolecular nucleophilic attack of the pendant amino group to the cyclopropanone carbonyl by the same face where the thioether is located proceeds with easily accessible activation barriers ($\Delta G^{\ddagger}_{TS2} = 14$ -17 kcal mol⁻¹) to give first zwitterionic (**HA**_{zw}) then neutral (**HA**) hemiaminal bicyclic adducts. As shown previously,³⁷ such transformations are strongly favored for cyclopropanones due to the significant strain release exerted by the carbonyl $sp^2 \rightarrow$ sp^3 rehybridization, despite the formation of 2-thia-5-azabicyclo[4.1.0]heptane structures. The OH-deprotonated form of these bicyclic intermediates (HA⁻) undergo ring-expansion reactions resembling those occurring fast in Favorskii rearrangements³⁷⁻³⁸ (ΔG^{\ddagger} TS3 = 6–8 kcal mol⁻¹). The resulting di-substituted carbanion in pathway A ($\Delta G_{TZ^{-}A} = -6$ kcal mol⁻¹), is more stable that the tri-substituted regioisomer generated in pathway B ($\Delta G_{TZ^-B} = -4$ kcal mol⁻¹), and negative hyperconjugation by neighboring sulfur atoms³⁹ can play an important role in the stabilization of both ring-expansion products. Finally, these species are protonated to irreversibly form the two experimentally observed 1,4-thiazepan-5-ones as racemic regioisomers with nearly identical energies ($\Delta G_{TZ} = -42 \text{ kcal mol}^{-1}$). Competitive but slower E₁cB-type elimination reactions were calculated from the same alkoxide intermediates (**HA**⁻) ($\Delta G^{\ddagger}_{TS4} = 10-15$ kcal mol⁻¹, **Figure S4**), in agreement with the observed preference for the ring-expansion reactions. In fact, this elimination mechanism would yield the type of α , β -unsaturated amides ((*Z*)- and (*E*)-crotonamides through pathway A and methacrylamide through pathway B) observed as ligation adducts with 1,2-aminophosphines, albeit through a different mechanism.²⁷

The activation barriers for the conjugate addition of 2-methylcyclopropenone **1'** and methanethiolate, an abbreviated model of internal cysteine residues and simple biological/unnatural thiols, are very similar to those calculated for the addition of cysteamine ($\Delta G^{\ddagger}_{TS1'}$ = 14–15 kcal mol⁻¹, **Figure S3**). However, the lack of adjacent free amino groups precludes cyclization and subsequent ring-expansion of the bicyclic intermediates and the associated ring strain release. Therefore, although internal cysteines and other thiols may compete with N-terminal cysteines for reacting with cyclopropenone derivatives, in the absence of such thermodynamic driving force the corresponding adducts would not be observed due to the reversibility of the intrinsically endergonic and reversible S-Michael reaction. Methylamine was also examined as a model nucleophile for the conjugate addition of cysteamine and N-terminal cysteine amino groups or lysine sidechains, to 2-methylcyclopropenone **1'**. The much higher activation barriers calculated for this reaction ($\Delta G^{\ddagger}_{TS1'N} \sim 20$ kcal mol⁻¹, **Figure S3b**) demonstrate the high preference for reactions with thiolates (and 1,2-aminothiolates) over primary amines.



Fig. 2 | a. Reaction of cyclopropenone **1** with L-cysteine ethyl ester hydrochloride and cysteamine; **b.** ORTEP plot (50% probability level) of compound **2a**. Hydrogen atoms are omitted for clarity; **c.** Proposed mechanism and minimum energy pathways for the reaction between 2-methylcyclopropenone (**1**') and cysteamine thiolate (**CA**) calculated with PCM(H₂O)/M06-2X/6-31+G(d,p). Note that each calculated structure represents a racemic compound constituted by two enantiomers with identical energies. *: protonation and deprotonation equilibria were calculated using tri-hydrated bicarbonate (HCO₃²⁻ · 3 H₂O) and carbonate (CO₃²⁻ · 3 H₂O) anions as an acid and a base, respectively. Given the intrinsic inaccuracy of such estimations, relative energies of charges/neutral species must be considered with caution. Relative free energies (Δ G) are given in kcal mol⁻¹. Numbers in brackets represent the relative

energies of transition structures. Three-dimensional structures for all lowest-energy stationary points are available in the **Supplementary Information**.

CPO-cysteine reaction is selective, and proceeds with fast kinetics in mild conditions.

Next, we examined the kinetics and selectivity of the CPO-cysteine reaction. To evaluate the reaction kinetics, we followed the HPLC chromatogram peak area of starting cyclopropenone 1 at different time points during the reaction (Figure S5). The second-order rate constant for this reaction was determined to be 3.0 M⁻¹s⁻¹ at 4 °C, which is comparable to the value reported for the CBT-cysteine reaction $(9.19 \text{ M}^{-1}\text{s}^{-1})$ and strain-promoted azide-alkyne cycloaddition reactions $(10^{-2} - 1 \text{ M}^{-1}\text{s}^{-1})$ performed at higher temperatures (37 °C).⁴⁰ In fact, the rate constant for the reaction between cyclopropenone **1** and cysteine extrapolated to 37 °C would be 67 M⁻¹ s⁻¹, which supersedes the aforementioned values. These results are in good agreement with the low activation barrier calculated for the reaction with cysteamine and confirm the high reactivity of monosubstituted cyclopropenones, making them ideal for protein modification with reduced amounts of labelling reagent and/or at particularly unreactive positions. Then, we tested the reactivity of probe **1** with other biological nucleophiles, including lysine, serine, threonine, tyrosine, glutathione, and cysteine methyl ester with tert-butoxycarbonyl protected amino group (Boc-Cys-OMe). As shown by LC–MS, compound 1 did not react with lysine, serine, threonine or tyrosine, but showed excellent selectivity for cysteine. In addition, these amino acids did not interfere with the reaction between CPO 1 and cysteine (Figure S8-S12). Boc-Cys-OMe reacted with compound **1** to form a complex mixture of products, although the reaction was slower relative to the one with cysteine that contains an unprotected amino group (Figure S14). In fact, when stoichiometric amounts of both cysteines (with free and Boc-protected amino groups) were treated with CPO probe 1, we observed high selectivity towards 1,2-aminothiol resulting in almost exclusive formation of 1,4-thiazepan-5-one products 2a-2d (Figure S14). Importantly, glutathione, the most abundant low molecular-weight thiol in cells, did not react with CPO probe 1, and did not interfere with the reaction between CPO 1 and cysteine (Figure S13), which confirms the computationally predicted high selectivity of cyclopropenones towards unprotected 1,2-aminothiols over simple thiols.

10

Cyclopropenone probes have a clear advantage over CBT probes that react with glutathione and other thiol nucleophiles²⁴ for the reaction with N-terminal cysteine.

Straightforward synthesis from easily accessible alkynes affords CPO probes with a range of payloads.

Before testing the applicability of the reaction in peptide and protein models, we designed and synthesised several cyclopropenone-based probes with different functionalities. We started with commercially available 5-hexynoic acid 4, which was converted into corresponding pentafluorophenol (PFP) ester 5. Ester 5 was subjected to cyclopropenation reaction to afford corresponding CPO-modified activated ester **CPO-PFP**. **CPO-PFP** is an easy-to-handle, stable (> 6 months) white solid that can be used to link the cyclopropenone unit to a molecule of interest through a primary amino group in a fast and efficient manner. With this intermediate, we prepared several labelling moieties of interest for peptide and protein bioconjugation. We optimised the reaction conditions by using a simple amine, benzylamine, to produce a model cvclopropenone with benzyl group (CPO-BN) in high yield. Next, we applied the optimised conditions to prepare cyclopropenone-based probes bearing polyethylene glycol (CPO-PEG), alkyne (CPO-PEG-Alkyne) and a fluorescent dye (CPO-EDANS). Overall, simple incubation of intermediate CPO-PFP with amine-bearing molecules of interest at room temperature for 20 min afforded CPO-based probes for protein and peptide labelling in high yields.



Fig. 3 | Synthesis of cyclopropenone-based probes with different functionalities.

CPO-probes selectively modify N-terminal cysteine on peptides, allowing dual cysteine labelling.

The high reactivity and selectivity of compound **1** towards 1,2-aminothiols prompted us to investigate the ability of cyclopropenone probes to modify N-terminal cysteine in peptides. Our investigation began with two unprotected 5-mer model peptides with Nterminal cysteines: CAIAI (**P1**) and CAIKI (**P2**). Notably, the **P2** contained a lysine residue to test the selectivity of the CPO reagent for cysteines over lysines. Treatment of both peptides (2 mM) with **CBO-BN** linker (2 equiv.) in NaP_i buffer (20 mM, pH 7)/acetonitrile resulted in complete conversion into the expected products after 1 h at 4 °C (**Figure 4a, S15**, and **S17**). Similarly, modification of peptides **P1** and **P2** with **CPO-PEG** linker resulted in formation of the expected PEGylated products, as confirmed by LC–MS (**Figure S16** and **S18**).

We then assayed the method with more complex peptides containing multiple nucleophilic side chains. To begin with, we focused on laminin-derived synthetic peptide **P3**, which inhibits tumor growth.⁴¹ This peptide contains nine amino acids, including N-terminal cysteine, tyrosine, serine, asparagine and arginine. Application of our optimized cysteine modification protocol resulted in full conversion of the starting peptide after 1 h, as confirmed by LC–MS (**Figure 4a** and **S19**). Then, we targeted the more challenging GTP-Binding Protein Fragment **P4**, which is a 16-mer peptide with multiple nucleophilic amino acids (one N-terminal cysteine, three lysines, two serines, one threonine, one methionine). The reaction of peptide **P4** with **CPO-BN** linker (2 equiv.) resulted in full conversion into products **P4-BNa** and **P4-BNb** within 1 h at 4 °C (**Figure 4a** and **S20**). No mass signal for double chemical modification was observed, which further highlights the chemoselectivity of the protocol.

As described above, N-terminal cysteine could be modified selectively even in the presence of other thiol nucleophiles. To test whether the same behaviour will be observed on the peptide level, we used **CPO-BN** to modify a peptide that contains both the N-terminal and internal cysteine residues. Vasopressin (**P5**), a 9-mer cyclic peptide, upon the reduction of the disulfide bridge generates cysteine residues in positions 1 and 6. In control experiments without the addition of the reducing agent, vasopressin showed no reaction with **CPO-BN** probe, or another common reagent for cysteine conjugation, *N*-methylmaleimide (NEM). We next reduced the disulfide bridge

with TCEP (Figure 4b and S21) and incubated the reduced peptide P5-SH with Nmethylmaleimide or CPO-BN probes. Cyclopropenones readily react with phosphines²⁷⁻²⁸ so care was taken to ensure that excess TCEP was removed from the reaction mixture before the addition of cyclopropenone (e.g., by using immobilised TCEP or limiting the amount used to 1 equiv.). As expected, NEM does not show any selectivity for the position of cysteine residue within the peptide, and incubation of P5-SH with NEM quickly resulted in modification of both cysteines (Figure S22). In contrast, incubation of the reduced vasopressin with CPO-BN probe resulted in selective modification of the N-terminal cysteine residue (Figure 4b and S24). The same result was obtained even if an excess of CPO probe (2 equiv.) was used. In further tests, we confirmed that the internal cysteine was still available for modification after selective labelling of the N-terminal cysteine with CPO-BN. Treatment of CPOmodified vasopressin with NEM allowed quantitative modification of the internal cysteine. In line with previous experiments, the internal cysteine could be efficiently conjugated to this reagent (Figure 4b and S25). LC-MS/MS analysis confirmed the sites of both modifications (Figure S26). Next, we examined whether CBT-based probes offer similar selectivity for the N-terminal cysteine of vasopressin. Under the conditions used for the reaction between CPO-BN and P5-SH, 2-cyano-6hydroxybenzothiazole (CBT) probe afforded a mixture of single- and double-modified peptides (Figure 4b and S23), which highlights the higher selectivity of the cyclopropenone-based reagents. This experiment demonstrates that CPO-labels can selectively modify N-terminal cysteine functionality even in the presence of other thiol groups, which can allow selective dual-cysteine labelling.



Fig. 4 | Chemo- and regioselective modification of 1,2-aminothiols on peptides. a. Site-selective modification of peptides **P1–P4** by **CPO-BN** reagent. Modification occurs only on N-terminal cysteine residue; **b. CPO-BN** probe selectively modifies N-terminal cysteine of vasopressin (**P5**) and leaves the internal cysteine unmodified and accessible for further functionalization.

CPO-probes react exclusively with 1,2-aminothiols on proteins.

To provide proof of concept for our CPO-reagent to be used for protein modification, we produced recombinant enhanced green fluorescent protein containing *N*-terminal

cysteine (**Cys-GFP**). Recombinant proteins with *N*-terminal cysteines are widely used for native chemical ligation technology, so various approaches have been developed for their direct production.⁴²⁻⁴⁴ To produce **Cys-GFP**, we engineered GFP protein variant with the tobacco etch virus (TEV) protease recognition sequence (ENLYFQ \downarrow C; arrow indicates the cleavage site) introduced after the hexa-His purification tag at the *N*-terminus. In this way, the cleavage of the expressed protein by TEV protease simultaneously removed the His-tag and generated GFP protein with the N-terminal cysteine. Incubation of Cys-GFP with CPO-BN (100 equiv.) in NaPi buffer (20 mM, pH 7.0) at room temperature for 2 h gave desired conjugate **GFP-CPO-BN** with high efficiency, as confirmed by LC-MS (Figure 5a, 5b, and S33). The high reactivity of GFP terminal cysteine required the use of DTT (25 equiv.) to avoid protein dimerization and thus an excess of cyclopropenone labelling reagent and a higher temperature compared to previous modification experiments was required in order to achieve complete conversion. Tryptic digestion of the conjugate and subsequent LC-MS/MS analysis confirmed the site of modification (Figure S36). Next, we evaluated the stability of GFP-CPO-BN conjugate with an excess of glutathione. After 24 h incubation at 37 °C with 5 mM glutathione, bioconjugate GFP-CPO-BN was shown to be stable with no de-conjugation observed (Figure S34 and S35). 0.5 µs Molecular dynamic simulations were performed on the four possible conjugates of GFP-CPO-BN (SI and Figure S49). The root-mean-square deviation values of the peptide backbone in all complexes range from 1.21 to 2.67 Å. Hence, these data suggest that the installation of CPO-BN scaffold does not cause relevant structural modifications on the protein, which is required for biological activity.

After demonstrating chemoselectivity and efficiency of **CPO-BN** for *N*-terminal cysteine bioconjugation, we expanded the scope of the reaction to other CPO-based linkers. Testing **CPO-PEG** and **CPO-EDANS** under the same reaction conditions as used for **CPO-BN** resulted in successful conversion of **Cys-GFP** into the expected products (**Figure 5c, 5d, S37** and **S38**). For **CPO-EDANS**, lower conversion (70%) is likely explained by the poor solubility of **CPO-EDANS**. We next assessed whether pre-purification of CPO-reagents is necessary for the conjugation reaction. For this purpose, we mixed **CPO-PFP** reagent with commercial amine-functionalised biotin to obtain **CPO-biotin** linker (**Figure S42**). Then, without any additional purification steps, the resulting solution was directly added to **Cys-GFP** (100 equiv. of **CPO-biotin** to 1

15

equiv. of **Cys-GFP**), and the reaction was monitored by LC–MS. To our delight, conversion of **Cys-GFP** into the expected conjugate was complete after incubation for 2 h (**Figure 5e** and **S36**). Thus, there is no need to purify the conjugation reagent before the labelling step, which is beneficial for the practical preparation of bioconjugates with payloads (eg. dyes, oligonucleotides, or drugs).

Notably, during our work on reaction optimization with **Cys-GFP**, addition of dithiothreitol (DTT) to protein stock solution was necessary to maintain the reactivity of cysteine residue, probably due to high susceptibility of the *N*-terminal cysteine to undergo oxidation and protein cross-linking. Nevertheless, an excess of DTT (500 equiv.) in the reaction mixture did not hinder the bioconjugation reaction of CPO probes with *N*-terminal cysteine. However, with similar NEM-based probes for **Cys-GFP** bioconjugation under the same reaction conditions, very low conversion rates were observed (0–15%; **Figure S39** and **S41**). This can be explained by the incompatibility of maleimides with excess DTT. Again, this highlights the selectivity of the cyclopropenone-based reagents towards 1,2-aminothiols for which 100 equiv. of the probe was enough to afford protein labelling with a range of probes even in the presence of excess of DTT (**Figure 5b** and **5e**).

Finally, we decided to explore whether CPO-based probes can selectively label a NCys-containing protein in the presence of other proteins bearing internal cysteines. We prepared a mixture of four proteins: **Cys-GFP** (3 free cysteines including N-terminal cysteine), engineered variant of the C2A domain of Synaptotagmin-I (**C2Am**, 1 free cysteine), Annexin V (**AnxV**, 1 free cysteine), engineered variant of a nanobody (**DesAB-HET**, 1 free cysteine), and incubated it with **CPO-biotin** linker or a similar maleimide-based linker (**MI-biotin**, **Figure S43**). **CPO-biotin** successfully conjugated to **Cys-GFP**, whereas **C2Am**, **AnxV**, and **DesAB-HET** proteins were unchanged. Similar results were observed regardless of whether or not DTT (500 equiv.) was present in the reaction mixture (**Figure 5f**, **S41** and **S43**). In contrast, **MI-biotin** probe fully modified 3 out of four proteins (**Cys-GFP**, **C2Am**, and **DesAB-HET**) when DTT was absent from the reaction mixture (**Figure 5f** and **S42**). **AnxV** was not modified because it usually requires larger excess of reagents, higher temperatures or longer reaction times for the cysteine modification to proceed.⁴⁵ As expected, in the presence of DTT, only minor modification of proteins (0–15%) with **MI-biotin** was observed



(**Figure 5f, S44**). Overall, these data confirm the high selectivity, orthogonality and highlight the suitability of CPO-reagents to label specific proteins in complex mixtures.

Fig. 5 | Chemo- and regioselective modification of 1,2-aminothiols on proteins. a. Site-selective bioconjugation of Cys-GFP with CPO-BN (b), CPO-PEG (c), CPO-EDANS (d), and CPO-biotin (e) probes; f. Modification of Cys-GFP, C2Am, AnxV, and DesAB-HET protein mixture (5 μ M each) with CPO-biotin or MI-biotin probes (50 equiv.) in NaP_i buffer (20 mM, pH 7.0) with or without DTT (500 equiv.).

CONCLUSIONS

We have developed an efficient method for N-terminal cysteine modification on peptide and proteins. The method is based on a reaction of monosubstituted cyclopropenones with 1,2-aminothiols, which results in formation of a stable 1,4thiazepa-5-none linkage. Synthesis of cyclopropenone-based conjugation reagents that bear various functional groups can be easily achieved in three steps starting from commercially available 5-hexynoic acid. The CPO-cysteine reaction proceeds with high efficiency in mild conditions (aqueous buffer, pH 7, 4–25 °C). The requirement for a nucleophile adjacent to the cysteine's thiol for the reaction to be irreversible, makes this transformation ideal for the selective labelling N-terminal cysteines. We have demonstrated that this approach can be used to modify N-terminal cysteine residues on a peptide or protein of interest even in the presence of remote nucleophilic residues, other biological thiols and reagents incompatible with common labelling reagents. Finally, we show the CPO-based probes can achieve selective labelling of NCys-containing protein in the presence of other proteins bearing internal cysteines. Further work is ongoing on the selective protein labelling in complex proteomes and in living cells. Overall, this ability to target specific cysteine residue on a protein will allow the straightforward construction of complex bioconjugates with well-defined structure, which makes this approach an important addition to the protein conjugation toolbox.

ASSOCIATED CONTENT

Supporting information

Detailed methods, characterization data and additional figures (PDF) CCDC 2010531 contains the supplementary crystallographic data for this paper

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENTS

This project has received funding from Fundação para a Ciência e a Tecnologia, Portugal (Investigator IF/00624/2015 to G.J.L.B), the Spanish Ministry of Science and Innovation (MCI) co-financed with FEDER funds (RTI2018-099592-B-C21 to F.C and RTI2018-099592-B-C22 to G.J.O) and Swiss National Science Foundation (Early Postdoc.Mobility P2BEP3_175253 to A.I). G.J.L.B. is a Royal Society University Research Fellow (URF\R\180019). The authors thank Dr Vikki Cantrill and Claudia Flandoli (draw.science) for their help with the editing of this manuscript and its figures, respectively. We also thank Prof. Michele Vendruscolo and Phil Lindstedt for the generous gift of DesAB-HET antibody, Dr. André Neves and Prof. Kevin Brindle for providing C2Am.

REFERENCES

1. Beck, A.; Goetsch, L.; Dumontet, C.; Corvaïa, N., Strategies and challenges for the next generation of antibody–drug conjugates. *Nat. Rev. Drug Discov.* **2017**, *16* (5), 315–337.

2. Chudasama, V.; Maruani, A.; Caddick, S., Recent advances in the construction of antibody–drug conjugates. *Nat. Chem.* **2016**, *8* (2), 114–119.

Pelegri-O'Day, E. M.; Lin, E.-W.; Maynard, H. D., Therapeutic Protein–
 Polymer Conjugates: Advancing Beyond PEGylation. *J. Am. Chem. Soc.* 2014, *136* (41), 14323–14332.

4. Spicer, C. D.; Pashuck, E. T.; Stevens, M. M., Achieving Controlled Biomolecule–Biomaterial Conjugation. *Chem. Rev.* **2018**, *118* (16), 7702–7743.

5. Zhang, G.; Zheng, S.; Liu, H.; Chen, P. R., Illuminating biological processes through site-specific protein labeling. *Chem. Soc. Rev.* **2015**, *44* (11), 3405–3417.

6. Xue, L.; Karpenko, I. A.; Hiblot, J.; Johnsson, K., Imaging and manipulating proteins in live cells through covalent labeling. *Nat. Chem. Biol.* **2015**, *11* (12), 917–923.

 Hoyt, E. A.; Cal, P. M. S. D.; Oliveira, B. L.; Bernardes, G. J. L., Contemporary approaches to site-selective protein modification. *Nat. Rev. Chem.* 2019, 3 (3), 147–171.

8. Taylor, M. T.; Nelson, J. E.; Suero, M. G.; Gaunt, M. J., A protein functionalization platform based on selective reactions at methionine residues. *Nature* **2018**, *562* (7728), *563*–*568*.

Zhang, C.; Welborn, M.; Zhu, T.; Yang, N. J.; Santos, M. S.; Van Voorhis, T.;
 Pentelute, B. L., π-Clamp-mediated cysteine conjugation. *Nat. Chem.* 2016, *8* (2), 120–128.

Matos, M. J.; Oliveira, B. L.; Martínez-Sáez, N.; Guerreiro, A.; Cal, P. M. S.
 D.; Bertoldo, J.; Maneiro, M.; Perkins, E.; Howard, J.; Deery, M. J.; Chalker, J. M.;
 Corzana, F.; Jiménez-Osés, G.; Bernardes, G. J. L., Chemo- and Regioselective

Lysine Modification on Native Proteins. *J. Am. Chem. Soc.* **2018**, *140* (11), 4004–4017.

11. Adusumalli, S. R.; Rawale, D. G.; Singh, U.; Tripathi, P.; Paul, R.; Kalra, N.; Mishra, R. K.; Shukla, S.; Rai, V., Single-Site Labeling of Native Proteins Enabled by a Chemoselective and Site-Selective Chemical Technology. *J. Am. Chem. Soc.* **2018**, *140* (44), 15114–15123.

Bernardim, B.; Cal, P. M. S. D.; Matos, M. J.; Oliveira, B. L.; Martínez-Sáez,
 N.; Albuquerque, I. S.; Perkins, E.; Corzana, F.; Burtoloso, A. C. B.; Jiménez-Osés,
 G.; Bernardes, G. J. L., Stoichiometric and irreversible cysteine-selective protein
 modification using carbonylacrylic reagents. *Nat. Commun.* 2016, 7 (1), 13128.

Vinogradova, E. V.; Zhang, C.; Spokoyny, A. M.; Pentelute, B. L.; Buchwald,
 S. L., Organometallic palladium reagents for cysteine bioconjugation. *Nature* 2015, 526 (7575), 687–691.

14. Chalker, J. M.; Bernardes, G. J. L.; Lin, Y. A.; Davis, B. G., Chemical Modification of Proteins at Cysteine: Opportunities in Chemistry and Biology. *Chem. Asian J.* **2009**, *4* (5), 630–640.

15. Rosen, C. B.; Francis, M. B., Targeting the N terminus for site-selective protein modification. *Nat. Chem. Biol.* **2017**, *13* (7), 697–705.

16. Dawson, P. E.; Muir, T. W.; Clark-Lewis, I.; Kent, S. B., Synthesis of proteins by native chemical ligation. *Science* **1994**, *266* (5186), 776.

17. Agouridas, V.; El Mahdi, O.; Diemer, V.; Cargoët, M.; Monbaliu, J.-C. M.; Melnyk, O., Native Chemical Ligation and Extended Methods: Mechanisms, Catalysis, Scope, and Limitations. *Chem. Rev.* **2019**, *119* (12), 7328–7443.

 Zhang, L.; Tam, J. P., Thiazolidine Formation as a General and Site-Specific Conjugation Method for Synthetic Peptides and Proteins. *Anal. Biochem.* **1996**, 233 (1), 87–93.

19. Casi, G.; Huguenin-Dezot, N.; Zuberbühler, K.; Scheuermann, J.; Neri, D., Site-Specific Traceless Coupling of Potent Cytotoxic Drugs to Recombinant Antibodies for Pharmacodelivery. *J. Am. Chem. Soc.* **2012**, *134* (13), 5887–5892.

20. Faustino, H.; Silva, M. J. S. A.; Veiros, L. F.; Bernardes, G. J. L.; Gois, P. M. P., Iminoboronates are efficient intermediates for selective, rapid and reversible N-terminal cysteine functionalisation. *Chem. Sci.* **2016**, *7* (8), 5052–5058.

Bandyopadhyay, A.; Cambray, S.; Gao, J., Fast and selective labeling of N-terminal cysteines at neutral pH via thiazolidino boronate formation. *Chem. Sci.* **2016**, 7 (7), 4589–4593.

22. Gao, J.; Li, K.; Wang, W., Fast and Stable N-Terminal Cysteine Modification via Thiazolidino Boronate Mediated Acyl Transfer. *Angew. Chem. Int. Ed.* **2020**, doi:10.1002/anie.202000837.

23. Podsiadły, R.; Grzelakowska, A.; Modrzejewska, J.; Siarkiewicz, P.; Słowiński,
D.; Szala, M.; Świerczyńska, M., Recent progress in the synthesis of firefly luciferin derivatives. *Dyes Pigm.* 2019, *170*, 107627.

24. Ren, H.; Xiao, F.; Zhan, K.; Kim, Y.-P.; Xie, H.; Xia, Z.; Rao, J., A Biocompatible Condensation Reaction for the Labeling of Terminal Cysteine Residues on Proteins. *Angew. Chem. Int. Ed.* **2009**, *48* (51), 9658–9662.

25. Yuan, Y.; Liang, G., A biocompatible, highly efficient click reaction and its applications. *Org. Biomol. Chem.* **2014,** *12* (6), 865–871.

Cohen, M.; Bretler, U.; Albeck, A., Peptidyl cyclopropenones: Reversible inhibitors, irreversible inhibitors, or substrates of cysteine proteases? *Protein Sci.* **2013**, 22 (6), 788–799.

27. Shih, H.-W.; Prescher, J. A., A Bioorthogonal Ligation of Cyclopropenones Mediated by Triarylphosphines. *J. Am. Chem. Soc.* **2015**, *137* (32), 10036–10039.

Row, R. D.; Shih, H.-W.; Alexander, A. T.; Mehl, R. A.; Prescher, J. A.,
 Cyclopropenones for Metabolic Targeting and Sequential Bioorthogonal Labeling. *J. Am. Chem. Soc.* 2017, 139 (21), 7370–7375.

29. Experiments show cyclopropenone is aromatic. *Chem. Eng. News* 1983, 61 (38), 33.

30. Potts, K. T.; Baum, J. S., Chemistry of cyclopropenones. *Chem. Rev.* **1974**, 74 (2), 189–213.

31. Krebs, A. W., Cyclopropenylium Compounds and Cyclopropenones. *Angew. Chem. Int. Ed.* **1965**, *4* (1), 10–22.

32. Kogen, H.; Kiho, T.; Tago, K.; Miyamoto, S.; Fujioka, T.; Otsuka, N.; Suzuki-Konagai, K.; Ogita, T., Alutacenoic Acids A and B, Rare Naturally Occurring Cyclopropenone Derivatives Isolated from Fungi: Potent Non-Peptide Factor XIIIa Inhibitors. *J. Am. Chem. Soc.* **2000**, *122* (8), 1842–1843. 33. Reber, K. P.; Gilbert, I. W.; Strassfeld, D. A.; Sorensen, E. J., Synthesis of (+)-Lineariifolianone and Related Cyclopropenone-Containing Sesquiterpenoids. *J. Org. Chem.* **2019**, *84* (9), 5524–5534.

34. Wang, F.; Luo, T.; Hu, J.; Wang, Y.; Krishnan, H. S.; Jog, P. V.; Ganesh, S. K.; Prakash, G. K. S.; Olah, G. A., Synthesis of gem-Difluorinated Cyclopropanes and Cyclopropenes: Trifluoromethyltrimethylsilane as a Difluorocarbene Source. *Angew. Chem. Int. Ed.* **2011**, *50* (31), 7153–7157.

35. Rullière, P.; Cyr, P.; Charette, A. B., Difluorocarbene Addition to Alkenes and Alkynes in Continuous Flow. *Org. Lett.* **2016**, *18* (9), 1988–1991.

36. Krenske, E. H.; Petter, R. C.; Houk, K. N., Kinetics and Thermodynamics of Reversible Thiol Additions to Mono- and Diactivated Michael Acceptors: Implications for the Design of Drugs That Bind Covalently to Cysteines. *J. Org. Chem.* **2016**, *81* (23), 11726–11733.

37. Salaun, J., Cyclopropanone hemiacetals. *Chem. Rev.* **1983**, *83* (6), 619–632.

38. Guijarro, D.; Yus, M., The Favorskii Rearrangement: Synthetic Applications. *Curr. Org. Chem.* **2005**, *9* (17), 1713–1735.

 Bernasconi, C. F.; Kittredge, K. W., Carbanion Stabilization by Adjacent Sulfur: Polarizability, Resonance, or Negative Hyperconjugation? Experimental Distinction Based on Intrinsic Rate Constants of Proton Transfer from (Phenylthio)nitromethane and 1-Nitro-2-phenylethane. *J. Org. Chem.* **1998**, *63* (6), 1944–1953.

40. Oliveira, B. L.; Guo, Z.; Bernardes, G. J. L., Inverse electron demand Diels– Alder reactions in chemical biology. *Chem. Soc. Rev.* **2017**, *46* (16), 4895–4950.

41. Sakamoto, N.; Iwahana, M.; Tanaka, N. G.; Osada, Y., Inhibition of Angiogenesis and Tumor Growth by a Synthetic Laminin Peptide, CDPGYIGSR-NH₂. *Cancer Res.* **1991**, *51* (3), 903–906.

42. Gentle, I. E.; De Souza, D. P.; Baca, M., Direct Production of Proteins with N-Terminal Cysteine for Site-Specific Conjugation. *Bioconjugate Chem.* **2004**, *15* (3), 658–663.

43. Tolbert, T. J.; Wong, C.-H., New Methods for Proteomic Research:
Preparation of Proteins with N-Terminal Cysteines for Labeling and Conjugation. *Angew. Chem. Int. Ed.* 2002, *41* (12), 2171–2174.

44. Liu, D.; Xu, R.; Dutta, K.; Cowburn, D., N-terminal cysteinyl proteins can be prepared using thrombin cleavage. *FEBS Lett.* **2008**, *582* (7), 1163–1167.

45. Cal, P. M. S. D.; Sieglitz, F.; Santos, F. M. F.; Parente Carvalho, C.; Guerreiro, A.; Bertoldo, J. B.; Pischel, U.; Gois, P. M. P.; Bernardes, G. J. L., Siteselective installation of BASHY fluorescent dyes to Annexin V for targeted detection of apoptotic cells. *Chem. Commun.* **2017**, *53* (2), 368–371.



graphical abstract.png (60.59 KiB)

view on ChemRxiv • download file

Supporting Information for

Selective N-terminal cysteine protein modification with cyclopropenones

Alena Istrate,¹ Claudio D. Navo,² Barbara B. Sousa,³ Marta C. Marques,³ Michael J. Deery,⁴ Andrew D. Bond,¹ Francisco Corzana,⁵ Gonzalo Jiménez-Osés,^{*,2} and Gonçalo J. L. Bernardes^{*,1,3}

¹Department of Chemistry, University of Cambridge, Lensfield Road, CB2 1EW Cambridge, United Kingdom.

²Center for Cooperative Research in Biosciences (CIC bioGUNE), Basque Research and Technology Alliance (BRTA), Bizkaia Technology Park, Building 801A, 48160 Derio, Spain

³Instituto de Medicina Molecular, Faculdade de Medicina da Universidade de Lisboa, Av. Prof. Egas Moniz, 1649-028 Lisboa, Portugal.

⁴Cambridge Centre for Proteomics, Cambridge Systems Biology Centre, Department of Biochemistry, University of Cambridge, Tennis Court Road, Cambridge CB2 1QW, United Kingdom.

⁵Departamento de Química, Universidad de La Rioja, Centro de Investigación en Síntesis Química, 26006 Logroño, Spain.

Correspondence should be sent to: gb453@cam.ac.uk (G.J.L.B.) and/or gjoses@cicbiogune.es (G.J.-O.)

Table of Contents

1.	List of abbreviations	.S2
2.	General methods	.S3
3.	Synthetic procedures	.S4
4.	X-ray structure analysis of 2a	.S13
5.	Quantum mechanical calculations	.S16
6.	Kinetic study for the reaction of cyclopropenone probe 1 with cysteine	.S29
7.	Stability studies of cyclopropenone probe 1 in buffer	.S31
8.	Stability of cyclopropenones towards nucleophiles	.S33
9.	Peptide modification	.S41
10.	Protein modification	.S54
11.	Molecular dynamics (MD) simulations	.S77
12.	NMR spectra	.S79
13.	References	.S91

1. List of abbreviations

CC	column chromatography
CPO-BN	4-(3'-oxocycloprop-1'-en-1'-yl)-N-phenylbutanamide
CPO-EDANS	Sodium 5-((2'-(4''-(3'''-oxocycloprop-1'''-en-1'''-
	yl)butanamido)ethyl)amino) naphthalene-1-sulfonate
CPO-PEG	N-(2-(2-(2-(2-hydroxyethoxy)ethoxy)ethoxy)ethyl)-4-(3'-oxocycloprop-1'-en-
	1'-yl)butanamide
CPO-PEG-Alkyne	4-(3'-oxocycloprop-1'-en-1'-yl)-N-(3'',6'',9'',12''-tetraoxapentadec-14''-yn-
	1''-yl)butanamide
CPO-PFP	perfluorophenyl 4-(3'-oxocycloprop-1'-en-1'-yl)butanoate
DIPEA	N,N-diisopropylethylamine
DMF	dimethylformamide
DTT	dithiothreitol
GSH	glutathione
HPLC	high-performance liquid chromatography
LC-MS	liquid chromatography-mass spectrometry
MD	molecular dynamics
MS	mass spectrometry
PE	petroleum ether
r.t.	room temperature
THF	tetrahydrofuran
TLC	thin layer chromatography
Tris HCl	tris(hydroxymethyl)aminomethane hydrochloride

2. General methods

Chemicals and characterization of the compounds

Chemicals were purchased and used without further purification. Anhydrous solvents for reactions were obtained by distillation over drying agents or purchased from Sigma Aldrich (DMF). All non-aqueous reactions were performed under an atmosphere of argon in oven-dried glassware. Argon gas was predried via passage through calcium chloride. Column chromatography (CC) was performed on silica gel (pore size 60 Å, 230-400 mesh particle size, Merck). All solvents for CC were of technical grade and distilled prior to use. Thin-layer chromatography (TLC) was performed on silica gel plates (TLC Silica gel 60 F254, Merck). Visualization was achieved either under UV light (254 nm) and/or by staining in dip solutions [Cer(IV)-sulfate (10.5 g), phosphormolybdenic acid (21 g), concentrated H₂SO₄ (60 ml), H₂O (900 ml) or KMnO₄(6 g), K₂CO₃ (40 g), 15% NaOH (3 ml) in H₂O (800 ml)] followed by heating with a heat gun. NMR spectra were recorded on a Bruker Avance III HD 400 MHz, Bruker Avance III HD 500 MHz spectrometers at 400 or 500 MHz (¹H), 100 or 125 MHz (¹³C), at 376.5 MHz (¹⁹F) of in either CDCl₃ or DMF- d_7 at room temperature. Chemical shifts (δ) are reported relative to the undeuterated residual solvent peak [CHCl₃: 7.26 ppm (¹H) and 77.16 ppm (¹³C); DMF-d₇: 2.92 ppm (¹H) and 163.15 ppm (¹³C)]. Signal assignments are based on DEPT or APT experiments, and on ¹H, ¹H and ¹H, ¹³C correlation experiments. Multiplicities are abbreviated as b (broad), s (singlet), d (doublet), t (triplet), q (quartet), m (multiplet) or combinations thereof. High-resolution mass spectra were performed by the Mass Spectrometry Service, Department of Chemistry, University of Cambridge using a Waters LCT Premier or a Waters Xevo G2-S spectrometer and ionized by ESI or ASAP.

LC-MS method for analysis

LC–MS was performed on a Waters SQ Detector 2 mass spectrometer coupled to an Acquity UPLC system using an Acquity UPLC BEH C18 column (130Å, 1.7 μ m, 2.1 mm X 50 mm). Solvents A, water with 0.1% formic acid and B, 71% acetonitrile, 29% water and 0.075% formic acid were used as the mobile phase at a flow rate of 0.2 mL min⁻¹. The gradient was programmed as follows: 100% A for 2 min, then 100% A to 100% B in 9 min, then 100% B for 5 min, and after that 100% A for 4 min. The electrospray source was operated with a capillary voltage of 3.0 kV and a cone voltage of 30 V. Nitrogen was used as the desolvation gas at a total flow of 800 L h⁻¹. *m/z* values of positively and negatively charged ions were measured in the mass analyzer, which was scanned between *m/z* 100-2000 for the generation of mass spectra. The major peak(s) were selected for integration and analyzed using MassLynx software (v. 4.1 from Waters).

3. Synthetic procedures

2-Phenethylcycloprop-2-en-1-one 1



In a glove box, sodium iodide (330 mg, 2.2 mmol, 2.2 equiv.) was added to an oven-dried flask equipped with a magnetic stir bar, after which the flask was closed with septa, transferred to a fume hood and filled with N₂. Next, dry THF (3 mL), starting alkyne (140 uL, 1 mmol, 1 equiv.), and TMSCF₃ (0.3 mL, 2 mmol, 2 equiv.) were added via syringe, and the reaction mixture was allowed to stir at r.t. for 24 h. Subsequently, the reaction mixture was poured into H₂O (10 ml) and extracted with Et₂O (3 x 10 ml). The combined organic phases were dried over MgSO₄, filtrated, evaporated, and the crude difluorocyclopropene intermediate was adsorbed on silica gel (1 g) and allowed to stand at r.t. overnight to generate the corresponding cyclopropenone product. Purification of the adsorbed material by CC (PE/EtOAc 2:1 \rightarrow EtOAc) afforded the product 1 (87 mg, 55%) as a dark-brown oil.

TLC (PE/EtOAc 2:3) $R_f = 0.17$.

¹H NMR (CDCl₃, 400 MHz): δ 8.34 (*s*, H-C(3)), 7.37 – 7.18 (*m*, 5H, H-arom), 3.09 – 2.98 (*m*, 4H, CH₂).

¹³C NMR (CDCl₃, 100 MHz): δ 169.47 (C(2)), 157.47 (C(1)), 149.24 (C(3)), 139.34 (C-arom), 128.94 (CH-arom), 128.41 (CH-arom), 126.96 (CH-arom), 31.89 (CH₂), 29.24 (CH₂).

ESI⁺-HRMS: calculated for $C_{11}H_{11}O([M+H]^+)$ 159.0810, found 159.0808.

Ethyl (3R,6R)-5-oxo-6-phenethyl-1,4-thiazepane-3-carboxylate 2a, ethyl (3R,6S)-5-oxo-6-phenethyl-1,4-thiazepane-3-carboxylate 2b, ethyl (3R,7R)-5-oxo-7-phenethyl-1,4-thiazepane-3-carboxylate 2c, ethyl (3R,7S)-5-oxo-7-phenethyl-1,4-thiazepane-3-carboxylate 2d



L-Cysteine ethyl ester hydrochloride (37.32 mg, 0.2 mmol, 1 equiv.) was dissolved in H₂O/CH₃CN (20 mL, 1/9) and cooled to 0°C. Next, Na₂CO₃ (21 mg, 0.4 mmol, 2 equiv.) and 2-phenethylcycloprop-2en-1-one (300 μ L of 0.67 M stock solution in CH₂Cl₂, 0.2 mmol of **1**, 1 equiv.), and the resulting solution was stirred for 30 min at 0°C. The reaction mixture was washed with 0.15 M HCl solution, and the aqueous phase was extracted with CH₂Cl₂ (3 x 10 mL). The combined organic phases were dried over MgSO₄, evaporated and the crude product was purified by CC (PE/ethyl acetate 8:2 \rightarrow 3:2) to give two fractions: **2a** (9.9 mg, 16%; 53.2 mg; 3R,6R-isomer) and **2b-d** (43.8 mg, 71%; mixture of three isomers) as white solids.

<u>2a</u> (3R,6R-isomer):

TLC (PE/EtOAc 2:3) $R_f = 0.64$.

¹H NMR (CDCl₃, 400 MHz): δ 7.32 – 7.26 (m, 2H, H-arom), 7.23 – 7.17 (m, 3H, H-arom), 6.58 (d, *J* = 6.1 Hz, 1H, NH), 4.49 (ddd, *J* = 9.4, 6.1, 1.5 Hz, 1H, H(3)), 4.28 (q, *J* = 7.1 Hz, 2H, CH₂-CH₃), 2.99 (d, *J* = 14.1, 1H, H(2a)), 2.86 – 2.70 (m, 4H, H(2b), H(6), H(7b), CH₂CH^aH^bPh), 2.66 – 2.57 (ddd, *J* = 13.7, 8.8, 6.4 Hz, 1H, CH₂CH^aH^bPh), 2.49 (d, *J* = 13.1, 1H, H(7a)), 2.34-2.23 (dddd, *J* = 13.9, 8.8, 7.6, 6.2 Hz, 1H, CH^aH^bCH₂Ph), 1.73 – 1.62 (m, 1H, CH^aH^bCH₂Ph), 1.32 (t, *J* = 7.1 Hz, 3H, CH₂-CH₃).

¹³C NMR (CDCl₃, 100 MHz): δ 175.73 (NHCO), 169.72 (COEt), 141.64 (C-arom), 128.65 (CH-arom), 128.58 (CH-arom), 126.13 (CH-arom), 62.89 (CH₂CH₃), 58.42 (C(3)), 47.78 (C(6)), 34.70 (C(2)), 33.77 (CH₂CH₂Ph), 33.63 (CH₂CH₂Ph), 30.45 (C(7)), 14.22 (CH₂CH₃).

 ESI^{+} -HRMS: calculated for $C_{16}H_{22}NO_{3}S$ ([M+H]⁺) 308.1320, found 308.1318.

<u>2b-d</u>:

TLC (PE/EtOAc 2:3) $R_f = 0.40$.

¹H NMR (CDCl₃, 400 MHz): δ 7.38 – 7.19 (m, 5H, H-arom), 6.62 – 6.46 (m, 1H), 4.53 – 4.44 (m, 1H), 4.36 – 4.25 (m, 2H), 3.27 – 2.58 (m, 7H), 2.35 – 1.89 (m, 2H), 1.42 – 1.29 (m, 3H).

¹H NMR and ¹³C NMR not assigned due to mixture of isomers.

ESI⁺-HRMS: calculated for $C_{16}H_{22}NO_3S$ ([M+H]⁺) 308.1320, found 308.1318.

6-phenethyl-1,4-thiazepan-5-one 3a,b (both enantiomers) and 7-phenethyl-1,4-thiazepan-5-one 3c,d (both enantiomers)



Cysteamine (22 mg, 0.29 mmol, 1.5 equiv.) was dissolved in H₂O/CH₃CN (90 mL, 1/1) and cooled to 0°C. Next, Na₂CO₃ (20 mg, 0.19 mmol, 1 equiv.) and 2-phenethylcycloprop-2-en-1-one (2.92 mL of 0.065 M stock solution in CH₃CN, 0.19 mmol of **1**, 1 equiv.), and the resulting solution was stirred for 1 h at 0°C. The reaction mixture was then concentrated to one-half volume and extracted with CH₂Cl₂ (3 x 20 mL). The combined organic phases were dried over MgSO₄, evaporated and the crude product was purified by CC (petrol/ethyl acetate 2:3 \rightarrow 2:6) to give products **3** (12.3 mg, 28%) and **3** (26.6 mg, 60%) as white solids.

<u>3a,b:</u>

TLC (PE/EtOAc 2:3) $R_f = 0.31$.

¹H NMR (CDCl₃, 400 MHz): δ 7.31 – 7.09 (*m*, 5H, H-arom), 6.14 (*s*, 1H, NH), 3.68 – 3.43 (*m*, 2H, H-C(3)), 2.87 (*dddd*, J = 9.7, 7.9, 6.1, 1.7 Hz, 1H, H-C(6)), 2.76 – 2.51 (*m*, 5H, CH₂CH₂Ph, 2xH-C(2), H-C(7)), 2.43 (*d*, J = 14.7 Hz, 1H, H-C(7)), 2.28 – 2.13 (*m*, 1H, CH^aH^bCH₂Ph), 1.74 – 1.58 (*m*, 1H, CH^aH^bCH₂Ph).

¹³C NMR (CDCl₃, 100 MHz): δ 178.43 (C(5)), 141.72 (*C*-arom), 128.67, 128.56, 126.11 (*C*H-arom), 47.42 (C(6)), 45.04 (C(3)), 33.69 (CH₂CH₂Ph), 33.54 (CH₂CH₂Ph), 31.17 (C(2)), 30.36 (C(7)).

ESI⁺-HRMS: calculated for C₁₃H₁₈NOS ([M+H]⁺) 236.1109, found 236.1104.

<u> 3c,d:</u>

TLC (PE/EtOAc 2:3) $R_f = 0.16$.

¹H NMR (CDCl₃, 400 MHz): δ 7.31 – 7.14 (*m*, 5H, H-arom), 6.62 (*s*, 1H, NH), 3.58 (*td*, *J* = 6.0, 4.3 Hz, 2H, H-C(3)), 3.01 – 2.90 (*m*, 2H, 2xH-C(6)), 2.86 – 2.73 (*m*, 4H, C(7), CH₂CH₂Ph, H-C(2)), 2.71 – 2.62 (*m*, 1H, H-C(2)), 2.03 – 1.84 (*m*, 2H, CH^aH^bCH₂Ph).

¹³C NMR (CDCl₃, 100 MHz): δ 176.24 (C(5)), 141.15 (*C*-arom), 128.65, 128.59, 126.19 (*C*H-arom), 46.56, C(6)), 45.90 (C(3)), 37.01 (C(7)), 36.58 (CH₂CH₂Ph), 33.37 (CH₂CH₂Ph), 29.98 (C(2)).

ESI⁺-HRMS: calculated for $C_{13}H_{18}NOS$ ([M+H]⁺) 236.1109, found 236.1102.



A solution of 5-hexynoic acid 4 (4.00 g, 35.7 mmol, 1 equiv.) in dry CH_2Cl_2 (130 ml) was cooled to 0°C. Next, DIPEA (14 mL, 80.4 mmol, 2.25 equiv.) and pentafluorophenyl trifluoroacetate (12 ml, 69.8 mmol, 1.95 equiv.) were added dropwise, and the resulting solution was allowed to warm up to r.t. After stirring for 1.5 h at r.t. the reaction mixture was concentrated on a rotovap, and the crude product was purified by CC (PE/Et₂O 98:2) to give product 5 (9.44 g, 95%) as a colourless liquid.

TLC (PE/Et₂O 4:1) *R*_f= 0.80.

¹H NMR (CDCl₃, 400 MHz): δ 2.84 (t, *J* = 7.4 Hz, 2H, H-C(2)), 2.36 (td, *J* = 6.9, 2.7 Hz, 2H, H-C(4)), 2.05 - 1.94 (m, 3H, H-C(6), H-C(3)).

 13 C NMR (CDCl₃, 100 MHz): δ 169.15 (C(1)), 142.67 – 142.12, 141.01 – 140.48, 140.13 – 139.69, 139.45 – 138.92, 138.53 – 137.94, 137.11 – 136.27, 125.44 – 124.70 (m, (C-arom)), 82.57 (C(5)), 69.90 (C(6)), 32.02 (C(2)), 23.46 (C(3)), 17.75 (C(4)).

¹⁹F NMR (CDCl₃, 376 MHz): δ -153.72 - -153.88 (m), -159.01 (t, *J* = 21.6 Hz), -163.24 - -163.44 (m).

ESI⁺-HRMS: calculated for $C_{12}H_8F_5O_2$ ([M+H]⁺) 279.0444, found 279.0435.

Perfluorophenyl 4-(3'-oxocycloprop-1'-en-1'-yl)butanoate 6



To an oven-dried, 10 ml pressure tube equipped with a magnetic stir bar anhydrous NaI (330 mg, 2.2 mmol) was added in a glove box. The flask was then filled with argon and transferred to a fume hood. Next, anhydrous THF (3.0 mL), perfluorophenyl hex-5-ynoate **5** (278 mg, 1.0 mmol), and trifluoromethyltrimethylsilane (300 μ L, 2.0 mmol) were added, and the reaction was stirred vigorously at room temperature for 24 h. After that, a second portion of trifluoromethyltrimethylsilane (150 μ L, 1.0 mmol) was added, and the reaction mixture was stirred for another 24 h. The reaction mixture was then diluted with H₂O (20 mL) and extracted with Et₂O (3 x 20 mL). The combined organic layers with crude difluorocyclopropene were dried over MgSO₄, filtered, concentrated, adsorbed on silica, and left overnight at room temperature to generate the corresponding cyclopropenone. The product was then purified by CC (PE/EtOAc 95:5 to 1:1 to 0:1) to afford the perfluorophenyl 4-(3'-oxocycloprop-1'-en-1'-yl)butanoate **6** (131 mg, 43%) as a white solid.

TLC of the difluorocyclopropene intermediate (PE/Et₂O 4:1) R = 0.45.

TLC of **6** (EtOAc) $R_f = 0.50$.

¹H NMR (CDCl₃, 500 MHz): δ 8.53 (s, 1H, H-C(2')), 2.89 – 2.82 (m, 4H, H-C(2), H-C(4)), 2.22 (p, *J* = 7.3 Hz, 2H, H-C(3)).

¹³C NMR (CDCl₃, 125 MHz): δ 168.85 (C(1)), 168.59 (C(1')), 157.21 (C(3')), 149.25 (C(2')), 142.30 – 141.82, 140.84 – 140.41, 140.16 – 139.86, 139.09 – 138.37, 137.07 – 136.71, 125.01 – 124.60 (m, (C-arom)), 32.25 (C(2)), 26.80 (C(4)), 20.97 (C(3)).

¹⁹F NMR (CDCl₃, 376 MHz): δ -152.03 - -153.11 (m), -157.47 (t, *J* = 21.7 Hz), -160.76 - -162.53 (m).

ESI⁺-HRMS: calculated for $C_{13}H_7F_5O_3Na$ ([M+Na]⁺) 329.0208, found 329.0200.

4-(3'-oxocycloprop-1'-en-1'-yl)-N-phenylbutanamide CPO-BN



Perfluorophenyl 4-(3'-oxocycloprop-1'-en-1'-yl)butanoate **6** (800 mg, 2.6 mmol) was dissolved in CH₂Cl₂ (32 mL), and portions of benzylamine (600 μ L in total, 5.5 mmol, 2.1 equiv.) were added every 5 – 10 min until there was no starting material left according to TLC. After that, the reaction mixture was concentrated* and purified by CC (EtOAc 100%; CH₂Cl₂/MeOH 95:5)** to give **CPO-BN** (542.5 mg, 91%) as a brown oil.

* Avoid complete evaporation of solvent as this can result in product decomposition if large excess of benzylamine is used.

** Column first washed with EtOAc to remove the by-product (pentafluorophenol) and the product then washed out with $CH_2Cl_2/MeOH$ 95:5.

TLC (EtOAc) $R_f = 0.17$.

¹H NMR (CDCl₃, 400 MHz): δ 8.40 (s, 1H, H-C(2')), 7.36 – 7.23 (m, 5H, H-arom), 6.42 (br, 1H, NH), 4.43 (d, J = 5.8 Hz, 2H, CH₂Ph), 2.74 (t, J = 7.0 Hz, 2H, H-C(4)), 2.40 (t, J = 7.0 Hz, 2H, H-C(2)), 2.08 (p, J = 7.0 Hz, 2H, H-C(3)).

¹³C NMR (CDCl₃, 125 MHz): δ 171.48 (C(1)), 169.60 (C(1')), 158.22 (C(3')), 148.14 (C(2')), 138.34 (C-arom), 128.86, 128.01, 127.68 (CH-arom), 43.82 (CH₂Ph), 34.90 (C(2)), 26.82 (C(4)), 21.74 (C(3)).

ESI⁺-HRMS: calculated for C₁₄H₁₆NO₂ ([M+H]⁺) 230.1176, found 230.1169.

N-(2-(2-(2-(2-hydroxyethoxy)ethoxy)ethoxy)ethyl)-4-(3'-oxocycloprop-1'-en-1'yl)butanamide CPO-PEG



Perfluorophenyl 4-(3'-oxocycloprop-1'-en-1'-yl)butanoate 6 (10 mg, 0.03 mmol) was dissolved in CH_2Cl_2 (1 mL), and portions of HO-PEG4-NH₂ (10 mg in total, 0.05 mmol, 1.6 equiv.) were added every 5 – 10 min until there was no starting material left according to TLC. After that, the reaction mixture was concentrated and purified by CC (CH₂Cl₂/MeOH 93:7 to 4:1) to give **CPO-PEG** (9.2 mg, 89%) as a colourless oil.

TLC (CH₂Cl₂/MeOH 93:7) $R_f = 0.27$.

¹H NMR (CDCl₃, 400 MHz): δ 8.46 (s, 1H, H-C(2')), 7.40 (br, 1H, NH), 3.75 – 3.70 (m, 4H, 2 x CH₂-PEG), 3.68 – 3.59 (m, 8H, 4 x CH₂-PEG), 3.54 – 3.51 (m, 2H, CH₂-PEG), 3.44 (q, *J* = 5.0 Hz, 2H, CH₂NH(PEG)), 2.75 (t, *J* = 7.1 Hz, 2H, H-C(4)), 2.34 (t, *J* = 7.2 Hz, 2H, H-C(2)), 2.06 (p, *J* = 7.1 Hz, 2H, H-C(3)).

¹³C NMR (CDCl₃, 125 MHz): δ 171.92 (C(1)), 169.77 (C(1')), 158.04 (C(3')), 148.57 (C(2')), 72.68, 70.82, 70.57, 70.32, 70.12, 70.07, 61.65 (7 x CH₂-PEG), 39.30 (CH₂NH(PEG)), 34.81 (C(2)), 27.11 (C(4)), 21.74 (C(3)).

ESI⁺-HRMS: calculated for $C_{15}H_{26}O_6N_1$ ([M+H]⁺) 316.1755, found 316.1744.
4-(3'-oxocycloprop-1'-en-1'-yl)-N-(3'',6'',9'',12''-tetraoxapentadec-14''-yn-1''-yl)butanamide CPO-PEG-Alkyne



Perfluorophenyl 4-(3'-oxocycloprop-1'-en-1'-yl)butanoate **6** (10.0 mg, 32.7 μ mol) was dissolved in CH₂Cl₂ (1 mL), and portions of alkyne-PEG4-NH₂ (15.1 mg in total, 65.3 μ mol, 2 equiv.) were added every 5 – 10 min until there was no starting material left according to TLC. After that, the reaction mixture was concentrated and purified by CC (CH₂Cl₂/MeOH 93:7 to 4:1) to give **CPO-PEG-Alkyne** (8.0 mg, 70%) as a colourless oil.

TLC (CH₂Cl₂/MeOH 9:1) $R_f = 0.20$.

¹H NMR (CDCl₃, 400 MHz): δ 8.47 (s, 1H, H-C(2')), 6.68 (br, 1H, NH), 4.19 (d, J = 2.4 Hz, 2H, H-C(13'')), 3.73 – 3.59 (m, 12H, 6 x CH₂-PEG), 3.57 – 3.52 (m, 2H, CH₂-PEG), 3.44 (q, J = 5.2 Hz, 2H, CH₂NH(PEG)), 2.76 (t, J = 7.1 Hz, 2H, H-C(4)), 2.44 (t, J = 2.3 Hz, 1H, H-C(15'')), 2.35 (t, J = 7.1 Hz, 2H, H-C(3)).

¹³C NMR (CDCl₃, 100 MHz): δ 171.82 (C(1)), 169.66 (C(1')), 158.03 (C(3')), 148.48 (C(2')), 79.61 (C(14'')), 74.84 (C(15'')), 70.65, 70.61, 70.44, 70.29, 69.92, 69.22 (6 x CH₂-PEG), 58.53 (C(13'')), 39.36 (CH₂NH(PEG)), 34.89 (C(2)), 26.96 (C(4)), 21.71 (C(3)).

ESI⁺-HRMS: calculated for C₁₈H₂₈O₆N₁ ([M+H]⁺) 354.1911, found 354.1902.

Sodium 5-((2'-(4''-(3'''-oxocycloprop-1'''-en-1'''-yl)butanamido)ethyl)amino) naphthalene-1-sulfonate CPO-EDANS



Perfluorophenyl 4-(3'-oxocycloprop-1'-en-1'-yl)butanoate 6 (10.0 mg, 32.7 μ mol) was dissolved in DMF (400 μ L), and N-(aminoethyl)-5-naphthylamine-1-sulfonic acid in DMF (18.8 mg in 600 μ L DMF, 65.2 μ mol, 2 equiv.) was added and the reaction mixture was stirred for 30 min, concentrated, and purified by CC (EtOAc/MeOH 1:0 to 1:1) to give **CPO-EDANS** (12.3 mg, 92%).

TLC (EtOAc/MeOH 7:3) $R_{f} = 0.23$.

¹H NMR (DMF- d_7 , 500 MHz): δ 8.97 (s, 1H, H-C(2^{''})), 8.43 (d, J = 8.7 Hz, 1H, H-arom), 8.27 (s, 1H, NH), 8.15 (d, J = 8.5 Hz, 1H, H-arom), 8.12 (d, J = 7.0 Hz, 1H, H-arom), 7.35 – 7.24 (m, 2H, H-arom), 6.59 (d, J = 7.5 Hz, 1H, H-arom), 6.13 (s, 1H, NH), 3.59 – 3.52 (m, 2H, H-C(2['])), 3.41 – 3.36 (m, 2H, H-C(1['])), 2.78 – 2.71 (m, 2H, H-C(4^{''})), 2.37 (t, J = 7.3 Hz, 2H, H-C(2^{''})), 2.04 – 1.95 (m, 2H, H-C(3^{''})).

¹³C NMR (DMF, 126 MHz): δ 173.58 (C(1'')), 170.79 (C(1''')), 158.10 (C(3''')), 151.04 (C(2''')), 146.25 (C-arom), 145.12 (C-arom), 131.97 (C-arom), 127.06 (CH-arom), 125.62 (CH-arom), 125.14 (C-arom), 123.46 (CH-arom), 123.45 (CH-arom), 117.61 (CH-arom), 103.75 (CH-arom), 45.16 (C(1')), 39.22 (C(2')), 35.39 (C(4'')), 27.59 (C(2'')), 22.93 (C(3'')).

ES⁻-HRMS: calculated for C₁₉H₁₉N₂O₅S ([M]⁻) 387.1015, found 387.1019.

4. X-ray structure analysis of 2a

Methods

Single-crystal X-ray diffraction data were collected using a D8-QUEST PHOTON-100 diffractometer equipped with an Incoatec IµS Cu microsource (CuK α radiation, λ = 1.5418 Å). The temperature was held at 180(2) K using an Oxford Cryosystems N₂ cryostat. Data integration and reduction were undertaken with SAINT in the APEX3 software suite. Multi-scan corrections were applied using SADABS. Structures were solved using SHELXT and refined using SHELXL.¹⁻²

X-Ray Crystallography

The crystal structure of compound 2a is refined as a 2-component non-merohedral twin (HKLF-5 refinement in SHELXL). The benzene ring C11–C16 is modelled as disordered over two components. The intermolecular contacts indicate that the two components must alternate along the short *b* axis, and the site occupancies of both components are therefore constrained to 0.5. There was no clear evidence of doubling of this cell axis. The displacement ellipsoid of atom C8 indicates clearly that this site has partial site occupancy. The atoms makes unreasonably short contacts to its symmetry equivalent generated by the action of 2-fold rotation axis, so its site occupancy factor was also constrained to 0.5. The disorder represents a mixture of EtO and MeO substituents.



Figure S1. Molecular structure of **2a** showing displacement ellipsoids at 50% probability. H atoms are omitted. One disorder component of the benzene ring C11–C16 is shown as semi-transparent. Atom C8 has 50% site occupancy.

 Table S1. Crystal data, data collection, and refinement parameters for 2a.

Compound	2a
CCDC Deposition No.	2010531
Cambridge Data No.	GB_B1_0001
Formula	C ₁₆ H ₂₁ NO ₃ S, C ₁₅ H ₁₉ NO ₃ S
M.w.	600.77
Crystal system	monoclinic
Space group (IT no.)	C2 (5)
a [Å]	22.5982 (14)
b [Å]	5.2231 (3)
<i>c</i> [Å]	13.4924 (8)
α [°]	90
β[°]	106.471 (4)
γ [°]	90
Z	2
V [Å ³]	1527.19(16)
D _x [g cm ⁻³]	1.306
Crystal size [mm]	0.30 x 0.05 x 0.01
Crystal colour, shape	colourless lath
μ [mm ⁻¹]	1.954
T_{\min}, T_{\max}	0.592, 0.981
Measured reflections	17502
Independent reflections (R_{int}^{a})	2572 (0.065)
Observed reflections [I> 2σ (I)]	2294
No. of parameters / restraints	226 / 147
R1 (observed data) ^b	0.065
$wR2(F^2)$ for all data	0.165
GOF°	1.05
Residual electron density [e·Å ⁻³]	-0.26, 0.30
Flack parameter	0.04 (5)

 ${}^{a}R_{int} = \Sigma |F_{o}^{2} - F_{o,mean}^{2}| / \Sigma F_{o}^{2}$ ${}^{b}R(F) = \Sigma ||F_{o}| - |F_{c}|| / \Sigma |F_{o}|; wR(F^{2}) = [\Sigma(w(F_{o}^{2} - F_{c}^{2})^{2}) / (\Sigma w(F_{o}^{2})^{2})]^{\frac{1}{2}}$ ${}^{c}GOF = [\Sigma(w(F_{o}^{2} - F_{c}^{2})^{2}) / (N_{diffrs} - N_{params})]^{\frac{1}{2}}$

5. Quantum mechanical calculations

Full geometry optimizations were carried out with Gaussian 16³ using the M06-2X⁴ hybrid functional and 6-31+G(d,p) basis set in combination with ultrafine integration grids. Bulk solvent effects in water were considered implicitly through the IEF-PCM polarizable continuum model.⁵ The possibility of different conformations was taken into account. Frequency analyses were carried out at the same level used in the geometry optimizations, and the nature of the stationary points was determined in each case according to the appropriate number of negative eigenvalues of the Hessian matrix. The quasiharmonic approximation reported by Truhlar et al. was used to replace the harmonic oscillator approximation for the calculation of the vibrational contribution to enthalpy and entropy.⁶ Scaled frequencies were not considered. Mass-weighted intrinsic reaction coordinate (IRC) calculations were carried out by using the Gonzalez and Schlegel scheme⁷⁻⁸ in order to ensure that the TSs indeed connected the appropriate reactants and products. Gibbs free energies (ΔG) were used for the discussion on the relative stabilities of the considered structures. Free energies calculated using the gas phase standard state concentration (1 atm = 1/24.5 M) were converted to reproduce the standard state concentration in solution (1 M) by adding or subtracting 1.89 kcal mol⁻¹ for bimolecular additions and decompositions, respectively. The lowest energy conformer for each calculated stationary point was considered in the discussion; all the computed structures can be obtained from authors upon request. Cartesian coordinates, electronic energies, entropies, enthalpies, Gibbs free energies, and lowest frequencies of the calculated structures are available in the Supporting Information.



Figure S2. Lowest-energy structures calculated with PCM(H₂O)/M06-2X/6-31+G(d,p)



Figure S3. a) Minimum energy pathways for the reaction between 2-methylcyclopropenone (1') and methanethiolate (**MT**) calculated with PCM(H₂O)/M06-2X/6-31+G(d,p). Note that each calculated structure represents a racemic compound constituted by two enantiomers with identical energies. b) Lowest energy transition structures calculated at the same level for the reactions between **1'** and methanethiolate (**TS1'** A and **TS1'** B) or methylamine (**TS1_N'** A and **TS1_N'**). *****: protonation and deprotonation equilibria were calculated using tri-hydrated bicarbonate (HCO₃²⁻ · 3 H₂O) and carbonate (CO₃²⁻ · 3 H₂O) anions as an acid and a base, respectively. Given the intrinsic inaccuracy of such estimations, relative energies of charges/neutral species must be considered with caution. Relative free energies (Δ G) are given in kcal mol⁻¹. Numbers in brackets represent the relative energies of transition structures.



Figure S4. Comparison between the ring-expansion reaction pathways (**TS3**) to give the experimentally observed 1,4-thiazepan-5-ones (**TZ**), and the elimination reaction pathways (**TS4**) to afford non-detected acrylamides, calculated with PCM(H₂O)/M06-2X/6-31+G(d,p). Both reaction channels are accessible from the same deprotonated hemiaminal intermediates (HA-). *: protonation and deprotonation equilibria were calculated using tri-hydrated bicarbonate (HCO₃²⁻ · 3 H₂O) and carbonate (CO₃²⁻ · 3 H₂O) anions as an acid and a base, respectively. Given the intrinsic inaccuracy of such estimations, relative energies of charges/neutral species must be considered with caution. Relative free energies (Δ G) are given in kcal mol⁻¹. Numbers in brackets represent the relative energies of transition structures.

Stanotuno	Eelec	$E_{elec} + ZPE$	Н	S	G	Lowest	# of imag.
Structure	(Hartree) ^a	(Hartree) ^a	(Hartree) ^a	(cal mol ⁻¹ K ⁻¹) ^b	(Hartree) ^{a,b}	freq. (cm ⁻¹)	freq.
1'	-229.894568	-229.827621	-229.822095	71.2	-229.853405	86.0	0
CA	-532.803161	-532.718809	-532.713469	70.3	-532.744529	161.5	0
CO3 ²⁻	-493.241249	-493.153831	-493.142707	106.1	-493.186493	21.0	0
HCO3 ⁻	-493.724975	-493.624401	-493.612104	111.2	-493.658113	26.9	0
TS1 A	-762.697276	-762.544737	-762.534279	101.4	-762.577754	-203.9	1
TS1 B	-762.696075	-762.543708	-762.533464	99.1	-762.576138	-198.1	1
CP-A	-762.704939	-762.551431	-762.541039	99.6	-762.584242	54.1	0
СР-В	-762.703922	-762.551200	-762.540847	99.1	-762.583429	27.3	0
CP Aanti	-763.193541	-763.026745	-763.016203	101.2	-763.059447	36.9	0
CP Asyn	-763.193715	-763.026639	-763.016390	97.8	-763.059106	63.3	0
СР В	-763.195992	-763.029221	-763.018959	97.2	-763.061417	62.5	0
TS2 Aanti	-763.185246	-763.017571	-763.008353	92.0	-763.048752	-163.1	1
TS2 A _{syn}	-763.182180	-763.014015	-763.004984	90.9	-763.045077	-227.8	1
ТS2 В	-763.186291	-763.018828	-763.009757	90.6	-763.049682	-160.4	1
HA _{zw} A _{anti}	-763.197557	-763.026260	-763.017442	90.5	-763.056830	51.7	0
HA _{zw} A _{syn}	-763.195426	-763.024040	-763.015438	88.6	-763.054511	91.5	0
HA _{zw} B	-763.199218	-763.028153	-763.019456	88.8	-763.058533	85.7	0
HA Aanti	-763.225435	-763.055351	-763.046300	90.8	-763.086129	71.9	0
HA A _{syn}	-763.225867	-763.055648	-763.046739	89.6	-763.086305	117.0	0
HAB	-763.226112	-763.056312	-763.047413	89.4	-763.086804	90.7	0
HA ⁻ Aanti	-762.727403	-762.571089	-762.562409	89.8	-762.601515	53.8	0
$HA^{-}A_{syn}$	-762.727193	-762.570507	-762.562032	88.2	-762.600794	79.6	0
HA ⁻ B	-762.727376	-762.571277	-762.562659	89.4	-762.601461	46.3	0
TS3 Aanti	-762.716572	-762.561878	-762.553359	87.9	-762.592162	-444.8	1
TS3 A _{syn}	-762.713517	-762.558268	-762.549783	88.2	-762.588623	-478.1	1
TS3 B	-762.712731	-762.558312	-762.549787	88.4	-762.588715	-383.2	1
TZ- A	-762.735858	-762.579768	-762.570995	89.3	-762.610385	95.9	0
TZ-B	-762.732639	-762.576799	-762.568003	89.3	-762.607399	95.9	0
TZ A	-763.279806	-763.108658	-763.099819	90.1	-763.139395	79.6	0
TZ B	-763.281038	-763.109631	-763.100892	89.3	-763.140201	85.7	0
TS4 Aanti	-762.703919	-762.549917	-762.541109	89.9	-762.580819	-333.8	1
TS4 A _{syn}	-762.701186	-762.546553	-762.538154	87.2	-762.576670	-362.4	1
TS4 B	-762.709512	-762.555732	-762.547245	87.1	-762.585720	-413.5	1

Table S2. Energies, entropies, and lowest frequencies of the lowest energy calculated structures.

(E)-Crotonamide	-762.792298	-762.636320	-762.626139	100.6	-762.669065	37.5	0
(Z)-Crotonamide	-762.787855	-762.631682	-762.621500	101.7	-762.664396	24.9	0
Methacrylamide	-762.790467	-762.634338	-762.624416	99.5	-762.666527	25.5	0
MT	-438.170103	-438.133138	-438.129472	55.2	-438.153846	715.7	0
TS1' A	-668.063096	-667.957948	-667.949298	91.7	-667.988765	-194.5	1
Т\$1' В	-668.062025	-667.956994	-667.948546	89.3	-667.987212	-198.0	1
СР'- А	-668.071038	-667.965047	-667.956305	92.0	-667.995658	26.4	0
СР'- В	-668.070953	-667.965492	-667.956962	88.6	-667.995543	55.4	0
CP' Aanti	-668.558509	-668.439023	-668.430411	90.2	-668.469169	31.1	0
CP' Asyn	-668.560304	-668.440529	-668.431914	90.1	-668.470901	43.9	0
CP' B	-668.563369	-668.443998	-668.435535	87.8	-668.474020	71.5	0
MA	-95.815844	-95.751246	-95.747258	56.6	-95.772265	303.9	0
TS1' _N A	-325.702132	-325.567645	-325.559027	90.6	-325.597637	-295.5	1
TS1' _N B	-325.702159	-325.567715	-325.559485	86.3	-325.597134	-296.1	1

^aEnergy values calculated with PCM_{H2O}/M06-2X/6-31+G(d,p). 1 Hartree = 627.51 kcal mol-1. ^bThermal corrections at 277.15 K.

Cartesian coordinates of the lowest-energy calculated structures.

1'				Н		1.385697	-1.629816	1.519438
С	0.553927	0.106997	-0.000058	Н		0.258920	-0.309928	1.202523
С	-0.200592	1.228355	-0.000011	С		2.330469	0.210867	0.907511
С	-0.859919	-0.019001	0.000013	Н		2,436076	0.550543	1,945965
0	-1 880816	-0 693094	0 000006	Н		3 267196	-0 291373	0 638006
C	1 917424	-0 464407	0 000003	N		2 185811	1 388943	0 051671
ч	2 044666	-1 099026	0.880794	н		2 085592	1 054014	-0 906190
11	2.044000	1 000262	0.000794	11		1 202227	1 0/1010	0.251457
п	2.044734	-1.099202	-0.0000391	п		1.295257	1.041910	0.231437
H	2.0/4/30	0.319984	-0.000012	m 01				
н	-0.182645	2.311389	0.000077	TSI	В	1 222020	0 626600	1 21 (102
CA				C		-1.323920 1 504052	0.030000	1.310102
CA	1 404057	0 217705	0 0 0 0 0 0 0	C		-1.J940J3	-0.097020	0.100779
S	1.494857	-0.317705	-0.062999	C		-1.1/323/	1.269832	0.098709
C	0.145407	0.882075	0.293348	0		-0.902850	2.262172	-0.593/32
H	0.428395	1.8/8346	-0.0663/3	S		0.363119	-1.522141	-0.464/62
Н	-0.000133	0.962880	1.3/8/88	С		1.504563	-0.824931	0./66139
С	-1.190719	0.499276	-0.34/4/0	Н		1.965525	-1.629676	1.349371
Н	-1.938612	1.270946	-0.123543	Н		0.908607	-0.218117	1.471308
Н	-1.068182	0.473856	-1.437271	С		2.600847	0.066677	0.173336
Н	-0.986120	-1.478995	-0.025911	H		3.251459	0.421402	0.982911
N	-1.735146	-0.793686	0.076122	Н		3.225155	-0.528547	-0.503375
Н	-1.935172	-0.756055	1.074171	N		2.112994	1.233459	-0.563933
				Н		1.429512	0.901768	-1.244654
CO32-				Н		1.570570	1.824247	0.064466
С	0.053655	-0.081861	-0.000940	С		-2.627599	-1.044864	-0.348105
0	0.281550	-1.340444	-0.001466	Н		-2.426303	-1.319342	-1.384762
0	1.023036	0.778619	0.001399	Н		-2.664735	-1.950719	0.259783
0	-1.175651	0.365905	-0.002758	Н		-3.603705	-0.547768	-0.297887
0	3.104947	-0.768906	0.000440	Н		-1.348915	0.631215	2.398177
0	-0.807729	2.967098	0.000635					
0	-2.492021	-1.923540	0.002049	CP-	A			
Н	2.553996	-1.563766	-0.000579	С		2.094457	0.085901	0.140741
Н	2.367579	-0.078622	0.000943	С		0.717036	-0.224533	0.594566
Н	0.151343	2.839708	0.002847	С		1.203488	1.054296	-0.025853
Н	-1.080834	2.003545	-0.001514	0		0.780126	2.189528	-0.406028
Н	-2.162001	-0.982357	0.000098	С		3.446010	-0.527957	0.035425
Н	-1.625069	-2.357191	0.001457	Н		3.830172	-0.832616	1.016604
				Н		3.432224	-1.423499	-0.598021
HCO ₃ ⁻				Н		4.160587	0.175977	-0.400686
С	-0.011258	0.045057	-0.018193	Н		0.436387	-0.331256	1.653320
0	-1.081607	0.706512	-0.027765	S		-0.337682	-1.408468	-0.388397
0	1.159465	0.496492	-0.008912	С		-1.997525	-1.084496	0.294992
0	-0.098963	-1.321056	-0.018236	Н		-2.600535	-1.964290	0.047907
0	0.323411	3.177070	0.022322	Н		-1.911760	-1.045681	1.388695
0	2.871284	-1.656853	0.022788	С		-2.697358	0.177180	-0.219862
0	-2.911116	-1.257834	-0.070052	Н		-2.702624	0.160449	-1.315523
Н	-0.515467	2.689693	0.005524	Н		-3.745469	0.128601	0.103148
Н	0.942491	2.425356	0.016546	Н		-2.112128	1.482850	1.233779
Н	2.316596	-0.846175	0.011112	Н		-1.187037	1.588830	-0.107622
Н	2.220082	-2.367566	0.014222	Ν		-2.147795	1.456793	0.217276
Н	-1.042123	-1.576954	-0.015973					
Н	-2.513395	-0.361251	-0.007902	CP-	A			
Н	-3.440421	-1.388092	0.724467	С		-1.157894	0.581619	1.342605
				С		-1.276272	-0.235736	0.120587
TS1 A				C		-1.059817	1.237845	0.179350
C	-1.931297	-0.017403	0.123667	0		-0.903291	2.267143	-0.533086
C	-1.269200	-0.199365	-1.085318	S		0.136781	-1.416937	-0.421372
C	-1.248875	1.057366	-0.401308	С С		1.512379	-0.940803	0.665191
0	-0.858604	2.239362	-0.368652	H		1.067883	-0.420293	1.524144
C	-2.777412	-0.686973	1.142535	Н		1.982547	-1.859073	1.029742
H	-2.223469	-1.523785	1.582878	C		2.556494	-0.043806	-0.005735
Н	-3.059663	0.000591	1.941442	н		3.375146	0.123762	0.704830
Н	-3.684391	-1.097073	0.688434	Н		2.985033	-0.566962	-0.867851
Н	-1.423639	-0.670130	-2.046328	Н		1.664665	1.757390	0.334271
S	0.846389	-1.434770	-0.835568	Н		1.276507	1.116804	-1.086224
C	1.178044	-0.798847	0.836074	N		2.065849	1.256588	-0.456523

С	-2.556165	-0.895954	-0.366920	Н	2.322604	-0.493594	1.866892
Η	-2.499856	-1.154071	-1.431471	Н	0.700638	-1.175401	2.032893
Н	-2.768087	-1.815677	0.191002	Н	3.226393	0.362972	-0.329281
Н	-3.394897	-0.205911	-0.228509				
н	-1.224406	0.542783	2.421890	TS2	Annti		
	1,00,100	0.012,000	2.122090		-1 345989	-0 249863	0 668754
CР	Δ			C	-0 574800	-0 725922	-0 597370
C	1 568666	0 710618	0 456861	C	-1 052555	0 669/35	-0 159953
c	1.500000	0.710010	0.450001	0	1 601104	1 540005	1 110001
C	0.070470	-0.204031	-0.433088	0	-1.301104	1.549605	-1.110001
C	2.094363	-0.482415	-0.199422	C	-2.762004	-0.724621	0.911922
0	3.021152	-1.217810	-0.414728	Н	-2.762272	-1.730567	1.341523
С	1.919678	2.104470	-0.022772	H	-3.282709	-0.055605	1.602415
Η	1.095141	2.791007	0.185455	Н	-3.327735	-0.750622	-0.024144
Η	2.815195	2.467269	0.486893	Н	-1.179591	-1.304303	-1.297570
Н	2.112906	2.110378	-1.098574	S	1.086602	-1.417946	-0.603613
Н	0.419773	0.216476	-1.426606	Н	0.527468	1.944586	1.290288
S	-0.552549	-1.275481	0.297800	Ν	0.786272	1,661294	0.347848
Ċ	-1.655636	-0.011944	1.015174	Н	-0.726153	-0.149269	1.555483
н	-2 406345	-0 578919	1 575183	C	1 961824	-0 461033	0 670752
ц Ц	-1 080363	0 585/15	1 730/85	с ц	1 478148	-0 604515	1 6/2097
п	-1.000303	0.000410	1.730403	п	2 0/7527	-0.004515	1.042097
C II	-2.323030	0.092117	-0.022012	п	2.94/33/	-0.930633	0.725445
н	-1.55/604	1.459340	-0.559/58	C	2.103839	1.015/11	0.336189
Н	-2.931770	1.626132	0.519976	Н	2.796389	1.482084	1.048782
Η	-3.928472	-0.272004	-0.528191	H	2.529929	1.119262	-0.664676
Η	-2.639102	-0.478039	-1.503465	Н	0.776396	2.498801	-0.227493
Ν	-3.176446	0.224332	-1.000257				
Η	1.377103	0.603629	1.524841	TS2	A _{syn}		
				С	-1.828358	-0.309849	-0.176098
CP	Asvn			С	-0.501176	-0.488807	-1.013626
С	-1.330990	0.938479	0.681330	С	-0.968274	0.835075	-0.491923
С	-0.868533	-0.580501	0.636288	0	-1.028936	1,992081	-0.876791
C	-2.191211	-0.130253	0.186720	C	-2.015601	-0.958700	1.173958
0	-3 186353	-0 431989	-0 420900	н	-2 775547	-0 415393	1 742411
Ĉ	-0 813169	1 978570	-0 288083	и Ц	-2 352068	-1 992403	1 051210
ц Ц	-1 527295	2 801335	-0 369745	и Ц	_1 088937	-0 966360	1 7/5923
п	-1.327293	2.001333	-0.309743	п	-1.000937	-0.900300	2 004000
п	0.143034	2.303097	1 204502	п	-0.043009	-0.014907	-2.004909
н	-0.672769	1.550390	-1.284502	5	0.824/26	-1.4//885	-0.329329
Н	-0.81/066	-1.035996	1.62458/	C	2.163340	-0.262368	-0.0358/5
S	0.278637	-1.256792	-0.556072	Н	2.747089	-0.702969	0.777408
С	1.833252	-1.088651	0.387548	Н	2.807331	-0.192390	-0.914497
Η	1.715704	-1.616710	1.338841	C	1.671602	1.133359	0.346344
Η	2.580385	-1.623559	-0.207618	Н	1.409104	1.697792	-0.552788
С	2.269430	0.359025	0.618774	Н	2.492872	1.663105	0.842983
Н	3.169066	0.343125	1.243180	Н	0.180720	1.982947	1.485759
Н	1.497515	0.880277	1.196647	Н	0.612234	0.476633	1.991348
Н	1.766039	1.097386	-1.218206	Ν	0.462676	1.058510	1.169178
н	3.345249	0.719517	-1.079811	н	-2.721160	-0.328293	-0.802097
N	2 560386	1 133601	-0 582525				
н	-1 482605	1 269907	1 706970	TS2	B		
	1.102000	1.20000	1.100010	 C	-0 994529	0 422558	1 369972
CP	В			C	-0 949522	-0.449297	0.080832
C	2 223856	0 300611	-0 7/9526	C	-0 981648	1 028956	0.012714
Ĉ	1 120091	-0 339601	0.749520	0	-1 501/13	1 0200/1	-0 623639
C	1 100051	1 12007	0.000400	0	-1.301413	1.929941	-0.023030
C	1.189251	1.126907	-0.013087	5	0.432702	-1.558147	-0.284102
0	0.68/688	2.184182	0.2/06/3	Н	1.3498//	2.08/365	0./32981
S	-0.141875	-1.115245	-0.942738	N	1.196912	1.546298	-0.114861
С	-1.553639	-1.145656	0.212669	H	-0.098234	0.464615	1.978810
Н	-2.369960	-1.580401	-0.373279	С	1.897501	-0.736762	0.407795
Н	-1.332115	-1.835367	1.032512	Н	1.762972	-0.575810	1.481810
С	-1.962663	0.225726	0.760154	Н	2.697842	-1.472131	0.287811
Н	-1.140618	0.650764	1.346531	С	2.259328	0.554467	-0.309655
Н	-2.793756	0.060424	1.454776	Н	3.230059	0.912732	0.056006
Н	-3.112777	0.844691	-0.814901	Н	2.355096	0.354842	-1.379551
Н	-1.593729	1.444960	-0.837523	H	1.152999	2.206271	-0.885866
N	-2.372845	1.217825	-0.224978	C	-2.246327	-1.128701	-0.325468
н	2.171803	0.364079	-1.835149	н	-2.234874	-1.380187	-1.390131
Ċ	1 544004	-1.059322	1.346871	н	-2 390768	-2.046801	0.252617
н	1,934599	-2.049081	1.094883	н	-3.095583	-0.464970	-0.142576
				11	2.02000	· · · · · / / / / / / / / / / / / / / /	/ V

Η	-1.928530	0.433481	1.929723
HA-w	Aanti		
C	-1.498594	-0.449710	0.513140
С	-0.413852	-0.720694	-0.540114
С	-0.731071	0.700584	-0.069552
0	-1.182467	1.661262	-0.807481
С	-2.948445	-0.606354	0.128321
Н	-3.242705	-1.659986	0.123636
Н	-3.593827	-0.072492	0.832381
Н	-3.121299	-0.188677	-0.866516
Н	-0.756135	-0.959073	-1.543876
S	1.101054	-1.508739	-0.008713
С	2.294617	-0.130950	-0.218145
Н	3.089268	-0.333519	0.502642
Н	2.727838	-0.159946	-1.218555
С	1.689196	1.260128	0.018329
Н	1.379275	1.727070	-0.917090
Н	2.404804	1.909914	0.522247
Н	0.167038	2.125776	1.139906
Н	0.581549	0.612253	1.670114
Ν	0.442251	1.188841	0.835476
Н	-1.239797	-0.791499	1.514162
$\mathtt{HA}_{\mathtt{zw}}$	A _{syn}		
С	-1.397132	-0.972009	0.024435
С	-0.290939	-0.608987	-0.973725
С	-1.138297	0.474609	-0.318433
0	-1.931423	1.280461	-0.954854
Н	-0.531064	-0.742415	-2.023820
S	1.454311	-0.939039	-0.685924
С	1.925386	0.389800	0.483/63
H	1.931953	0.003840	1.50/869
H	2.947253	0.676242	0.230037
С	1.034931	1.612126	0.397614
H	0.981/58	2.014244	-0.616206
H	1.390333	2.386606	1.0/91//
Н	-0.929273	2.093728	0.886594
Н	-0.3614/8	0.//5434	1.694905
N	-0.35/515	1.250300	0./86919
H	-2.304/11	-1.34/410	-0.444198
C	-1.05/366	-1.62/511	1.342881
H	-1.800/58	-1.38/298	2.109279
H	-1.02961/	-2./150/9	1.22/300
н	-0.068884	-1.33/228	1./15036
HAzw	В		
С	-1.488674	0.215797	1.199323
С	-0.806302	-0.599667	0.100497
С	-0.742892	0.926106	0.093075
0	-1.215847	1.710790	-0.821123
S	0.738549	-1.447129	0.494593
С	1.727787	-0.484425	-0.700251
Η	2.739624	-0.891735	-0.702995
Η	1.294479	-0.643526	-1.689553
С	1.763391	1.016686	-0.363457
Η	1.622924	1.617812	-1.261563
Η	2.700926	1.301253	0.114854
Η	0.599498	2.406296	0.650717
Η	0.877306	1.020827	1.507341
Ν	0.662212	1.387859	0.577491
Н	-1.089252	0.157693	2.208646
С	-1.663045	-1.270299	-0.947022
Н	-1.087332	-1.455197	-1.860272
Н	-2.045971	-2.227394	-0.580567
Η	-2.507339	-0.626048	-1.205345
Н	-2.571955	0.267566	1.138805

на	A _{anti}		
С	-1.279393	-0.505235	0.663268
С	-0.342410	-0.740772	-0.504322
0	-0.703279	0.656817	-0.106/63
C	-2.734567	-0.884837	0.536079
Н	-2.874265	-1.949286	0.745127
Н	-3.350683	-0.315046	1.238098
Η	-3.095851	-0.680361	-0.474819
H	-0.820799	-1.083804	-1.417992
S	1.328894 2.106921	-1.396622	-0.3//203
Н	2.190021 2.141613	-0.023884	1 551354
Н	3.241956	-0.111647	0.161889
С	1.629179	1.339330	0.086485
Н	1.676781	1.481542	-0.998275
Η	2.237304	2.119886	0.552590
N	0.245586	1.521019	0.517369
н ч	-0.835435 -1.888924	-0.653191	-0 611957
н	0.170975	1.489685	1.529095
עז	7		
HA C	A _{syn} -1 284171	-1 034727	0 065829
C	-0.283296	-0.546362	-0.969002
С	-1.074114	0.433966	-0.138358
0	-2.154831	1.055161	-0.790833
Н	-0.591363	-0.619939	-2.006525
S	1.497316	-0.771941	-0.812039
C u	1.922964	0.429679	1 475396
н	2.940094	0.761406	0.280266
C	0.965779	1.615771	0.522655
Н	0.994988	2.131121	-0.443594
Н	1.298424	2.326069	1.284748
Ν	-0.417438	1.258468	0.814293
H	-1.844242	L.906982	-1.1269//
н Н	-0.550286	$-1 \ 401072$	-0 366989
C	-0.842431	-1.744953	1.320777
Н	-1.604558	-1.663717	2.101365
Н	-0.675880	-2.806201	1.115130
Η	0.092605	-1.337793	1.713653
на	В		
С	1.005211	0.061191	1.395343
С	0.555413	0.726482	0.110751
C	0.851515	-0.756011	0.147019
S	2.075480 -1 134482	-1.165889	-0.406491
C	-2.105971	-0.159765	0.167991
Н	-2.220149	-0.392530	1.231410
Η	-3.093318	0.059208	-0.245798
С	-1.447474	-1.322786	-0.566126
H	-1.288841	-1.048478	-1.614909
H N	-2.11323/ -0 177461	-2.190104 -1 736407	-0.340685
H	1.888577	-1.508454	-1.291314
Н	-0.312435	-2.252942	0.878613
С	1.529053	1.645032	-0.596248
Η	1.331714	1.664579	-1.672217
H	1.436566	2.664609	-0.208638
H IJ	2.553477	1.305132	-0.435680
H	2.013863	0.281387	1.734014
	_		
HA -	A _{anti}		0 500401
Ċ	-1.4026/6	-0.44959/	0.389421

С	-0.376780	-0.685002	-0.511534	С	2.1799
С	-0.741028	0.747348	-0.068623	Н	3.1900
0	-1.397794	1.571322	-0.875470	Н	2.2519
Н	-0./88112	-0.951/8/	-1.482131	C	1.3/16
ъ С	2 296593	-1.4/32/1	-0.211101	п	1 2121
н	2.290393	-0 148091	1 293510	N	0 0856
н	3.209344	-0.140170	-0.359611	Н	0.2204
С	1.617421	1.285248	-0.001566	H	-0.6530
Н	1.479697	1.448937	-1.082582	С	-2.5607
Н	2.283137	2.070229	0.372042	Н	-2.5436
Н	0.078271	2.313843	0.843358	Н	-3.1996
Ν	0.348467	1.340305	0.725362	Н	-3.0105
Н	-1.037631	-0.687656	1.587578		
С	-2.84866/	-0./84882	0.321331	TS3 A	Asyn 1 CEOC
н ц	-3.038946	-1.859524	1 01/001	C	-1.6535
н	-3 123043	-0.237900 -0.481149	-0 692666	C	-0.4387
	3.120010	0.101119	0.092000	0	-0.7539
HA-	Asvn			H	-0.4191
С	-1.438302	-0.900567	-0.005695	S	0.9186
С	-0.305058	-0.528126	-0.961353	С	2.1019
С	-1.087494	0.564121	-0.210816	Н	3.0157
0	-1.904146	1.378149	-0.876901	Н	2.3377
H	-0.536802	-0.571327	-2.021312	С	1.5327
S	1.415592	-0.964053	-0.660526	H	L./392
Ч	1 904/12	-0 024441	1 496705	п	2.01/4
Н	2.977747	0.645527	0.172835	H	-0.1000
С	1.035014	1.573230	0.433988	H	-2.4432
Н	1.039380	2.008641	-0.579534	С	-2.2034
Н	1.424357	2.328452	1.124822	Н	-2.7693
Н	-0.831928	2.018018	1.110322	Н	-1.4062
Ν	-0.305213	1.183418	0.867261	Н	-2.8695
H	-2.349715	-1.206428	-0.519199		
C	-1.138096	-1.663091	1.259852	TS3 E	3 0 000/
п u	-1.907373	-2 727945	1.970910	C	0.9904
Н	-0.241562	-1 268163	1 744230	C	1 0147
	0.011000	1.200100	1,11200	0	2.0631
HA-	В			S	-1.1732
С	-1.294511	-0.194613	1.314031	С	-1.9396
С	-0.589541	-0.724352	0.068072	Н	-3.0227
С	-0.941577	0.770849	0.195762	Н	-1.6936
0	-1.871281	1.322631	-0.574923	С	-1.4878
S	1.114612 2 135164	-1.321/25	0.207742	H	-2.1213
н	2 689430	0.100377	0.761258	N	-0 1070
Н	2.845290	-0.090928	-0.936699	Н	-0.0019
С	1.278650	1.357480	-0.554778	Н	0.2549
Н	0.839971	1.180969	-1.550903	С	1.5347
Н	1.928870	2.235514	-0.627357	Н	2.0984
Η	-0.078572	2.551419	0.380336	Н	1.0507
Ν	0.251299	1.592702	0.458725	H	2.2933
Н	-0.716020	-0.166520	2.233137	Н	1.9871
U U	-1.409394	-1.543964	-0.902550	Ͳ 7- ۸	
Н	-1.445798	-2.598988	-0.606175	т и А С	0.0572
Н	-2.429124	-1.152764	-0.936111	C	-2.1077
Н	-2.341128	-0.473354	1.422765	C	1.0006
				С	-1.1967
TS3	A _{anti}			С	1.1320
C	-1.148328	-0.469522	0.549082	H	-3.1366
C	-0.234926	-0.992624	-0.545045	H	-1.6553
C	-U.946U58 -1 782000	U.198239 1 273604	-U.IXIYIY -1 012702	H u	-2.0862
Н	-0.676907	-1.377928	-1.457492	0	2.1059
S	1.434166	-1.371454	-0.202344	N	0.1164

С Н Н С Н Н Н Н Н Н Н		2. 3. 2. 1. 1. 0. 0. -0. -2. -3.	179997 190085 251976 371669 956684 212145 085637 220449 653066 560732 543682 199685 010586	0.302374 0.181492 0.724775 1.237238 2.142648 0.753120 1.688693 2.357600 -0.496907 -1.019404 -2.053611 -0.428687 -1.007447	-0.132320 0.272228 -1.140977 0.771617 0.965094 1.741440 0.237363 -0.515326 1.526127 0.609937 0.969282 1.276017 -0.386295
Г ЗСССОНЗСННСННИННСННН	Asyn	-1. -0. -0. -0. 2. 3. 2. 1. 1. 2. -0. -2. -2. -2. -2. -2. -2. -2.	653935 438703 729149 753901 419128 918676 101944 015769 337773 532789 739207 017416 081837 100099 443209 203462 769304 406210 869588	-0.174889 -0.743185 0.974668 1.976563 -0.679288 -1.494374 -0.068363 -0.451135 0.222110 1.121041 2.056357 1.178951 1.058475 0.269663 0.032394 -0.889403 -1.771374 -1.246060 -0.242668	-0.269262 -0.988639 -0.198739 -0.995019 -2.071361 -0.203195 -0.073067 0.390789 -1.099636 0.727704 0.200957 1.706220 0.972565 1.583963 -0.994890 0.953214 0.633529 1.612361 1.534113
rs3 C C	В	0.	990451 521726	-0.018777 0.968778 -0.983287	1.254078 0.205135 0.141568
СОЅСННСННИНСНННН		$\begin{array}{c} 1 \\ 1 \\ 2 \\ -1 \\ -1 \\ -1 \\ -1 \\ -1 \\ -1$	014762 063145 173267 939617 022776 693642 487818 121303 633919 107092 001968 254942 534790 098488 050725 293320 987126	-1.227538 1.364238 -0.137967 -0.017011 -0.209895 -1.404268 -2.244242 -1.272366 -1.820144 -2.196709 -0.208656 1.726667 1.070035 2.502367 2.228478 0.174621	-0.526625 0.149921 -0.583392 -0.478525 -1.648910 0.147130 -0.157040 1.224891 -0.104052 -1.041896 2.042472 -0.599122 -1.289148 -1.201954 0.027982 1.672384

	0 000001	0 550017	0 207470		0 700000	0 545706	0 000040
н	0.222231	2.33821/	-0.39/4/2	Н	-0.733630	-2.545/86	0.989342
S	-1.661654	-1.233265	-0.126919	Н	0.952431	-2.371783	1.522924
н	0 264650	-1 198279	-1 543244	н	-0 194655	-1 037731	1 739170
11	0.204030	1.190279	1.010211	11	0.194055	1.057751	1.755170
Н	0.553812	-0.559051	1.419152	Н	-0.3925/1	-0.169/1/	-1.9415/4
С	2.387959	-1.246571	0.528191	Н	0.853223	-2.153872	-0.920318
ц	3 083773	-0 623175	1 101202				
11	0.000110	0.025175	1.101292		-		
Н	2.300113	-2.220685	1.015854	TS4	A _{anti}		
Н	2.815880	-1.399017	-0.467344	С	-1.457324	-0.285200	0.666876
				C	-0 606452	-0 330630	_0 525350
				C	0.000452	0.550050	0.525555
TZ- B				C	-0.410548	1.110706	-0.297622
С	0.419416	-1.395812	-0.367255	0	-0.951594	2.027192	-0.997001
C	1 443674	1 211241	0 567400	н	-1 186521	-0 484850	-1 437115
0	1.1130/1	1.211241	0.307400	11	1.100521	0.404050	1.45/115
C	-0./53836	-0./24282	-1.0892/8	S	0.913/31	-1.482830	-0./06259
С	0.271924	1.930937	-0.093451	С	1.832920	-0.695658	0.654877
C	-1 581373	0 197557	-0 200981	н	1 338908	-0 921917	1 606421
	1.001070	1 0057001	0.200901	11	1.990900	1 125101	1.000121
Н	2.341297	1.835/01	0.201868	Н	2.832457	-1.135191	0.6/04//
Н	0.237006	2.971815	0.241006	С	1.925354	0.813900	0.465465
н	-1 475060	-1 471783	-1 444018	н	2 385108	1 017264	-0 515410
	1.000007	1 041410	1 000700	11	2.505100	1 000000	1 005714
Н	1.222297	1.041418	1.626/86	Н	2.584368	1.228098	1.235/14
Н	0.412543	1.943828	-1.180602	Н	0.701332	2.473257	0.630544
0	-2 712354	-0 123891	0 196143	N	0 620168	1 462355	0 617621
N	1 025122	1 205000	0.225442	11	0.050177	0.054000	1 (20400
IN	-1.035125	1.303210	0.225445	н	-0.9581//	-0.254922	1.030400
Н	-1.615616	1.878172	0.876700	С	-2.844112	-0.873740	0.601675
S	1.863626	-0.377893	-0.296179	н	-2.853438	-1,969444	0.455976
C	0.004074	1 022201	0.0000077	11	2 412514	0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	1 514500
C	-0.004274	-1.933291	0.992237	н	-3.413514	-0.00/92/	1.514590
Н	-0.841802	-2.637989	0.883281	Н	-3.417676	-0.449374	-0.234254
Н	0.813846	-2.474218	1,481078				
 TT	0 250204	1 160246	1 71 27 / 7	me 4	7		
п	-0.330294	-1.100240	1./13/4/	154	Asyn		
Н	-0.390731	-0.163845	-1.960253	C	1.176800	1.422110	0.051378
				С	0.453787	0.536724	-0.890662
ጥ 7 7				C	1 13/551	-0 622170	_0 278218
12 7	0 0 0 0 0 0 0 0	1 0000000	0 505050	C	1.134331	0.022170	0.270210
C	-0.0593/3	-1.3223/2	-0.505259	0	2.130112	-1.210664	-0.801216
С	-2.033108	0.677108	-0.224521	Н	0.804293	0.652160	-1.917933
C	0 964067	-0 617700	0 410281	S	-1 435931	0 495942	-1 035487
Č	1 000001	1 510005	0.410201	5	1 701 600	0.40042	1.033407
C	-1.089084	1.519985	0.632038	С	-1./81680	-0.3313/8	0.553889
С	1.223126	0.798705	-0.096972	Н	-1.659934	0.372511	1.379885
н	-3 064826	0 853469	0 091143	н	-2 826540	-0 648438	0 518131
	1 515700	0.000100	0.001110		2.020010	1 546510	0.010101
н	-1.515/39	2.518903	0./43361	C	-0.889296	-1.546510	0./6391/
Н	-1.943912	0.968553	-1.275389	Н	-1.071336	-2.265178	-0.051541
н	-1.003050	1.091185	1.635824	н	-1.170928	-2.030806	1.704476
0	2 202007	1 105272	0 644216	11	1 001172	1 060127	1 170210
0	2.203907	1.1033/3	-0.044310	п	1.0911/2	-1.909127	1.1/2310
N	0.230671	1.708610	0.044777	N	0.527345	-1.182908	0.866337
Н	0.435246	2.613298	-0.360713	Н	1.801150	2.171896	-0.438242
C	_1 000515	_1 110002	_0 053057	C	0 575900	1 0/06/0	1 361/72
3	-1.009515	-1.110902	-0.033937	C	0.575800	1.040040	1.301472
Н	0.091384	-1.030302	-1.550362	Н	1.248357	2.543830	1.877170
Н	0.542749	-0.556518	1.421012	Н	-0.398323	2.364695	1.264257
C	2 268787	-1 409818	0 457582	н	0 404909	0 994509	2 033979
	2.200707	1.102010	1 050000	11	0.101000	0.001000	2.0000779
н	3.023200	-0.888615	1.020839				
Н	2.086401	-2.390381	0.904514	TS4	В		
Н	2.670222	-1.552966	-0.549254	С	-0.999799	-0.097087	1.561389
ц	0 00811/	-2 102902	-0 115171	C	-0 739519	-0 504761	0 1589/3
11	0.090114	2.402902	0.4434/1	C	0.759519	0.304701	0.130943
				С	-0.734157	0.964130	-0.055001
TZ B				0	-1.664162	1.613582	-0.634096
C	0 481366	-1 321965	-0 314200	S	0 791155	-1 501638	-0 303489
ä	1 01000	1 004000	0.511200	~	0.010001	1.001000	0.000000
C	1.316/52	1.264682	0.56/99/	C	2.018991	-0.286994	0.260996
С	-0.729231	-0.733469	-1.065916	Н	2.003834	-0.227164	1.354757
C	0.123434	1,934901	-0.111225	н	2.999220	-0.650750	-0.054962
Č	_1 607014	0 100100	_0 100004	~	1 705070	1 000000	-0 242502
C	-1.02/314	0.122160	-0.190004	C	1.135913	T.000308	-0.342396
Н	2.168825	1.949741	0.560981	Н	1.721830	0.984139	-1.440787
Н	0.061056	2.970521	0.229238	Н	2.548298	1.765903	-0.077218
 []	_1 257607	_1 557405	_1 /11770	11	0 260064	2 600000	_0 10(120
п	-1.33/08/	-1.35/495	-1.411/19	Н	0.309834	2.009090	-0.100130
H	1.078860	1.040203	1.611802	N	0.497653	1.647781	U.190546
Н	0.269877	1.957997	-1.196386	Н	-0.164138	0.188100	2.190051
0	-2 739215	-0 278512	0 163750	C	-1 916009	-1 215413	-0 485239
	1 1 5 6 4 5 6	1 204012	0.10010		1 000000	1 0000000	1 575002
IN	-1.1594/2	1.324919	0.213319	Н	-1.82/826	-1.22//56	-1.5/589/
Н	-1.780540	1.834340	0.829296	Н	-1.968607	-2.247335	-0.121727
S	1.941746	-0.216466	-0.284490	Н	-2.843683	-0.704561	-0.219076
Ċ	0 107000	_1 0/0500	1 060224	11	_1 060401	_0 551754	2 05/005
\cup	0.10/000	エ・ロサラフムの	1.009004	п	1.000421	0.001/04	2.004000

(E)-	Crotonamide		
С	-3.195928	-0.126680	-0.007110
С	-1.891630	-0.406426	0.048966
С	-0.863653	0.669015	-0.056305
0	-1 158760	1 866097	-0 181964
ц	-1 5//808	-1 /30775	0 173407
	2 160700	1 210040	0.106101
3	3.100/90	-1.210940	-0.100191
C	2.788063	0.449353	0.512428
Н	2.621492	0.368215	1.592784
Н	3.639156	1.121061	0.365505
С	1.563771	1.110620	-0.115986
Η	1.758245	1.329692	-1.175109
Н	1.330464	2.055260	0.385435
Н	0.621845	-0.755434	0.005228
Ν	0.410277	0.238422	0.007058
н	-3.482715	0.916989	-0.132240
C	-4 292821	-1 138383	0 089641
ц	-1 9/8835	_0 911060	0.0000041
11	4.017077	1 111000	0.950572
н	-4.91/8//	-1.111986	-0.808984
Н	-3.898287	-2.149524	0.213155
(Z)-	Crotonamide		
С	3.106150	-1.039999	0.131219
С	1.776032	-0.937811	0.257739
С	0.961654	0.297038	0.063511
0	1.442807	1.430219	-0.078646
Н	1.215644	-1.833453	0.516154
S	-3.308777	-0.906151	-0.238569
С	-2.703094	0.694845	0.434118
Н	-3,422452	1,491094	0.220156
Н	-2.608576	0.627182	1.524096
C	-1 354795	1 123003	-0 140620
ч	-0 995500	2 034986	0 346760
ц	-1 454391	1 332304	_1 21/000
11	0 7/1702	1.332304	0 070602
п	-0.741793	-0.005595	0.070000
IN	-0.369538	0.079665	0.073388
C	4.102948	0.01/652	-0.231149
Н	3.625633	0.948824	-0.527060
Н	4.759160	0.211487	0.625421
Н	4.745941	-0.347555	-1.038406
Н	3.527699	-2.028628	0.311432
Metn		0 000011	0 000770
C	3.491//6	0.222211	-0.009779
C	2.18/360	0.504640	0.02/2/4
C	1.21/093	-0.650835	-0.052972
0	1.600692	-1.824844	-0.144258
S	-2.901615	0.930098	-0.221241
С	-2.436500	-0.695263	0.504763
Н	-2.288164	-0.588163	1.585422
Н	-3.242512	-1.420357	0.358319
С	-1.164549	-1.285233	-0.100983
Н	-1.336247	-1.551397	-1.153077
Н	-0.861301	-2.192417	0.430792
Н	-0.388860	0.648408	-0.033654
Ν	-0.084405	-0.320289	-0.005460
Н	3.828942	-0.805286	-0.095983
Н	4.238376	1.008350	0.045430
C	1 655512	1 909253	0 143912
н	1 036911	2 177889	-0 720018
н Ц	1 0400/1	2 021160	1 041903
п u	1.040041 2 /70706	2.031109	1.041903 A 100719
11	2.4/9/90	2.022310	U • I J J / I Z
мт			
<u>с</u>	0 00000	0.00000	-1.130939
н	-0 000000	1.019044	-1.528954
н	0 882518	-0 509522	-1 528954
Н	-0 882518	-0.509522	-1.528954
± ±	0.002010	0.00/022	エ・シムシンフィ

S		0.000000	0.000000	0.710781
TS1'	A			
СССОСННН SСННН		$\begin{array}{c} 1.278244\\ 0.454076\\ 1.157182\\ 1.410153\\ 1.807304\\ 0.972631\\ 2.447415\\ 2.378458\\ 0.196154\\ -1.944072\\ -1.622804\\ -2.053159\\ -0.537017\\ -2.024575\end{array}$	0.441555 - 0.264543 - 0.919651 - 1.976169 1.790566 2.474340 1.767358 2.196626 - 0.309630 - 0.030016 0.399288 1.369555 0.460777 - 0.352702	-0.209496 -1.075776 -0.018397 0.585337 0.113958 0.304442 0.997416 -0.726489 -2.124557 -0.510294 1.228182 1.489386 1.398840 1.912142
TS1' C C C C S C H H C H H H H H	В	1.103387 0.830564 1.324401 1.673798 -1.600000 -1.960478 -2.785658 -1.072654 1.018794 0.673462 0.480337 2.088502 1.136630 -2.211006	-0.375331 0.550921 -0.756137 -1.580262 0.511848 -1.070926 -0.978867 -1.399874 2.005485 2.262224 2.613064 2.234099 -0.546323 -1.855877	1.223091 0.225096 -0.085432 -0.942439 -0.407920 0.410992 1.121755 -0.050943 -1.053495 0.678855 0.024507 2.291278 -0.307661
СР'- ССССС И Н Н Н Н Н Н	A	-1.042207 -0.132025 -1.499208 -2.334777 -1.119198 -0.415196 -2.122985 -0.884445 0.048093 1.391576 2.432219 2.570055 1.972140 3.407835	0.706341 - 0.317360 - 0.535948 - 1.455247 2.172372 2.498347 2.460750 2.740384 - 0.392673 - 0.878650 0.594995 0.792956 1.464880 0.413326	0.077048 0.663097 0.052103 -0.198577 -0.166692 -0.942535 -0.492811 0.741910 1.744820 -0.220482 0.019411 1.085191 -0.454114 -0.435930
СР'- СССО ССО К Н Н С Н Н Н Н Н Н	В	0.935230 0.432073 1.236165 1.757319 -1.405013 -1.862022 -2.931125 -1.656795 0.943079 0.733613 0.483282 2.026526 0.969464 -1.310466	-0.340053 0.529988 -0.722348 -1.487246 0.530423 -1.155519 -1.274172 -1.318959 1.933621 2.241621 2.670174 1.962415 -0.451826 -1.892187	1.255652 0.169688 0.010518 -0.849043 -0.300176 0.178433 -0.005875 1.238102 -0.115134 -1.146669 0.555456 0.038505 2.331295 -0.410601

CP' A_{anti}

СССОНЅСНННСННН	-1.224469 -0.151297 -0.749551 -0.732221 -0.530614 1.574731 2.080328 3.137287 1.528358 1.958861 -0.837169 -2.588230 -2.554767 -3.332792 -2.907776	-0.282401 -0.407504 0.880302 2.076906 -0.868367 -0.825957 0.502132 0.333631 0.459733 1.480042 -0.332937 -0.906034 -1.973047 -0.430386 -0.787574	0.521447 -0.612206 -0.236852 -0.368934 -1.523636 -0.341933 0.789981 0.999951 1.730409 0.321682 1.541246 0.304641 0.537629 0.947145 -0.734082
СР' 2 С С С С С О Н В С С Н Н Н Н Н Н Н Н Н	A _{syn} 1.579125 0.050516 0.807105 0.732876 -0.141888 -1.189242 -2.439264 -3.267334 -2.023640 -2.800359 2.196205 2.090317 3.096960 2.130986 1.447147	-0.175577 -0.033901 1.018142 2.056444 0.063146 -0.924216 0.388708 -0.023764 1.250709 0.682956 -0.109119 -1.111975 -0.813215 -2.136542 -1.090646	0.499044 0.871004 -0.431037 -0.070828 -0.213108 -0.791407 -0.736166 0.773335 1.393593 -0.576463 -0.877782 -0.197546 -1.458206
СР' н ССССО КССО КССО КССО КССО КССО КССО КС	B 0.847136 0.263437 1.538485 2.492199 -1.091377 -2.384293 -3.266674 -2.630741 0.257305 0.253068 -0.629880 1.141563 -2.066509 0.358940 1.232247	0.018926 0.376498 -0.333037 -0.837268 -0.657490 -0.122524 -0.728923 0.929812 1.832899 1.911748 2.336147 2.349749 -0.296356 -0.770224 0.849459	$\begin{array}{c} 1.364124\\ -0.052340\\ 0.124167\\ -0.407515\\ -0.630099\\ 0.526882\\ 0.317414\\ 0.380824\\ -0.473665\\ -1.563025\\ -0.078675\\ -0.089629\\ 1.557003\\ 1.930825\\ 1.951956\end{array}$
MA C H H N H H	-0.708121 -1.118455 -1.061351 -1.118453 0.751111 1.144607 1.144607	0.000000 -0.881672 -0.00007 0.881679 -0.000000 0.811873 -0.811873	0.017287 -0.480981 1.056829 -0.480969 -0.125811 0.341037 0.341037
TS1' C C C C C H H H	A 1.034503 0.087819 0.710405 0.820538 1.829064 2.146511 1.231547 2.712473	-0.363758 0.104845 0.966064 2.121397 -1.553312 -2.123929 -2.218459 -1.254643	0.122030 1.034380 0.064024 -0.387610 -0.283519 0.593450 -0.915638 -0.851123

Н С Н Н Н Н Н	-0.113620 -1.927356 -2.912815 -1.167660 -1.732343 -1.809719 -2.052281 -2.414693	0.024858 -0.360627 -0.695624 -1.040214 0.636615 -0.319289 -1.213380 0.389342	2.096544 -0.903230 -1.235354 -1.297493 -1.301972 0.553607 0.974020 0.961088
TS1' _№	В		
С	0.760714	-0.183064	1.254778
С	0.385100	0.604831	0.160530
С	1.129780	-0.624626	0.013471
0	1.688523	-1.420638	-0.760736
С	-1.957512	-1.006146	0.135548
Н	-3.013340	-1.195271	-0.071391
Н	-1.794505	-1.016165	1.215964
С	0.451357	2.050876	-0.220293
Н	0.159593	2.208708	-1.260827
Н	-0.185084	2.659793	0.427395
Н	1.484677	2.388096	-0.097444
Н	0.772079	-0.234380	2.334984
Н	-1.355451	-1.801364	-0.308477
Ν	-1.508331	0.280233	-0.393837
Н	-2.088030	1.043650	-0.051244
Н	-1.546439	0.299188	-1.410417

6. Kinetic study for the reaction of cyclopropenone probe 1 with cysteine



2-phenethylcycloprop-2-en-1-one **1** in CH₃CN (160 μ L of 25 mM solution), L-cysteine ethyl ester hydrochloride in H₂O (40 μ L of 0.1 M solution) and the internal standard 2-phenylacetamide in CH₃CN (200 μ L of 15 mM solution) were added to CH₃CN (640 μ L) and H₂O (936 μ L) at 4 °C (in a cold room). The reaction was then initiated by adding Na₂CO₃ in H₂O (24 μ L of 0.25 M solution) affording the starting reactant concentrations of 1.5 mM (2-phenylacetamide), 2 mM (for cysteine and cyclopropenone **1**) and 3 mM (for Na₂CO₃), and the reaction mixture was stirred at 4 °C. 125 μ L of reactant mixture was quenched at different time points with 1 μ L of 1M HCl solution. The sample was then analyzed by LC–MS as described earlier (page S3). The concentration of 2-phenethylcycloprop-2-en-1-one **1** was then determined at each time point and used to calculate the rate constant as shown in **Figure S5**. Applying the similar kinetic analysis reported for the CBT-cysteine reaction,⁹ the resulted 1/[concentration] values for cyclopropenone **1** were plotted against the time and fitted to a linear equation to obtain the second order constant *k* from the slope. The experiment was repeated three times. Data were processed using GraphPad Prism (GraphPad Software, La Jolla, CA, USA).

In a separate experiment, LC–MS analysis was used to confirm that the reaction did not proceed after the quenching with 1 μ L of 1M HCl solution, and that the reaction did not start before the initiation with the addition of Na₂CO₃.





Figure S5. a) Typical chromatograms of the reaction mixture at different time points. b) Ratio of cyclopropenone 1 peak area to the peak area of the internal standard was uses to determine cyclopropenone concentration at different time points. c) Concentration of cyclopropenone 1 vs time for reaction of probe 1 (2 mM) with L-cysteine ethyl ester (2 mM) in CH₃CN/H₂O at 4 °C. Error bars indicate the standard deviations of three replicate experiments. D) Linear regression analysis of 1/[cyclopropenone] vs. time of the cyclopropenone 1 reaction with cysteine. Error bars indicate the standard deviations of three replicate experiments.

7. Stability studies of cyclopropenone probe 1 in buffer

To determine stability in aqueous buffer, cyclopropenone probe 1 (5 mM) was dissolved in 50% CD_3CN/NaP_i buffer (50 mM, pH 7 or 8), incubated at 37 °C, and monitored by ¹H NMR for one week.



Figure S6. Cyclopropenone **1** is stable in 50% CD₃CN/NaP_i buffer (50 mM, pH 7) at 37 °C for at least 7 days. *H/D exchange was observed for cyclopropenone-ring proton.



Figure S7. Cyclopropenone **1** is stable in 50% CD₃CN/NaP_i buffer (50 mM, pH 8) at 37 °C for at least 7 days. *H/D exchange was observed for cyclopropenone-ring proton.

8. Stability of cyclopropenones towards nucleophiles

Selectivity studies and competition experiments. General method

Preparation of stock solutions

To determine stability of cyclopropenones towards biological nucleophiles, cyclopropenone probe **1** was incubated with various nucleophiles under conditions used for the reaction with cysteine. To this end, stock solutions of L-cysteine ethyl ester hydrochloride (100 mM in H₂O), L-lysine ethyl ester hydrochloride (100 mM in H₂O), L-tyrosine methyl ester hydrochloride (100 mM in H₂O), L-tyrosine methyl ester hydrochloride (100 mM in H₂O), L-tyrosine (100 mM in H₂O), N-(tert-butoxycarbonyl)-L-cysteine methyl ester (100 mM in CH₃CN), cyclopropenone **1** (50 mM in CH₃CN), and Na₂CO₃ (0.25 M in H₂O) were prepared.

Control experiment

Cyclopropenone 1 stock (10 μ L, 0.50 μ mol) was mixed with CH₃CN (52.5 μ L), H₂O (61 μ L), Na₂CO₃ stock solution (1.5 μ L, 0.375 μ mol), incubated at 4 °C, and analyzed by LC–MS after 30 min and 24 h. LC–MS analysis was performed as described on page S3.

In a separate experiment, L-cysteine ethyl ester hydrochloride stock solution (2.5 μ L, 0.25 μ mol, 1 equiv.) was mixed with CH₃CN (52.5 μ L), H₂O (58.5 μ L), cyclopropenone **1** (10 μ L, 0.50 μ mol, 2 equiv.), Na₂CO₃ (1.5 μ L, 0.375 μ mol, 1.5 equiv.), incubated at 4 °C, and analyzed by LC–MS after 1 h and 24 h. LC–MS analysis was performed as described on page S3.

Stability of cyclopropenone 1 towards lysine, threonine, tyrosine and serine

Next, nucleophile stock solution (2.5 μ L, 0.25 μ mol, 1 equiv.) was mixed with CH₃CN (52.5 μ L), H₂O (58.5 μ L), cyclopropenone **1** (10 μ L, 0.50 μ mol, 2 equiv.), Na₂CO₃ (1.5 μ L, 0.375 μ mol, 1.5 equiv.), incubated at 4 °C, and analyzed by LC–MS after 30 min and 24 h. LC–MS analysis was performed as described on page S3.

For competition experiments, nucleophile stock solution (1.25 μ L, 0.125 μ mol, 1 equiv.) and L-cysteine ethyl ester hydrochloride stock solution (1.25 μ L, 0.125 μ mol, 1 equiv.) were mixed with CH₃CN (52.5 μ L), H₂O (58.5 μ L), cyclopropenone **1** (10 μ L, 0.50 μ mol, 4 equiv.), Na₂CO₃ (1.5 μ L, 0.375 μ mol, 3 equiv.), incubated at 4 °C, and analyzed by LC–MS after 30 min. LC–MS analysis was performed as described on page S3.

Stability of cyclopropenone 1 towards thiol nucleophiles (glutathione, N-(tert-butoxycarbonyl)-protected cysteine)

Next, nucleophile stock solution (2.5 μ L, 0.25 μ mol, 1 equiv.) was mixed with CH₃CN (52.5 μ L), H₂O (59.5 μ L), cyclopropenone **1** (10 μ L, 0.50 μ mol, 2 equiv.), Na₂CO₃ (0.5 μ L, 0.125 μ L, 0.5 equiv.)*, incubated at 4 °C, and analyzed by LC–MS after 30 min and 24 h. LC–MS analysis was performed as described on page S3.

For competition experiments, nucleophile stock solution (1.25 μ L, 0.125 μ mol, 1 equiv.) and L-cysteine ethyl ester hydrochloride stock solution (1.25 μ L, 0.125 μ mol, 1 equiv.) were mixed with CH₃CN (52.5 μ L), H₂O (59.5 μ L), cyclopropenone **1** (5 μ L, 0.25 μ mol, 2 equiv.), Na₂CO₃ (1 μ L, 0.250 μ L, 2 equiv.),* incubated at 4 °C, and analyzed by LC–MS after 30 min. LC–MS analysis was performed as described on page S3.

^{*}compared to the experiment above, 1 equiv. less of Na₂CO₃ was used to account for the neutralization of hydrochloride in stock solutions of L-cysteine, L-lysine, L-threonine, L-tyrosine and L-serine.

Control experiment



Figure S8. a. 2-Phenethylcycloprop-2-en-1-one **1** is stable under reaction conditions for at least 24 h. **b.** Cysteine reacts with 2-phenethylcycloprop-2-en-1-one forming products **2a–d** after incubation for 30 min. After 24 h, ester hydrolysis is observed (products **2e–h**) while thiazepanone ring is stable.

Selectivity in the presence of lysine



Figure S9. a. 2-Phenethylcycloprop-2-en-1-one **1** shows no reactivity with lysine after 30 min or 24 h incubation. **b.** Lysine does not interfere with the reaction between 2-phenethylcycloprop-2-en-1-one **1** and cysteine.

Selectivity in the presence of tyrosine



Figure S10. a. 2-Phenethylcycloprop-2-en-1-one **1** shows no reactivity with tyrosine after 30 min or 24 h incubation. **b.** Tyrosine does not interfere with the reaction between 2-phenethylcycloprop-2-en-1-one **1** and cysteine.

Selectivity in the presence of threonine



Figure S11. a. 2-Phenethylcycloprop-2-en-1-one **1** shows no reactivity with threonine after 30 min or 24 h incubation. **b.** Threonine does not interfere with the reaction between 2-phenethylcycloprop-2-en-1-one **1** and cysteine.

Selectivity in the presence of serine



Figure S12. a. 2-Phenethylcycloprop-2-en-1-one **1** shows no reactivity with serine after 30 min or 24 h incubation. **b.** Serine does not interfere with the reaction between 2-phenethylcycloprop-2-en-1-one **1** and cysteine.

Selectivity in the presence of glutathione



Figure S13. a. 2-Phenethylcycloprop-2-en-1-one **1** shows no reactivity with glutathione after 30 min or 24 h incubation. **b.** Glutathione does not interfere with the reaction between 2-phenethylcycloprop-2-en-1-one **1** and cysteine.

Selectivity in the presence of N-(tert-butoxycarbonyl)-protected cysteine



Figure S14. a. 2-Phenethylcycloprop-2-en-1-one 1 reacts with N-(tert-butoxycarbonyl)-protected cysteine forming a complex mixture of products. b. Even in the presence of N-(tert-butoxycarbonyl)-protected cysteine, 2-phenethylcycloprop-2-en-1-one 1 predominantly reacts with cysteine with free amino group.

9. Peptide modification

General procedures and characterization methods

LC–MS method for analysis of peptide conjugation

LC–MS was performed on a Waters SQ Detector 2 mass spectrometer coupled to an Acquity UPLC system using an Acquity UPLC BEH C18 column (130Å, 1.7 μ m, 2.1 mm X 50 mm). Solvents A, water with 0.1% formic acid and B, 71% acetonitrile, 29% water and 0.075% formic acid were used as the mobile phase at a flow rate of 0.2 mL min⁻¹. The gradient was programmed as follows: 100% A for 2 min, then 100% A to 100% B in 9 min, then 100% B for 5 min, and after that 100% A for 4 min. The electrospray source was operated with a capillary voltage of 3.0 kV and a cone voltage of 30 V. Nitrogen was used as the desolvation gas at a total flow of 800 L h⁻¹. *m/z* values of positively and negatively charged ions were measured in the mass analyzer, which was scanned between *m/z* 100–2000 for the generation of mass spectra. The major peak(s) were selected for integration and analyzed using MassLynx software (v. 4.1 from Waters).

LC-MS/MS analysis of modified peptides

Peptide solutions were prepared to approximate concentrations of 1 pmol/uL in 50:50 acetonitrile:water + 0.1% formic acid. The solutions were pulled into a syringe and loaded into the mass spectrometer source by means of a syringe pump.

All experiments were performed using an LTQ Orbitrap Velos hybrid ion trap mass spectrometer (Thermo Scientific, Waltham, MA). m/z values of positively charged ions were measured in the Orbitrap mass analyzer, set at a resolution of 30000 which was scanned between m/z 380–1500 for the generation of mass spectra. Ions were then selected manually for MS/MS using higher energy collisional dissociation (HCD, NCE:20%) in the HCD collision cell. Measurement of the resulting fragment ions was performed in the Orbitrap analyser, set at a resolution of 30000 and which was scanned between m/z 50-1500. The resulting MS/MS spectra were annotated manually.

Modification of peptides by cyclopropenone-based probes:

Typically, peptide (25 μ L, 0.3 – 1 mM) in NaP_i buffer (pH 7, 20 mM) was incubated with CPO-probe (required amount of 15 mM stock solution in CH₃CN, 2 equiv.) at 4 °C for indicated time before being analyzed by LC-MS as described above. For peptides that are poorly soluble in NaP_i buffer, CH₃CN (up to 50%) or DMF (up to 10%) were added to the reaction mixture.

Entry	Sequence	Chemical Formula	Molecular Weight	Purchased from
P1	CAIAI-NH ₂	$C_{21}H_{40}N_6O_5S$	488.6	Biomatik
P2	CAIKI-NH ₂	C24H47N7O5S	545.7	Biomatik
P3	CDPGYIGSR (Laminin)	$C_{40}H_{62}N_{12}O_{14}S$	967.07	Kindly provided by Prof. Pedro M. P. Gois
P4	CGAGESGKSTIVKQMK (GTP-Binding Protein Fragment, G alpha)	$C_{66}H_{118}N_{20}O_{23}S_2$	1623.91	APExBIO Technology
Р5	CYFQNCPRG-NH ₂ (Disulfide bridge: 1-6; Vasopressin)	C46H64N14O13S2	1085.22	Abcam

Peptides used in this study:

Reactions and characterization of peptide conjugates



Reaction of peptide P1 with CPO-BN to give P1-BNa and P1-BNb

Figure S15. HPLC and MS analysis of reaction between peptide P1 (15 μ L of 0.5 mM stock solution in in NaP_i buffer (pH 7, 20 mM)/DMF 9/1) and **CPO-BN** (10 μ L of 1.5 mM stock solution in CH₃CN, 2 equiv.). The reaction was complete after incubation for 1 h at 4 °C.

Reaction of peptide P1 with CPO-PEG to give P1-PEGa and P1-PEGb



Figure S16. HPLC and MS analysis of reaction between peptide P1 (15 μ L of 0.5 mM stock solution in in NaP_i buffer (pH 7, 20 mM)/DMF 9/1) and **CPO-PEG** (10 μ L of 1.5 mM stock solution in CH₃CN, 2 equiv.). The reaction was complete after incubation for 1 h at 4 °C.

Reaction of peptide P2 with CPO-BN to give P2-BNa and P2-BNb



Figure S17. HPLC and MS analysis of reaction between peptide P2 (20 μ L of 0.5 mM stock solution in in NaP_i buffer (pH 7, 20 mM)) and **CPO-BN** (2.5 μ L of 8 mM stock solution in CH₃CN, 2 equiv.). The reaction was complete after incubation for 1 h at 4 °C.

Reaction of peptide P2 with CPO-PEG to give P2-PEGa and P2-PEGb



Figure S18. HPLC and MS analysis of reaction between peptide P2 (20 μ L of 0.5 mM stock solution in in NaP_i buffer (pH 7, 20 mM)) and **CPO-PEG** (2.5 μ L of 8 mM stock solution in CH₃CN, 2 equiv.). The reaction was complete after incubation for 1 h at 4 °C.

Reaction of peptide P3 with CPO-BN to give P3-BNa and P3-BNb



Figure S19. HPLC and MS analysis of reaction between peptide P3 (25 μ L of 0.4 mM stock solution in in NaP_i buffer (pH 7, 20 mM)) and **CPO-BN** (2.5 μ L of 8 mM stock solution in CH₃CN, 2 equiv.). The reaction was complete after incubation for 1 h at 4 °C.
Reaction of peptide P4 with CPO-BN to give P4-BNa and P4-BNb



Figure S20. HPLC and MS analysis of reaction between peptide P4 (25 μ L of 0.4 mM stock solution in in NaP_i buffer (pH 7, 20 mM)) and **CPO-BN** (2.5 μ L of 8 mM stock solution in CH₃CN, 2 equiv.). The reaction was complete after incubation for 1 h at 4 °C.



Reduction of peptide P5 with tris(2-carboxyethyl)phosphine (TCEP) to give P5-SH

Figure S21. HPLC and MS analysis of reaction between peptide **P5** (50 μ L of 2 mM stock solution in in NaP_i buffer (pH 7, 20 mM)) and TCEP (1 μ L of 100 mM stock solution in MQ-H₂O, 1 equiv.). The reaction was complete after incubation for 1 h at 25 °C.



Reaction of peptide P5-SH with N-methylmaleimide to give P5-MI

Figure S22. HPLC and MS analysis of reaction between peptide **P5-SH** (25 μ L of 0.5 mM stock solution in in NaP_i buffer (pH 7, 20 mM)) and *N*-methylmaleimide (3.9 μ L of 8 mM stock solution in CH₃CN, 2.5 equiv.). The reaction was complete after incubation for 1 h at 25 °C, both cysteine residues were modified.



Reaction of peptide P5-SH with 2-cyano-6-hydroxybenzothiazole to give P5-CBTa and P5-CBTb

Figure S23. HPLC and MS analysis of reaction between peptide **P5-SH** (30 μ L of 0.3 mM stock solution in in NaP_i buffer (pH 7, 20 mM)) and 2-cyano-6-hydroxybenzothiazole (1.5 μ L of 12 mM stock solution in CH₃CN, 2.0 equiv.). The reaction was analyzed after incubation for 1 h at 4 °C, peptide conjugates with one modification (**P5-CBTa**) and two modifications (**P5-CBTb**) were observed.





Figure S24. HPLC and MS analysis of reaction between peptide **P5-SH** (30 μ L of 0.3 mM stock solution in in NaP_i buffer (pH 7, 20 mM)) with **CPO-BN** (2.25 μ L of 8 mM stock solution in CH₃CN, 2.0 equiv.). The reaction was complete after incubation for 1 h at 4 °C with only one cysteine residue being modified.

Reaction of peptide conjugates P5-BNa and P5-BNb with N-methylmaleimide to give P5-BNa-MI and P5-BNb-MI



Figure S25. HPLC and MS analysis of reaction between peptides **P5-BNa** and **P5-BNb** (15 μ L of the reaction mixture prepared as described above) with *N*-methylmalemide (1.05 μ L of 8 mM stock solution in CH₃CN, 2 equiv.). The reaction was complete after incubation for 1 h at 4 °C.

LC-MS/MS analysis of P5-BNa-MI and P5-BNb-MI confirming the modified sites



Figure S26. MS/MS spectrum of the m/z 713.8 doubly charged ion of the peptide CYFQNCPRG containing a cyclopropenone modification at the N-terminal cysteine residue and an N-methylmaleimide modification at the internal cysteine.

10. Protein modification

General procedures and characterization methods

LC–MS method for analysis of protein conjugation. LC–MS was performed on a Waters SQ Detector 2 mass spectrometer coupled to an Acquity UPLC system using an Acquity UPLC Protein BEH C4 column (300Å, 1.7 μ m, 2.1 mm × 50 mm). Solvents A, water with 0.1% formic acid and B, 71% acetonitrile, 29% water and 0.075% formic acid were used as the mobile phase at a flow rate of 0.2 mL min⁻¹. The gradient was programmed as follows: 28% B to 71.2% B in 12 min, then 71.2% B for 1 min, 100% B for 2 min, and after that 28% A for 4 min. The electrospray source was operated with a capillary voltage of 3.0 kV and a cone voltage of 30 V. Nitrogen was used as the desolvation gas at a total flow of 800 L h⁻¹. Total mass spectra were reconstructed from the ion series using the MaxEnt algorithm preinstalled on MassLynx software (v. 4.1 from Waters) according to the manufacturer's instructions. To obtain the ion series described, the major peak(s) of the chromatogram were selected for integration and further analysis.

Analysis of protein conjugation by LC–MS. A typical analysis of a conjugation reaction by LC–MS is described below. The total ion chromatogram, combined ion series and deconvoluted spectra are shown for the product of the reaction. Identical analyses were carried out for all the conjugation reactions performed in this work.

Before conjugation

After conjugation

Total ion chromatogram for conjugation reaction. Region containing protein marked in black:



Figure S27. A typical analysis of a conjugation reaction by LC–MS is described for the reaction of Cys-GFP protein with cyclopropenone derivative CPO-Bn. The total ion chromatogram, combined ion series and deconvoluted spectra are shown for the starting material and the product of the reaction of Cys-GFP with 100 equiv. of **CPO-BN**. Identical analyses were carried out for all the conjugation reactions performed in this work.

Enzymatic digestion and LC-MS/MS analysis

Protein solutions were subjected to enzymatic digestion in 50 mM ammonium bicarbonate (pH 8) with trypsin overnight at 37 °C. After digestion, the supernatant was pipetted into a sample vial and loaded onto an autosampler for automated LC–MS/MS analysis.

All LC–MS/MS experiments were performed using a Dionex Ultimate 3000 RSLC nanoUPLC (Thermo Fisher Scientific Inc, Waltham, MA, USA) system and a QExactive Orbitrap mass spectrometer (Thermo Fisher Scientific Inc, Waltham, MA, USA). Separation of peptides was performed by reverse-phase chromatography at a flow rate of 300 nL/min and a Thermo Scientific reverse-phase nano Easy-spray column (Thermo Scientific PepMap C18, 2 μ m particle size, 100 Å pore size, 75 μ m i.d. x 50cm length). Peptides were loaded onto a pre-column (Thermo Scientific PepMap 100 C18, 5 μ m particle size, 100 Å pore size, 300 μ m i.d. x 5mm length) from the Ultimate 3000 autosampler with 0.1% formic acid for 3 minutes at a flow rate of 10 μ L/min. After this period, the column valve was switched to allow elution of peptides from the pre-column onto the analytical column. Solvent A was water + 0.1% formic acid and solvent B was 80% acetonitrile, 20% water + 0.1% formic acid. The linear gradient employed was 2–40% B in 30 minutes.

The LC eluant was sprayed into the mass spectrometer by means of an Easy-Spray source (Thermo Fisher Scientific Inc.). All m/z values of eluting ions were measured in an Orbitrap mass analyzer, set at a resolution of 70000 and was scanned between m/z 380-1500. Data dependent scans (Top 20) were employed to automatically isolate and generate fragment ions by higher energy collisional dissociation (HCD, NCE:25%) in the HCD collision cell and measurement of the resulting fragment ions was performed in the Orbitrap analyser, set at a resolution of 17500. Singly charged ions and ions with unassigned charge states were excluded from being selected for MS/MS and a dynamic exclusion window of 20 seconds was employed.

Post-run, the data was processed using Protein Discoverer (version 2.1., Thermo Scientific). Briefly, all MS/MS data were converted to mgf files and the files were then submitted to the Mascot search algorithm (Matrix Science, London UK) and searched against a custom database containing the GFP and a common contaminant sequences (123 sequences; 40594 residues). Variable modifications of oxidation (M), deamidation (NQ) and the **CPO-BN** modification were applied. The peptide and fragment mass tolerances were set to 100 ppm and 0.1 Da, respectively. A significance threshold value of p<0.05 and a peptide cut-off score of 20 were also applied.

Proteins used in this study

Cys-GFP

Enhanced Green fluorescent protein with N-terminal cysteine (Cys-GFP, for the sake of simplicity) was expressed and purified as described below:

Genetic construct

The pBAD-EGFP vector containing the EGFP gene was purchased from Addgene (Catalog #54762). This vector contains a N-terminal 6x His-T7-TEV-EGFP gene construct. To introduce a cysteine at the N-terminal of the enhanced green fluorescent protein (EGFP), a single point mutation in the glycine residue of the TEV sequence (ENLYFQ/G) was generated by the NZYMutagenesis kit (nzytech) using the mutagenic primers 5'- gaacctgtacttccagtgctcgagcatggtgag-3' and 5'- ctcaccatgctcgagcactggaagtacaggttc-3'. The amplification products were transformed into *E. coli* NZY5 α and the (His-Cys-EGFP) plasmid construct was confirmed by PCR and sequencing at Eurofins Genomics Facility.

Cell growth and protein purification

The vector encoding the His-Cys-EGFP gene was used to transform competent E. coli cells (BL21.A1 strain). For protein expression, cells were inoculated into Luria Bertani (LB) media containing 100 µg/mL of carbenicillin and grown at 37 °C, until an optical density of 0.4–0.6 was reached. Expression of the EGFP protein was induced by adding 1 mM isopropyl β -D-1-thiogalactopyranoside (ITPG) and 0.2% of L-arabinose, and the culture is left to incubate with shaking at 18 °C for 4-6 hours. Cells were harvested by centrifugation and the pellet was resuspended in 1:3 of cold lysis buffer containing 50 mM Tris-HCl pH 7.5, 150 mM NaCl, 20 mM imidazole, the protease inhibitor cocktail (cOmplete^{TMGFP}-Roche), and 1 mM DTT. The cells were lysed by sonication for 4 min in total (20s on, 20s off/cycle) and the lysate was clarified by centrifugation at 18000 rpm, for 30 min. The supernatant was then loaded into a HisTrapTM HP affinity column (GE Healthcare Life Sciences), pre-equilibrated with 20 mM Tris-HCl pH 7.5, 150 mM NaCl, 20 mM imidazole, and 1 mM DTT. The His-Cys-GFP protein was eluted with 20 mM Tris-HCl pH 7.5, 150 mM NaCl, 20 mM imidazole, 1 mM DTT and 500 mM of imidazole. Fractions enriched with His-Cys-GFP, were pooled and treated with TEV protease (in house) to remove the 6x His-tag. The treated sample was loaded into a HisTrapTM column, and the flowthrough (containing the TEV treated Cys-GFP), was pooled and concentrated to 4 mg/mL. Samples containing the Cys-GFP protein were analyzed by SDS-PAGE (Figure S28) and LC-MS (Figure S29).



Figure S28. 15% SDS-PAGE of Cys-GFP protein.

Before modification reactions, a batch of Cys-GFP (100 μ L, 30 μ M) was treated with TCEP (1 μ L, 30 mM, 10 equiv.) for 1 h at 25 °C. Small molecules were removed Amicon Ultra-0.5 mL Centrifugal Filter (MWCO 3 KDa) to afford the Cys-GFP (80 μ L, 33 μ M) with a reactive cysteine in NaP_i buffer (pH 7, 20 mM). For storage, DTT (0.88 μ L, 1.5 M, 500 equiv.) was added to the buffer to prevent cysteine oxidation (final DTT concentration is 16.5 mM).

Amino acid sequence (20 lysines, 3 cysteines, no disulfides):

 $\underline{C} SSMVSKGEELFTGVVPILVELDGDVNGHKFSVSGEGEGDATYGKLTLKFICTTGKLPVPWP TLVTTLTYGVQCFSRYPDHMKQHDFFKSAMPEGYVQERTIFFKDDGNYKTRAEVKFEGDTL VNRIELKGIDFKEDGNILGHKLEYNYNSHNVYIMADKQKNGIKVNFKIRHNIEDGSVQLADH YQQNTPIGDGPVLLPDNHYLSTQSALSKDPNEKRDHMVLLEFVTAAGITLGMDELYK$



Isotopically Averaged Molecular Weight = 27199 Da

Figure S29. Combined ion series and deconvoluted mass spectra of Cys-GFP. Calcd. mass, 27199 Da, found: 27196 Da.

C2Am

Engineered variant of the C2A domain of Synaptotagmin-I Cys 95 mutant (C2Am) was kindly provided by Dr. André Neves.¹⁰

Amino acid sequence (14 lysines, 1 free cysteine, no disulfides):

 $GSPGISGGGGGILDSMVEKLGKLQYSLDYDFQNNQLLVGIIQAAELPALDMGGTSDPYVKVF\\ LLPDKKKKFETKVHRKTLNPVFNEQFTFKVPY\underline{C}ELGGKTLVMAVYDFDRFSKHDIIGEFKVP\\ MNTVDFGHVTEEWRDLQSAEK$

Isotopically Averaged Molecular Weight = 16222 Da



Figure S30. Combined ion series and deconvoluted mass spectra of C2Am. Calcd. mass, 16222 Da, found: 16224 Da.

Annexin V

Annexin V was expressed and purified as previously reported.¹¹

Amino acid sequence (22 lysines, 1 free cysteine, no disulfides):

AQVLRGTVTDFPGFDERADAETLRKAMKGLGTDEESILTLLTSRSNAQRQEISAAFKTLFGRD LLDDLKSELTGKFEKLIVALMKPSRLYDAYELKHALKGAGTNEKVLTEIIASRTPEELRAIKQ VYEEEYGSSLEDDVVGDTSGYYQRMLVVLLQANRDPDAGIDEAQVEQDAQALFQAGELKW GTDEEKFITIFGTRSVSHLRKVFDKYMTISGFQIEETIDRETSGNLEQLLLAVVKSIRSIPAYLAE TLYYAMKGAGTDDHTLIRVMVSRSEIDLFNIRKEFRKNFATSLYSMIKGDTSGDYKKALLLL CGEDD

Isotopically Averaged Molecular Weight = 35805 Da



Figure S31. Combined ion series and deconvoluted mass spectra of Annexin V. Calcd. mass, 35805 Da, found: 35814 Da.

Nanobody DesAB-HET¹²

Nanobody Cys57 mutant was kindly provided by Philip Lindstedt and Prof. Michele Vendruscolo. Amino acid sequence (7 lysines, 1 free cysteine, 1 disulfide):

MRGSHHHHHHGMASMTGGQQMGRDLYDDDDKDPKLEVQLVESGGGLVQPGGSLRLSCAA SGFNIKDTYIGWVRRAPGKGKEWVASIYPTNGYTRYADSVKGRFTISADTSKNTAYLQMNSL RAEDTAVYYCAAGSCETLTLREEEAAAWGQGTLVTVSSGT



Isotopically Averaged Molecular Weight = 17391 Da

Figure S32. Combined ion series and deconvoluted mass spectra of DesAB-HET nanobody. Calcd. mass, 17391 Da, found: 17394 Da.

Reactions and characterization of Cys-GFP-conjugates

Reaction of Cys-GFP with CPO-BN to give GFP-CPO-BN



A mixture of 8 μ L of Cys-GFP stock solution (30 μ M in 20 mM pH 7 NaP_i buffer containing 16.5 mM DTT), 12 μ L of 20 mM pH 7 NaP_i buffer, and 2.5 μ L of cyclopropenone reagent **CPO-BN** (16 mM in CH₃CN, 100 equiv.) was shaken in an 1.5 mL microcentrifuge tube at 25 °C for 2 h. The modified product was characterized by LC-MS analysis and complete conversion to the expected product was observed (calculated mass, 27428 Da; observed mass, 27434 Da).



Figure S33. The total ion chromatogram, combined ion series and deconvoluted mass spectrum of the reaction between Cys-GFP and CPO-BN after 2 h at 25 °C.

Stability of GFP-CPO-BN in glutathione (GSH).

A 30 μ L aliquot of GFP-CPO-BN (5 μ M) in 20 mM NaP_i buffer at pH 7.0 was thawed. Then, a freshly prepared solution of L-Glutathione reduced form (3 μ L, 55 mM stock in MQ-H₂O) was added at room temperature and the resulting mixture vortexed for 10 seconds (final GSH concentration is 5 mM). The resulting reaction mixture was then shaken at 37 °C. After 1 and 24 h, an aliquot of the reaction mixture was analyzed by LC–MS.



Figure S34. LC–MS trace, combined ion series and deconvoluted mass spectrum of GFP-CPO-BN in GSH. Incubation of the bioconjugate GFP-CPO-BN with 5 mM with GSH shows no degradation after 1 h at 37 °C.



Figure S35. LC–MS trace, combined ion series and deconvoluted mass spectrum of GFP-CPO-BN in GSH. Incubation of the bioconjugate GFP-CPO-BN with 5 mM with GSH shows no degradation after 24 h at 37 °C.

Enzymatic digestion of GFP-CPO-BN and LC-MS/MS analysis confirming the modified site



Figure S36. MS/MS spectrum of the m/z 485.72 doubly charged ion of the tryptic peptide <u>CSSMVSKL</u> from GFP containing a modification at the N-terminal cysteine residue. The underscore relates to the modified amino acid.

Reaction of Cys-GFP with CPO-PEG



A mixture of 8 μ L of Cys-GFP stock solution (30 μ M in 20 mM pH 7 NaP_i buffer containing 16.5 mM DTT), 12 μ L of 20 mM pH 7 NaP_i buffer, and 2.5 μ L of cyclopropenone reagent **CPO-PEG** (16 mM in CH₃CN, 100 equiv.) was shaken in an 1.5 mL microcentrifuge tube at 25 °C for 2 h. The modified product was characterized by LC-MS analysis and complete conversion to the expected product was observed (calculated mass, 27514 Da; observed mass, 27516 Da).



Figure S37. The total ion chromatogram, combined ion series and deconvoluted mass spectrum of the reaction between Cys-GFP and CPO-PEG after 2 h at 25 °C.

Reaction of Cys-GFP with CPO-EDANS



A mixture of 8 μ L of Cys-GFP stock solution (30 μ M in 20 mM pH 7 NaP_i buffer containing 16.5 mM DTT), 12 μ L of 20 mM pH 7 NaP_i buffer, and 2.5 μ L of cyclopropenone reagent **CPO-EDANS** (16 mM in DMF, 100 equiv.) was shaken in an 1.5 mL microcentrifuge tube at 25 °C for 2 h. The modified product was characterized by LC-MS analysis and 70% conversion to the expected product was observed (calculated mass for [M-Na⁺], 27586 Da; observed mass, 27589 Da).



Figure S38. The total ion chromatogram, combined ion series and deconvoluted mass spectrum of the reaction between Cys-GFP and CPO-EDANS after 2 h at 25 °C.

Reaction of Cys-GFP with N-methylmaleimide



A mixture of 8 μ L of Cys-GFP stock solution (30 μ M in 20 mM pH 7 NaP_i buffer containing 16.5 mM DTT), 12 μ L of 20 mM pH 7 NaP_i buffer, and 2.5 μ L of *N*-methylmaleimide (16 mM in CH₃CN, 100 equiv.) was shaken in an 1.5 mL microcentrifuge tube at 25 °C for 2 h. The modified product was characterized by LC-MS analysis and no conversion to the product was observed (calculated mass, 27310 Da; observed mass, 27206 Da (starting protein only)).



Figure S39. The total ion chromatogram, combined ion series and deconvoluted mass spectrum of the reaction between Cys-GFP and *N*-methylmaleimide after 2 h at 25 °C. No reaction was observed.

Reaction of Cys-GFP with CPO-Biotin



A mixture of 3 μ L of Cys-GFP stock solution (25 μ M in 20 mM pH 7 NaP_i buffer), 10.65 μ L of 20 mM pH 7 NaP_i buffer, 0.6 μ L of DTT stock solution (62.5 mM in MQ-H₂O, 500 equiv.), and 0.75 μ L of cyclopropenone reagent **CPO-Biotin** (10 mM in CH₃CN, 100 equiv., prepared as described on page S69) was shaken in an 1.5 mL microcentrifuge tube at 25 °C for 2 h. The modified product was characterized by LC-MS analysis and complete conversion to the product was observed (calculated mass, 27695 Da; observed mass, 27702 Da).



Figure S40. The total ion chromatogram, combined ion series and deconvoluted mass spectrum of the reaction between Cys-GFP and CPO-Biotin after 2 h at 25 °C.

Reaction of Cys-GFP with Maleimide-Biotin



A mixture of $3 \mu L$ of Cys-GFP stock solution (25 μ M in 20 mM pH 7 NaP_i buffer), 10.65 μ L of 20 mM pH 7 NaP_i buffer, 0.6 μ L of DTT stock solution (62.5 mM in MQ-H₂O, 500 equiv.), and 0.75 μ L of cyclopropenone reagent **MI-Biotin** (10 mM in CH₃CN, 100 equiv., prepared as described on page S70) was shaken in an 1.5 mL microcentrifuge tube at 25 °C for 2 h. The modified product was characterized by LC-MS analysis and 13% conversion to the product was observed (calculated mass, 27766 Da; observed mass, 27771 Da).



Figure S41. The total ion chromatogram, combined ion series and deconvoluted mass spectrum of the reaction between Cys-GFP and **MI-Biotin** after 2 h at 25 °C.

Selective modification of GFP with N-terminal cysteine in a mixture of proteins:

Preparation of CPO-Biotin probe:



To a glass vial containing 28 μ L of CH₃CN, CPO-probe **6** (2.4 mg, 7.84 μ mol, 1 equiv.), DIPEA (1.3 μ l 7.5 μ mol, 1 equiv.) and N-biotinyl-3,6-dioxaoctane-1,8-diamine trifluoroacetate salt solution (167 μ L of 25 mg/mL reagent in DMSO, 8.55 μ mol, 1.1 equiv.) were added, and the reaction mixture was incubated for 1 h at 25 °C. Next, 1 μ L aliquot of the reaction mixture was mixed with 39 μ L of CH₃CN, the resulting sample was analyzed by LC–MS, and complete conversion of CPO-probe **6** to CPO-biotin probe was observed (**Figure S42**). The resulting CPO-biotin solution was diluted to 5 mM with CH₃CN (assuming quantitative conversion) and used for protein selectivity experiment without further purification.



Figure S42. HPLC and MS analysis of reactions between CPO-probe **6** and N-biotinyl-3,6-dioxaoctane-1,8-diamine trifluoroacetate salt solution. The reaction was complete after incubation for 1 h at 25 °C.



To a glass vial containing 32 μ L of CH₃CN, perfluorophenyl 6-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1yl)hexanoate (3.3 mg, 8.75 μ mol, 1 equiv.), DIPEA (1.5 μ L, 8.61 μ mol, 1 equiv.) and N-biotinyl-3,6dioxaoctane-1,8-diamine trifluoroacetate salt solution (188 μ L of 25 mg/mL reagent in DMSO, 9.62 μ mol, 1.1 equiv.) were added, and the reaction mixture was incubated for 1 h at 25 °C. Next, 1 μ L aliquot of the reaction mixture was mixed with 39 μ L of CH₃CN, the resulting sample was analyzed by LC–MS, and complete conversion of perfluorophenyl ester to maleimide-biotin (**MI-biotin**) probe was observed (**Figure S43**). The resulting maleimide-biotin solution was diluted to 5 mM with CH₃CN (assuming quantitative conversion) and used for protein selectivity experiment without further purification.



Figure S43. HPLC and MS analysis of reactions between perfluorophenyl 6-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)hexanoate and N-biotinyl-3,6-dioxaoctane-1,8-diamine trifluoroacetate salt solution. The reaction was complete after incubation for 1 h at 25 °C.

Preparation of protein stocks:

To an eppendorf containing 100 μ L of 30 μ M Cys-GFP protein stock solution in Tris HCl buffer (20 mM, pH 7.5), 10 equiv. of TCEP (1 μ L of 30 mM stock in MQ H₂O) was added, and the reaction mixed for 2 h at 25 °C. The resulting sample was buffer-exchanged using an Amicon[®] Ultra 3K 0.5 mL centrifugal filter device (*Merck*) by centrifugation (3 x 15 min) using NaP_i buffer (pH 7, 20 mM). Next, protein concentration was assessed by measuring absorbance at 280 nm and was determined to be 28 μ M. The freshly reduced and buffer-exchanged protein stock solution was then used for protein selectivity experiment.

C2Am, Annexin V and Nanobody DesAB-HET protein stocks were buffer-exchanged using an Amicon[®] Ultra 3K 0.5 mL centrifugal filter device (*Merck*) by centrifugation (3 x 15 min) using NaP_i buffer (pH 7, 20 mM). Next, protein concentration was assessed by measuring absorbance at 280 nm and was determined to be 55 μ M (C2Am), 49 μ M (Annexin V) and 48 μ M (DesAB-HET). The protein stock solutions were then used for protein selectivity experiment.

Protein modification with CPO-Biotin or maleimide-biotin probes:

To an eppendorf containing 13 μ L of NaP_i buffer (20 mM, pH 7.0), 4.5 μ L of a stock solution of Cys-GFP (28 μ M, final concentration 5 μ M), 2.6 μ L of a stock solution of Annexin V (49 μ M, final concentration 5 μ M), 2.6 μ L of a stock solution of DesAB-HET (48 μ M, final concentration 5 μ M), 2.3 μ L of a stock solution of C2Am (55 μ M, final concentration 5 μ M) were added and the resulting mixture was vortexed for 5 seconds. Afterwards, 50 equiv. (relative to Cys-GFP) of CPO-biotin or maleimide-biotine probe (1.25 μ l of 5 mM stock solution) was added to the solution, and the reaction mixed for 1 h at 25 °C. In a control experiment, 1.25 μ l of acetonitrile was added instead of the probes. Then, a 10 μ L aliquot was analyzed by LC–MS and the conversion for each protein was analyzed (**Figures S45–46**).

Protein modification with CPO-Biotin or maleimide-biotin probes in the presence of DTT:

To an eppendorf containing 12 μ L of NaP_i buffer (20 mM, pH 7.0), 4.5 μ L of a stock solution of Cys-GFP (28 μ M, final concentration 5 μ M), 2.6 μ L of a stock solution of Annexin V (49 μ M, final concentration 5 μ M), 2.6 μ L of a stock solution of DesAB-HET (48 μ M, final concentration 5 μ M), 2.3 μ L of a stock solution of C2Am (55 μ M, final concentration 5 μ M) were added and the resulting mixture was vortexed for 5 seconds. Afterwards, 500 equiv. (relative to Cys-GFP) of DTT (1 μ l of 62.5 mM stock solution in MQ H₂O) and 50 equiv. (relative to Cys-GFP) of CPO-biotin or maleimide-biotine probe (1.25 μ l of 5 mM stock solution) were added to the solution, and the reaction mixed for 1 h at 25 °C. In a control experiment, 1.25 μ l of acetonitrile was added instead of the probes. Then, a 10 μ L aliquot was analyzed by LC–MS and the conversion for each protein was analyzed (**Figures S47–S48**).

Table S3. Calculates masses for Cys-GFP, C2Am, AnxV and DesAB-HET proteins, and their conjugates with CPO-biotin or MI-biotin.

	Calculates mass		
Protein	Unmodified	Conjugate with CPO-biotin	Conjugate with MI-biotin
DesAB-HET	17391 Da	17887 Da	17958 Da
Cys-GFP	27199 Da	27695 Da	27766 Da
C2Am	16222 Da	16718 Da	16789 Da
AnxV	35805 Da	36301 Da	36372 Da



Figure S44. LC–MS trace, combined ion series and deconvoluted mass spectrum of the DesAB-HET 1, Cys-GFP 2, C2Am 3 and AnxV 4 protein mixture after incubation for 1 h at 25 °C.



Figure S45. LC–MS trace, combined ion series and deconvoluted mass spectrum of the DesAB-HET 1, Cys-GFP 2, C2Am 3 and AnxV 4 protein mixture after incubation with 50 equiv. of CPO-Biotin probe for 1 h at 25 °C. Cys-GFP protein is modified (>90%) while the other proteins remain unchanged.



Figure S46. LC–MS trace, combined ion series and deconvoluted mass spectrum of the DesAB-HET **1**, Cys-GFP **2**, C2Am **3** and AnxV **4** protein mixture after incubation with 50 equiv. of Maleimide-Biotin probe for 1 h at 25 °C. As expected, full conjugation is observed for DesAB-HET, Cys-GFP, and C2Am proteins, while AnxV protein remains unchanged.



Figure S47. LC–MS trace, combined ion series and deconvoluted mass spectrum of the DesAB-HET 1, Cys-GFP 2, C2Am 3 and AnxV 4 protein mixture after incubation with 50 equiv. of CPO-Biotin probe in the presence of 2.5 mM DTT for 1 h at 25 °C. Cys-GFP protein is modified (>95%) while the other proteins remain unchanged.



Figure S48. LC–MS trace, combined ion series and deconvoluted mass spectrum of the DesAB-HET 1, Cys-GFP 2, C2Am 3 and AnxV 4 protein mixture after incubation with 50 equiv. of Maleimide-Biotin probe in the presence of 2.5 mM DTT for 1 h at 25 °C. DesAB-HET and AnxV remain unreacted, while for Cys-GFP and C2Am 10–15% modification is observed.

11. Molecular dynamics (MD) simulations

The crystal structure of GFP (PDB id: 1GFL)¹³ was used as starting coordinates for the proteins in the simulations. The missing amino acids in the peptide sequence were added with PyMOL 2.0 (https://pymol.org/2/) and modeled with Chimera.¹⁴ MD simulations were carried out with AMBER 18 package,¹⁵ implemented with ff14SB,¹⁶ and GAFF¹⁷ force fields. Parameters for the unnatural residues were generated with the antechamber module of AMBER, using GAFF force field and with partial charges set to fit the electrostatic potential generated with HF/6-31G(d) by RESP.¹⁸ The charges were calculated according to the Merz-Singh-Kollman scheme using Gaussian 16.³ Each conjugate was immersed in a water box with a 10 Å buffer of TIP3P water molecules¹⁹ and neutralized by adding explicit Na⁺ ions. A two-stage geometry optimization approach was performed. The first stage minimizes only the positions of solvent molecules and ions, and the second stage is an unrestrained minimization of all the atoms in the simulation cell. The systems were then heated by incrementing the temperature from 0 to 300 K under a constant pressure of 1 atm and periodic boundary conditions. Harmonic restraints of 10 kcal·mol⁻¹ were applied to the solute, and the Andersen temperature coupling scheme²⁰ was used to control and equalize the temperature. The time step was kept at 1 fs during the heating stages, allowing potential inhomogeneities to self-adjust. Hydrogen atoms were kept fixed through the simulations using the SHAKE algorithm.²¹ Long-range electrostatic effects were modelled using the particle-mesh-Ewald method.²² An 8 Å cutoff was applied to Lennard-Jones interactions. Each system was equilibrated for 2 ns with a 2-fs time step at a constant volume and temperature of 300 K. Production trajectories were then run for additional 0.5 µs under the same simulation conditions (Figure S49).



Figure S49. Structural ensembles derived from 0.5 μ MD simulations performed on the four stereoisomers of **GFP-CPO-BN**. The numbers indicate the root-mean-square deviation (RMSD) for heavy-atom superimposition of the backbone of GFP relative to the average structure through the entire trajectory. The low RMSD values indicate that the structure of the proteins remains unaltered upon the chemical modifications, which is essential to retain their biological activity.

12. NMR spectra

¹H NMR (400 MHz, CDCl₃) of 2-phenethylcycloprop-2-en-1-one 1



¹³C NMR (CDCl₃, 100 MHz) of 2-phenethylcycloprop-2-en-1-one 1





¹H NMR (400 MHz, CDCl₃) of ethyl (3R,6R)-5-oxo-6-phenethyl-1,4-thiazepane-3-carboxylate 2a

¹³C NMR (CDCl₃, 100 MHz) of ethyl (3R,6R)-5-oxo-6-phenethyl-1,4-thiazepane-3-carboxylate 2a



¹H NMR (400 MHz, CDCl₃) of ethyl (3R,6S)-5-oxo-6-phenethyl-1,4-thiazepane-3-carboxylate **2b**, ethyl (3R,7R)-5-oxo-7-phenethyl-1,4-thiazepane-3-carboxylate **2c**, ethyl (3R,7S)-5-oxo-7-phenethyl-1,4-thiazepane-3-carboxylate **2d**



¹³C NMR (CDCl₃, 100 MHz) of ethyl (3R,6S)-5-oxo-6-phenethyl-1,4-thiazepane-3-carboxylate **2b**, ethyl (3R,7R)-5-oxo-7-phenethyl-1,4-thiazepane-3-carboxylate **2c**, ethyl (3R,7S)-5-oxo-7-phenethyl-1,4-thiazepane-3-carboxylate **2d**





¹H NMR (400 MHz, CDCl₃) of 6-phenethyl-1,4-thiazepan-5-one **3a,b** (both enantiomers)

¹³C NMR (CDCl₃, 100 MHz) of 6-phenethyl-1,4-thiazepan-5-one **3a,b** (both enantiomers)




¹H NMR (400 MHz, CDCl₃) of 7-phenethyl-1,4-thiazepan-5-one **3c,d** (both enantiomers)

¹³C NMR (CDCl₃, 100 MHz) of 7-phenethyl-1,4-thiazepan-5-one **3c,d** (both enantiomers)





¹H NMR (400 MHz, CDCl₃) of perfluorophenyl hex-5-ynoate **5**

¹³C NMR (100 MHz, CDCl₃) of perfluorophenyl hex-5-ynoate 5



¹⁹F NMR (376.5 MHz, CDCl₃) of perfluorophenyl hex-5-ynoate 5



¹H NMR (500 MHz, CDCl₃) of perfluorophenyl 4-(3'-oxocycloprop-1'-en-1'-yl)butanoate 6





¹³C NMR (125 MHz, CDCl₃) of Perfluorophenyl 4-(3'-oxocycloprop-1'-en-1'-yl)butanoate 6

¹⁹F NMR (376.5 MHz, CDCl₃) of Perfluorophenyl 4-(3'-oxocycloprop-1'-en-1'-yl)butanoate 6



¹H NMR (400 MHz, CDCl₃) of 4-(3'-oxocycloprop-1'-en-1'-yl)-N-phenylbutanamide CPO-BN



¹³C NMR (125 MHz, CDCl₃) of 4-(3'-oxocycloprop-1'-en-1'-yl)-N-phenylbutanamide CPO-BN



¹H NMR (400 MHz, CDCl₃) of N-(2-(2-(2-(2-hydroxyethoxy)ethoxy)ethoxy)ethyl)-4-(3'-oxocycloprop-1'-en-1'-yl)butanamide **CPO-PEG**



¹³C NMR (125 MHz, CDCl₃) of N-(2-(2-(2-(2-hydroxyethoxy)ethoxy)ethoxy)ethyl)-4-(3'-oxocycloprop-1'-en-1'-yl)butanamide **CPO-PEG**



¹³H NMR (400 MHz, CDCl₃) of 4-(3'-oxocycloprop-1'-en-1'-yl)-N-(3'',6'',9'',12''tetraoxapentadec-14''-yn-1''-yl)butanamide **CPO-PEG-Alkyne**



¹³C NMR (100 MHz, CDCl₃) of 4-(3'-oxocycloprop-1'-en-1'-yl)-N-(3'',6'',9'',12''-tetraoxapentadec-14''-yn-1''-yl)butanamide **CPO-PEG-Alkyne**



¹³H NMR (500 MHz, DMF-*d*₇) of Sodium 5-((2'-(4''-(3'''-oxocycloprop-1'''-en-1'''-yl)butanamido)ethyl)amino) naphthalene-1-sulfonate **CPO-PEG-EDANS**



¹³C NMR (126 MHz, CDCl₃) of of Sodium 5-((2'-(4''-(3'''-oxocycloprop-1'''-en-1'''-yl)butanamido)ethyl)amino) naphthalene-1-sulfonate **CPO-PEG-EDANS**



13.References

1. Sheldrick, G., SHELXT - Integrated space-group and crystal-structure determination. *Acta Crystallogr. A* 2015, *71* (1), 3–8.

2. Sheldrick, G., Crystal structure refinement with SHELXL. Acta Crystallogr. C 2015, 71 (1), 3-8.

3. Frisch, M. J.; Trucks, G. W.; Schlegel, H. B.; Scuseria, G. E.; Robb, M. A.; Cheeseman, J. R.; Scalmani, G.; Barone, V.; Petersson, G. A.; Nakatsuji, H.; Li, X.; Caricato, M.; Marenich, A. V.; Bloino, J.; Janesko, B. G.; Gomperts, R.; Mennucci, B.; Hratchian, H. P.; Ortiz, J. V.; Izmaylov, A. F.; Sonnenberg, J. L.; Williams; Ding, F.; Lipparini, F.; Egidi, F.; Goings, J.; Peng, B.; Petrone, A.; Henderson, T.; Ranasinghe, D.; Zakrzewski, V. G.; Gao, J.; Rega, N.; Zheng, G.; Liang, W.; Hada, M.; Ehara, M.; Toyota, K.; Fukuda, R.; Hasegawa, J.; Ishida, M.; Nakajima, T.; Honda, Y.; Kitao, O.; Nakai, H.; Vreven, T.; Throssell, K.; Montgomery Jr., J. A.; Peralta, J. E.; Ogliaro, F.; Bearpark, M. J.; Heyd, J. J.; Brothers, E. N.; Kudin, K. N.; Staroverov, V. N.; Keith, T. A.; Kobayashi, R.; Normand, J.; Raghavachari, K.; Rendell, A. P.; Burant, J. C.; Iyengar, S. S.; Tomasi, J.; Cossi, M.; Millam, J. M.; Klene, M.; Adamo, C.; Cammi, R.; Ochterski, J. W.; Martin, R. L.; Morokuma, K.; Farkas, O.; Foresman, J. B.; Fox, D. J. *Gaussian 16 Rev. B.01*, Wallingford, CT, 2016.

4. Zhao, Y.; Truhlar, D. G., The M06 suite of density functionals for main group thermochemistry, thermochemical kinetics, noncovalent interactions, excited states, and transition elements: two new functionals and systematic testing of four M06-class functionals and 12 other functionals. *Theor. Chem. Acc.* **2008**, *120* (1), 215–241.

Scalmani, G.; Frisch, M. J., Continuous surface charge polarizable continuum models of solvation.
I. General formalism. J. Chem. Phys. 2010, 132 (11), 114110.

6. Ribeiro, R. F.; Marenich, A. V.; Cramer, C. J.; Truhlar, D. G., Use of Solution-Phase Vibrational Frequencies in Continuum Models for the Free Energy of Solvation. *J. Phys. Chem. B.* **2011**, *115* (49), 14556–14562.

7. Gonzalez, C.; Schlegel, H. B., An improved algorithm for reaction path following. *J. Chem. Phys.* **1989**, *90* (4), 2154–2161.

8. Gonzalez, C.; Schlegel, H. B., Reaction path following in mass-weighted internal coordinates. *J. Phys. Chem.* **1990**, *94* (14), 5523–5527.

9. Ren, H.; Xiao, F.; Zhan, K.; Kim, Y.-P.; Xie, H.; Xia, Z.; Rao, J., A Biocompatible Condensation Reaction for the Labeling of Terminal Cysteine Residues on Proteins. *Angew. Chem. Int. Ed.* **2009**, *48* (51), 9658–9662.

 Alam, I. S.; Neves, A. A.; Witney, T. H.; Boren, J.; Brindle, K. M., Comparison of the C2A Domain of Synaptotagmin-I and Annexin-V As Probes for Detecting Cell Death. *Bioconjug. Chem.* 2010, *21* (5), 884--891.

Freedy, A. M.; Matos, M. J.; Boutureira, O.; Corzana, F.; Guerreiro, A.; Akkapeddi, P.; Somovilla,
V. J.; Rodrigues, T.; Nicholls, K.; Xie, B.; Jiménez-Osés, G.; Brindle, K. M.; Neves, A. A.; Bernardes,

G. J. L., Chemoselective Installation of Amine Bonds on Proteins through Aza-Michael Ligation. J. Am. Chem. Soc. 2017, 139 (50), 18365–18375.

12. Aprile, F. A.; Sormanni, P.; Perni, M.; Arosio, P.; Linse, S.; Knowles, T. P. J.; Dobson, C. M.; Vendruscolo, M., Selective targeting of primary and secondary nucleation pathways in A β 42 aggregation using a rational antibody scanning method. *Sci. Adv.* **2017**, *3* (6), e1700488.

13. Yang, F.; Moss, L. G.; Phillips, G. N., The molecular structure of green fluorescent protein. *Nat. Biotechnol.* **1996**, *14* (10), 1246–1251.

14. Pettersen, E. F.; Goddard, T. D.; Huang, C. C.; Couch, G. S.; Greenblatt, D. M.; Meng, E. C.; Ferrin, T. E., UCSF Chimera—A visualization system for exploratory research and analysis. *J. Comput. Chem.*2004, *25* (13), 1605–1612.

15. D.A. Case, I. Y. B.-S., S.R. Brozell, D.S. Cerutti, T.E. Cheatham, III, V.W.D. Cruzeiro, T.A. Darden, R.E. Duke, D. Ghoreishi, M.K. Gilson, H. Gohlke, A.W. Goetz, D. Greene, R Harris, N. Homeyer, S. Izadi, A. Kovalenko, T. Kurtzman, T.S. Lee, S. LeGrand, P. Li, C. Lin, J. Liu, T. Luchko, R. Luo, D.J. Mermelstein, K.M. Merz, Y. Miao, G. Monard, C. Nguyen, H. Nguyen, I. Omelyan, A. Onufriev, F. Pan, R. Qi, D.R. Roe, A. Roitberg, C. Sagui, S. Schott-Verdugo, J. Shen, C.L. Simmerling, J. Smith, R. Salomon-Ferrer, J. Swails, R.C. Walker, J. Wang, H. Wei, R.M. Wolf, X. Wu, L. Xiao, D.M. York and P.A. Kollman, AMBER 2018, University of California, San Francisco. **2018**.

16. Maier, J. A.; Martinez, C.; Kasavajhala, K.; Wickstrom, L.; Hauser, K. E.; Simmerling, C., ff14SB: Improving the Accuracy of Protein Side Chain and Backbone Parameters from ff99SB. *J. Chem. Theory Comput.* **2015**, *11* (8), 3696–3713.

17. Wang, J.; Wolf, R. M.; Caldwell, J. W.; Kollman, P. A.; Case, D. A., Development and testing of a general amber force field. *J. Comput. Chem.* **2004**, *25* (9), 1157–1174.

18. Bayly, C. I.; Cieplak, P.; Cornell, W.; Kollman, P. A., A well-behaved electrostatic potential based method using charge restraints for deriving atomic charges: the RESP model. *J. Phys. Chem.* **1993**, *97* (40), 10269–10280.

19. Jorgensen, W. L.; Chandrasekhar, J.; Madura, J. D.; Impey, R. W.; Klein, M. L., Comparison of simple potential functions for simulating liquid water. *J. Chem. Phys.* **1983**, *79* (2), 926–935.

20. Andersen, H. C., Molecular dynamics simulations at constant pressure and/or temperature. *J. Chem. Phys.* **1980**, *72* (4), 2384–2393.

21. Miyamoto, S.; Kollman, P. A., Settle: An analytical version of the SHAKE and RATTLE algorithm for rigid water models. *J. Comput. Chem.* **1992**, *13* (8), 952–962.

22. Darden, T.; York, D.; Pedersen, L., Particle mesh Ewald: An N·log(N) method for Ewald sums in large systems. *J. Chem. Phys.* **1993**, *98* (12), 10089–10092.