

Dual Electrocatalysis Enables Enantioselective Hydrocyanation of Conjugated Alkenes

Lu Song, Niankai Fu, Brian G. Ernst, Wai-Hang Lee, Michael O. Frederick, Robert A. DiStasio Jr., [Song Lin](#)

Submitted date: 09/09/2019 • Posted date: 10/09/2019

Licence: CC BY-NC-ND 4.0

Citation information: Song, Lu; Fu, Niankai; Ernst, Brian G.; Lee, Wai-Hang; Frederick, Michael O.; DiStasio Jr., Robert A.; et al. (2019): Dual Electrocatalysis Enables Enantioselective Hydrocyanation of Conjugated Alkenes. ChemRxiv. Preprint.

Chiral nitriles and their derivatives are prevalent in pharmaceuticals and bioactive compounds. Enantioselective alkene hydrocyanation represents a convenient and efficient approach for synthesizing these molecules. However, a generally applicable method featuring a broad substrate scope and high functional group tolerance remains elusive. Here, we address this long-standing synthetic problem using an electrocatalytic strategy. Electrochemistry allows for the seamless combination of two classic radical reactions—cobalt-mediated hydrogen-atom transfer and copper-promoted radical cyanation—to accomplish highly enantioselective hydrocyanation without the need for stoichiometric oxidant. We harness electrochemistry's unique feature of precise potential control to optimize the chemoselectivity of challenging substrates. Computational analysis sheds light on the origin of enantioinduction, for which the chiral catalyst imparts a combination of attractive and repulsive non-covalent interactions that direct the enantio-determining C-CN bond formation. This discovery demonstrates the power of electrochemistry in accessing new chemical space and providing solutions to pertinent challenges in synthetic chemistry.

File list (2)

01_Lin_MS.pdf (1.70 MiB)

[view on ChemRxiv](#) • [download file](#)

00_Lin_TOC.png (333.52 KiB)

[view on ChemRxiv](#) • [download file](#)

Dual Electrocatalysis Enables Enantioselective Hydrocyanation of Conjugated Alkenes

Lu Song¹, Niankai Fu¹, Brian G. Ernst¹, Wai Hang Lee¹, Michael O. Frederick²,
Robert A. DiStasio Jr.^{1*}, Song Lin^{1*}

¹Department of Chemistry and Chemical Biology, Cornell University, Ithaca, NY 14853, USA

²Small Molecule Design and Development, Eli Lilly and Company, Indianapolis, IN 46285, USA

*Correspondence to: songlin@cornell.edu, distasio@cornell.edu

Abstract: Chiral nitriles and their derivatives are prevalent in pharmaceuticals and bioactive compounds. Enantioselective alkene hydrocyanation represents a convenient and efficient approach for synthesizing these molecules. However, a generally applicable method featuring a broad substrate scope and high functional group tolerance remains elusive. Here, we address this long-standing synthetic problem using an electrocatalytic strategy. Electrochemistry allows for the seamless combination of two classic radical reactions—cobalt-mediated hydrogen-atom transfer and copper-promoted radical cyanation—to accomplish highly enantioselective hydrocyanation without the need for stoichiometric oxidant. We harness electrochemistry’s unique feature of precise potential control to optimize the chemoselectivity of challenging substrates. Computational analysis sheds light on the origin of enantioinduction, for which the chiral catalyst imparts a combination of attractive and repulsive non-covalent interactions that direct the enantio-determining C–CN bond formation. This discovery demonstrates the power of electrochemistry in accessing new chemical space and providing solutions to pertinent challenges in synthetic chemistry.

Introduction

Alkene hydrocyanation, in which H and CN groups are added across a C=C π -bond, is a highly useful transformation,^{1–4} as it provides versatile nitriles that are key intermediates in the synthesis of polymers, agrochemicals, cosmetics, and pharmaceuticals (Figure 1A).⁵ This reaction is used in the industrial production of adiponitrile on a million-ton annual scale⁶ and has been explored by scientists at DuPont in the synthesis of naproxen, a top-selling anti-inflammatory medicine.^{7–9} The development of a general catalytic alkene hydrocyanation method with broad substrate scope, precise chemoselectivity, and high stereochemical control would have significant impact on synthetic organic chemistry across both academia and industry. Despite extensive studies on the enantioselective hydrocyanation of polar π -bonds (e.g., C=O and C=N),¹⁰ analogous methods for the hydrocyanation of alkenes remain underdeveloped.¹¹ Following early examples of highly enantioselective hydrocyanation of 2-methoxy-6-vinylnaphthalene in the context of naproxen synthesis,^{7–9} RajanBabu,¹² Schmalz,^{13,14} and Vogt^{14,15} have recently made seminal contributions using Ni catalysis (Figure 1B). To date, however, a general catalytic approach that meets the criteria of both broad reaction scope and high enantioselectivity remain elusive. In particular, internal alkenes, which would lead to a substantially wider range of useful products, have proven to be challenging substrates.

Recently, we have established electrocatalysis^{16–19} as a broadly applicable strategy for the difunctionalization of alkenes.²⁰ This approach merges the electrochemical generation of radical intermediates with catalyst-controlled radical addition to the alkene. Specifically, we devised anodically coupled electrolysis—a process that combines two anodic events to form two distinct

radical intermediates—for the synthesis of a diverse suite of vicinally heterodifunctionalized structures.^{21,22} Further applying electrocatalysis to the *hydrofunctionalization* of alkenes, such as hydrocyanation, would substantially expand the scope of electrosynthesis. Such transformations would enable direct access to a specific monofunctionalized product in one step from the alkene precursor, and are therefore complementary to difunctionalization reactions in the context of target-oriented synthesis. Achieving this reaction enantioselectively would further increase its synthetic value and grant access to valuable chiral targets for applications in organic synthesis and medicinal chemistry. To date, electrochemically driven hydrofunctionalization of alkenes is unknown, and electrosynthetic methods that enable asymmetric alkene functionalization remain elusive.²³⁻²⁵ Against this backdrop, we describe an electrochemical approach for the enantioselective hydrocyanation of conjugated alkenes powered by a Co/Cu dual electrocatalytic process.




Figure 1. Enantioselective hydrocyanation: synthetic significance (A) and proposed strategy (B).

Results and Discussion

Reaction design. Achieving the desired hydrocyanation using our electrocatalytic strategy would require the parallel generation of two open-shell species that serve as $\text{H}\cdot$ and $\text{CN}\cdot$ equivalents. Toward this goal, we envisioned combining two metal-mediated elementary radical reactions in an anodically coupled electrocatalytic system (Figure 2A). In the proposed catalytic cycle, $[\text{Co}^{\text{III}}]-\text{H}$ (formally), generated from a Co^{III} precursor and a hydrosilane,²⁶⁻³⁰ reacts with an alkene substrate via hydrogen atom transfer (HAT) to produce carbon-centered radical (**I**).³¹⁻³³ This intermediate then enters the cyanation cycle and undergoes single-electron oxidative addition to $[\text{Cu}^{\text{II}}]-\text{CN}$ to form **II**, a formally Cu^{III} intermediate. Subsequent reductive elimination completes the hydrocyanation reaction.^{34,35} This reaction design was inspired by Shenvi's seminal work showing that HAT-initiated alkene functionalization could intercept a second organometallic cycle in the context of Co/Ni-catalyzed, overall redox-neutral hydroarylation.³⁶ Critically, our desired hydrocyanation reaction is an overall oxidative transformation, which requires the turnover of both catalysts via a pair of single-electron oxidation events to return the resultant Co^{II} and Cu^{I} species

back to their reactive Co^{III} and Cu^{II} oxidation states. The key to successfully achieving this reaction thus relies on the identification of conditions that seamlessly accommodate both the Co HAT and Cu cyanation cycles.




Figure 2. Reaction design (A), discovery, and optimization (B). ^aYields determined by ¹H NMR. ^b U_{cell} : cell voltage applied between the cathode and anode; $U_{\text{cell}} = 2.3$ V corresponds to *ca.* 0.24 V vs $\text{Fc}^{+/-}$ anodic potential (E_{anode}) and gives *ca.* 2 mA initial current. ^cIsolated yield.

We note that the attributes of electrochemistry render it uniquely capable of facilitating the merger of the two aforementioned catalytic processes into a broadly useful hydrocyanation protocol. Co-catalyzed hydrofunctionalization can encounter chemoselectivity issues with styrene-type substrates (*vide infra*), particularly in the presence of an oxidant, likely because the benzylic radical formed upon HAT is susceptible to over-oxidation to the benzyl cation³⁷ (except when the reaction between the benzylic radical and oxidant is desirable). Cu-catalyzed cyanation *requires* a potent chemical oxidant^{34,35} which can also limit the reaction scope to substrates lacking oxidatively labile functional groups. By contrast, electrochemical strategies, which allow one to dial in a minimally sufficient potential, circumvent the need for stoichiometric oxidants and eliminate formation of undesired side products.

Reaction discovery and development. We set out to evaluate various combinations of Co and Cu catalysts. The electrochemical properties of many Co complexes have been well-documented.³⁸ However, the electrochemical compatibility and behavior of Cu complexes with common chiral ligands are largely unknown. In addition, Cu ions are highly susceptible to electroplating on the cathode [$E(\text{Cu}^{2+}/\text{Cu}^0) \approx -0.3$ V vs ferrocenium/ferrocene, $\text{Fc}^{+/-}$]. In an undivided electrochemical cell (e.g., a standard laboratory flask), an ideal vessel for preparative-scale electrochemical reactions, Cu complexes could be readily reduced at the cathode and lose catalytic activity.³⁹ Through systematic optimization, we found that Co(salen) complex **3** (0.5 mol%) and Cu(OTf)₂

(5 mol%) together with bisoxazoline (BOX) type ligands (**5–7**; 10 mol%) in the presence of PhSiH₃ and TMSCN promoted the conversion of 4-*tert*-butylstyrene (**1**) to the desired product (**2**) in good yield and moderate enantioselectivity (50–72% ee; Figure 2B). To further optimize enantioselectivity, we surveyed a catalog of chiral bidentate *N,N*- and *P,P*-ligands, including several that are known to be excellent ligands for Cu in enantioselective Lewis acid⁴⁰ or radical catalysis.^{34,35,41,42} However, the optimal ee could not be improved above 72% (ligand **7**). During our reaction optimization, we discovered that ester-substituted bisoxazolines (e.g., **4**) are excellent ligands in the desired hydrocyanation reaction. These ligands are readily prepared from serine and have not been studied previously in asymmetric catalysis.⁴³ The use of ligand **4** under otherwise identical conditions drastically improved the reaction enantioselectivity to 91%. We therefore hypothesize that the second-sphere ester groups provide attractive noncovalent interactions to stabilize the enantio-determining transition state for the C–CN bond formation (*vide infra*).


A number of control experiments serve to highlight the importance of each component of the reaction system. For example, using MeCN instead of DMF as the solvent, LiClO₄ instead of TBABF₄ as the electrolyte, or EtOH instead of HOAc as the proton source led to decreased product yield (Figure 2B). In each of these variations, we observed metallic Cu deposition on the cathode. Notably, the product was also formed in substantially lower ee in the presence of LiClO₄, showing that the polarity of the reaction medium influences the reaction enantioselectivity. When Co loading was increased from 0.5 to 2 mol%, the yield of **2** was dramatically decreased and a large amount of hydrogenated side product was observed.³³ This result highlights the importance of balancing the rates of the HAT and cyanation events to ensure optimal product selectivity. Finally, the reaction requires both Cu and Co to operate, as excluding either catalyst led to minimal conversion to the hydrocyanation product.

Reaction scope and application. Under optimal conditions, a variety of vinylarenes underwent hydrocyanation to furnish desired arylpropionitriles with excellent enantioselectivity (Figure 3A). The efficiency of the transformation was found to be relatively independent of the electronic properties of the aryl substituents. Importantly, the very mild reaction conditions imparted by the combination of electrochemistry and a radical mechanism made accessible a broad scope of enantioenriched nitriles with a diverse range of functional groups. In particular, functionalities that are susceptible to oxidative degradation under chemical conditions, such as electron-rich arenes (e.g., **8, 15, 20**), sulfides (**11**), and aldehydes (**13**), can be readily engaged in the hydrocyanation. Furthermore, groups including benzyl chlorides (**12**), aryl halides (**9, 25**), and aryl boronates (**17**), which might induce catalyst promiscuity under Ni-promoted conditions, were well tolerated in our reaction. Alkenes with various heterocycles (**20–23, 25**) also proved excellent substrates.


Importantly, our reaction is also applicable to internal alkenylarenes (Figure 3B), which are typically challenging substrates in previously reported enantioselective hydrocyanation reactions due to their low reactivity.¹³ Our protocol shows very broad substrate generality, furnishing products with a variety of synthetically useful functional handles (**28, 29, 34, 38**) and *N*-heterocyclic motifs (**43**). The stereochemistry of the starting alkenes is not important, and when applicable, we subject to the reaction a mixture of *E/Z*-isomers obtained directly from the olefination reaction. Both electron-rich (**37**) and electron-deficient alkenes (**39–44**) proved suitable substrates. In particular, hydrocyanation was achieved for cinnamoyl-type structures including esters (**39, 43, 44**), ketones (**40, 41**), and nitriles (**42**). These results demonstrate that our reaction can serve as a more general alternative to the enantioselective Michael addition en route to similar products. Considering that numerous bioactive natural products and their synthetic precursors

contain cinnamoyl side chains (e.g., taxol, phyllanthocin), our method provides a convenient means to derivatize these compounds for potential medicinal chemistry studies.

(A) Terminal alkenes:



(B) Internal alkenes:



(C) Other conjugated alkenes:




Figure 3. Reaction scope. Conditions: alkene (0.2 mmol, 1 equiv), PhSiH₃ (1.1 equiv), TMSCN (2.0 equiv), Co **3** (0.5 mol%), Cu(OTf)₂ (5 mol%), ligand **4** (10 mol%), TBABF₄ (2.0 equiv), HOAc (5.0 equiv), DMF (4.0 mL), C anode, Pt cathode, undivided cell, 0 °C, $U_{\text{cell}} = 2.3$ V. Isolated yields are reported, unless otherwise noted. ^aOptimal yield obtained at $U_{\text{cell}} = 1.8$ V (see discussion below). ^bWith 1 mol% Co **3**. ^cWith 2 mol% Co **3**. ^dCondition A: with 1.5 equiv TMSCN, 1.1 equiv PhSiH₃; yield determined by ¹H NMR. ^eCondition B: with 4.0 equiv TMSCN, 2.2 equiv PhSiH₃. Abbreviations: rr = regiometric ratio, de = diastereomeric excess.

In addition to representing a conceptual advance, our electrochemical hydrocyanation—with its highly general reaction scope—is also synthetically valuable and provides a complementary route to existing methods for the synthesis of chiral nitriles. For examples, chiral arylpropionitriles of similar structures can, in theory, be constructed via a recently developed method that relies on Cu-catalyzed benzylic C–H cyanation using *N*-fluorobenzenesulfonimide (NFSI) as the terminal oxidant.³⁴ Although an elegant reaction design and a powerful method, this protocol employs strongly oxidizing NFSI and necessitates the intermediacy of an electrophilic N-centered radical as the H-atom abstractor. These requirements impose several limitations on the reaction scope. Our electrochemical approach provides an opportunity to overcome these limitations and makes accessible a complementary scope of enantioenriched nitriles. For example, substrates with oxidatively labile functional groups, such as sulfides (**11**) and electron-rich (hetero)arenes (e.g., **8, 15, 20**), can be converted to the corresponding nitriles without undesired oxidative degradation. In addition, our reaction design engages mild Co-mediated HAT to access the key benzylic radicals, thus circumventing the C–H abstraction pathway required for the chemical approach. As a result, substrates with multiple benzylic sites (e.g., **16, 26**) can take part in the cyanation reaction in a site-specific manner. Furthermore, substrates with electron-deficient arenes (e.g., **13, 18, 21**), which are challenging in the chemical cyanation reaction likely because the benzylic C–H is acidic and polarity-mismatched with the NFSI-derived H-atom abstractor, can be readily engaged in our hydrocyanation reaction.

We further extended the scope of our reaction to other types of unsaturated substrates (Figure 3C). For example, dienes were converted to mono-hydrocyanated products in high efficiency and good to excellent enantioselectivity. In the case of linear dienes (**45–49**), the reaction selectively functionalizes the terminal C=C π -bond. Interestingly, in these transformations, enantioselective Cu-mediated C–CN bond formation is achieved via an allyl radical species, a process that is unknown in the literature. Structurally analogous enynes⁴⁴ also reacted chemoselectively in good ee (**50**). Finally, we expanded the reaction scope to allene substrates. The hydrocyanation of phenyllallene (**51**) yielded cinnamyl cyanide (**52**) as the predominant product. This result suggests that the terminal C=C is more reactive and the HAT step preferentially generates a primary allyl radical intermediate.⁴⁵ This selectivity allowed us to achieve double hydrocyanation through the use of an excess of reagents to generate 1,3-dinitrile **53** with high enantioselectivity. A radical rearrangement experiment provided key support for the intermediacy of the organic radical species. Exposure of cyclopropane-derived alkene **54** to the reaction conditions led to the expected ring opening, furnishing alkene-containing nitrile **55** in high yield and enantioselectivity.

Alkyl nitriles are highly versatile synthetic intermediates that can be readily transformed to a variety of useful functional groups, such as aldehydes, alcohols, carboxylic acids, esters, amines, and heterocycles. Here we showcase the synthetic value of our reaction products through three examples (Figure 4). Product **15**, synthesized on a 2-mmol scale (~0.4 grams), was readily converted to either naproxen methyl ester **56** via methanolysis or to benzindoline **57** via reduction followed by Pd-catalyzed C–H amination.⁴⁶ Both derivatizations occurred with complete retention of stereochemistry. Nitrile **18** was subjected to reduction and subsequent cyclization to yield dihydroisoquinolone **58**. These products bear structural resemblance to high-value drug molecules, again highlighting the significance of the enantioselective hydrocyanation reaction.





Figure 4. Product derivatization.


Electrochemical control of reaction chemoselectivity. A key advantage of electrochemistry is the ability to achieve external control over the electrode potential input. This feature allowed us to regulate the hydrocyanation chemoselectivity of electron-rich vinylarenes (Figure 5A). For example, using 3-vinyl-*N*-Ts-indole as the substrate, under our standard conditions with the application of a cell voltage of 2.3 V ($E_{\text{anode}} \sim 0.24$ V vs $\text{Fc}^{+/0}$), we observed full conversion of the alkene but minimal formation of desired nitrile **20**. Instead, several side products were formed, including those that arose from over-oxidation of the aryl radical intermediate to the corresponding cation followed by nucleophilic trapping (e.g., formyl ester and ketone products). By lowering the voltage input from 2.3 V to 1.8 V ($E_{\text{anode}} \sim 0.08$ V), we successfully suppressed overoxidation and obtained hydrocyanation product **20** in 71% yield. This principle was also applied to the optimization of reactions forming **8** and **59**.

For comparative purposes, we examined the hydrocyanation of alkene **1** under traditional chemical conditions without an electrical input. None of the chemical oxidants we surveyed promoted the reaction with efficiencies or enantioselectivities comparable to our protocol (Figure 5B). In cases where a low yield of **2** was obtained (9-24%), the enantioselectivity was also substantially lower (40-68% ee). This erosion of ee is likely a result of a racemic carbocation pathway caused by over-oxidation of the key benzylic radical intermediate. We note that, although these results could potentially be improved through extensive optimization of reaction conditions, such optimization studies would need to address the challenges associated with the use of a terminal chemical oxidant. *Electrochemistry, with its ability to enable electron transfer with a minimally sufficient potential input, provides an ideal means to circumvent issues with chemical oxidation and harness multiple redox catalytic cycles for productive chemistry.*

(A) Improving chemoselectivity by potential control



(B) Comparison with common chemical oxidants



Entry	Oxidant	Conversion (%) ^a	Yield (%) ^a	Ee (%)
1	anode	92	87	85
2	NFSI	38	<5	— ^c
3	TBHP	85	15	68
4	[F ⁺]	78	24	63
5	PhI(OAc) ₂	63	<5	— ^c
6	Mn(OAc) ₃	86	9	40
7	Cu(OAc) ₂	50	15	52
8	O ₂ (balloon)	>95	<5	— ^c
9	[I ⁺]	79	14	62
10	CAN	75	<5	— ^c
11	Fc ⁺ BF ₄ [−]	8	<5	— ^c




Figure 5. Electrochemical tuning of reaction chemoselectivity (A) and comparison with chemical methods (B). ^aDetermined by ¹H NMR. ^bIsolated yield. ^cNot determined.

Stereochemical model. Finally, we enlisted theoretical methods to elucidate the role played by serine-derived BOX ligands in directing the highly enantioselective Cu-mediated cyanation. A previous study showed that the radical combination to generate intermediate **II** (Figure 2) is facile, whereas reductive elimination to construct the C–CN bond is slow and thus enantio-determining.³⁴ Traditionally, BOX-type ligands are thought to induce enantioselectivity through primarily *repulsive* steric interactions imparted by the bulky substituents α -to the N coordination sites. Considering that the ester-derived ligands (**4**, **60**) perform substantially better than the canonical BOX ligands with various steric profiles (e.g., **5**–**7**), we hypothesized that the ester groups must also provide *attractive* electrostatic interactions with the substrate assembly in the key C–CN

forming transition state. Density functional theory computations are fully consistent with this hypothesis (Figure 6 and Supplementary Information). Using the reaction forming **10** as an example, the lowest energy transition state structures leading to the major (*R*) and minor (*S*) products (TS_{*R*} and TS_{*S*}, respectively) both display a favorable C–H···π interaction between the acidic proton α -to one of the ester groups and the aryl group on the substrate. This attractive non-covalent interaction dictates the geometry of both transition states by positioning the alkyl substituent on the substrate toward the 'Bu-ester on the opposite half of the catalyst. However, this arrangement causes a more severe steric interaction between these two adjacent groups in TS_{*S*} than TS_{*R*}. In fact, to mitigate this detrimental interaction in TS_{*S*}, the ester group deviates from its optimal planar geometry (dihedral angle $\theta = 22^\circ$), therefore breaking conjugation. Based on these findings, we further hypothesize that this interplay between attractive and repulsive non-covalent interactions in the transition state determines the reaction enantioselectivity.




Figure 6. Computational stereochemical model. Numbers in red indicate interatomic distances that are shorter than the sum of the van der Waals radii of the two atoms or dihedral angles that show significant deviation from their optimal values.

This stereochemical model is consistent with our experimental observations. For example, reactions with internal alkenes are in general more enantioselective than those with terminal alkenes, which can be explained by the increased steric interaction between the substrate alkyl group and the catalyst ester group in the minor TS for the internal alkenes. In addition, substrates with more electron-rich aryl groups (e.g., **2**, **8**) in general lead to higher enantioselectivity than those with electron-deficient groups (e.g., **9**, **13**), which can be rationalized by an enhanced C–H···π interaction involving the aryl group.⁴⁷

Conclusion

The electrocatalytic hydrocyanation described herein reveals the unique advantages of harnessing electrochemistry in exploring new chemical space and developing solutions to challenging synthetic problems. This new transformation will enhance chemists' access to a diverse array of enantioenriched nitriles. On a fundamental level, the realization of electrochemical hydrofunctionalization and demonstration of enantioselective electrocatalysis will improve the scope of electrosynthesis and its adoption in modern organic chemistry.

References:

1. Rajanbabu, T. V. Hydrocyanation of alkenes and alkynes. *Org. React.* **75**, 1-73 (2011).
2. Nugent, W. A. & McKinney, R. J. Nickel-catalyzed Markovnikov addition of hydrogen cyanide to olefins. Application to nonsteroidal anti-inflammatories. *J. Org. Chem.* **50**, 5370-5372 (1985).
3. Gaspar, B. & Carreira, E. M. Mild cobalt-catalyzed hydrocyanation of olefins with tosyl cyanide. *Angew. Chem. Int. Ed.* **46**, 4519-4522 (2007).
4. Fang, X., Yu, P. & Morandi, B. Catalytic reversible alkene-nitrile interconversion through controllable transfer hydrocyanation. *Science* **351**, 832-836 (2016).
5. Pollak, P., Romeder, G., Hagedorn, F. & Gelbke, H. "Nitriles," in *Ullman's Encyclopedia of Industrial Chemistry* (Wiley-VCH, Weinheim, Germany, ed. 5, 1985), vol. A17, p. 363.
6. Tolman, C. A., McKinney, R. J., Seidel, W. C., Druliner, J. D. & Stevens, W. R. Homogeneous nickel-catalyzed olefin hydrocyanation. *Adv. Catal.* **33**, 1-46 (1985).
7. RajanBabu, T. V. & Casalnuovo, A. L. Tailored ligands for asymmetric catalysis. *J. Am. Chem. Soc.* **114**, 6265-6266 (1992).
8. RajanBabu, T. V. & Casalnuovo, A. L. Role of electronic asymmetry in the design of new ligands: the asymmetric hydrocyanation reaction. *J. Am. Chem. Soc.* **118**, 6325-6326 (1996).
9. Casalnuovo, A. L., RajanBabu, T. V., Ayers, T. A. & Warren, T. Ligand electronic effects in asymmetric catalysis: enhanced enantioselectivity in asymmetric hydrocyanation of vinylarenes. *J. Am. Chem. Soc.* **116**, 9869-9882 (1994).
10. Kurono, N. & Ohkuma, T. Catalytic asymmetric cyanation reactions. *ACS Catal.* **6**, 989-1023 (2016).
11. Bini, L., Müller, C. & Vogt, D. Ligand development in the Ni-catalyzed hydrocyanation of alkenes. *Chem. Commun.* **46**, 8325-8334 (2010).
12. Saha, B. & RajanBabu, T. V. Nickel(0)-catalyzed asymmetric hydrocyanation of 1,3-dienes. *Org. Lett.* **8**, 4657-4659 (2006).
13. Falk, A., Göderz, A.-L. & Schmalz, H.-G. Enantioselective nickel-catalyzed hydrocyanation of vinylarenes using chiral phosphine-phosphite ligands and TMS-CN as a source of HCN. *Angew. Chem. Int. Ed.* **52**, 1576-1580 (2013).
14. Falk, A., Cavalieri, A., Nichol, G. S., Vogt, D. & Schmalz, H.-G. Enantioselective nickel-catalyzed hydrocyanation using chiral phosphine-phosphite ligands: recent improvements and insights. *Adv. Synth. Catal.* **357**, 3317-3320 (2015).
15. Wilting, J., Janssen, M., Müller, C. & Vogt, D. The enantioselective step in the Nickel-catalyzed hydrocyanation of 1,3-cyclohexadiene. *J. Am. Chem. Soc.* **128**, 11374-11375 (2006).
16. Francke, R. & Little, R. D. Redox catalysis in organic electrosynthesis: basic principles and recent developments. *Chem. Soc. Rev.* **43**, 2492-2521 (2014).
17. Jutand, A. Contribution of electrochemistry to organometallic catalysis. *Chem. Rev.* **108**, 2300-2347 (2008).

18. Meyer, T. H., Finger, L. H. Finger., Gandeepan, P. & Ackermann, L. Resource economy by metallaelectrocatalysis: merging electrochemistry and C–H activation. *Trends Chem.* **1**, 63-76 (2019).
19. Yan, M., Kawamata, Y. & Baran, P. S. Synthetic organic electrochemical methods since 2000: on the verge of a renaissance. *Chem. Rev.* **117**, 113230-13319 (2017).
20. Fu, N., Sauer, G. S., Saha, A., Loo, A. & Lin, S. Metal-catalyzed electrochemical diazidation of alkenes. *Science* **357**, 575-579 (2017).
21. Ye, K., Pombar, G., Fu, N., Sauer, G. S., Keresztes, I. & Lin, S. Anodically coupled electrolysis for the heterodifunctionalization of alkenes. *J. Am. Chem. Soc.* **140**, 2438-2441 (2018).
22. Fu, N., Song, L., Liu, J., Shen, Y., Siu, J. C. & Lin, S. Mn-catalyzed electrochemical chloroalkylation of alkenes. *ACS Catal.* **9**, 746-754 (2019).
23. Nguyen, B. H., Redden, A. & Moeller, K. D. Sunlight, electrochemistry, and sustainable oxidation reactions. *Green Chem.* **16**, 69-72 (2014).
24. Nguyen, B. H., Perkins, R. J., Smith, J. A. & Moeller, K. D. Photovoltaic-driven organic electrosynthesis and efforts toward more sustainable oxidation reactions. *Beilstein J. Org. Chem.* **11**, 280-287 (2015).
25. Tanaka, H., Kuroboshi, M., Takeda, H., Kanda, H. & Torii, S. Electrochemical asymmetric epoxidation of olefins by using an optically active Mn-salen complex. *J. Electroanal. Chem.* **507**, 75-81 (2001).
26. Crossley, S. W. M., Obradors, C., Martinez, R. M. & Shenvi, R. A. Mn-, Fe-, and Co-catalyzed radical hydrofunctionalizations of olefins. *Chem. Rev.* **116**, 8912-9000 (2016).
27. Isayama, S. & Mukaiyama, T. Hydration of olefins with molecular oxygen and triethylsilane catalyzed by bis-(trifluoroacetylacetonato)cobalt(II). *Chem. Lett.* **18**, 569-572 (1989).
28. Waser, J., Nambu, H. & Carreira, E. M. Cobalt-catalyzed hydroazidation of olefins: convenient access to alkyl azides. *J. Am. Chem. Soc.* **127**, 8294-8295 (2005).
29. Renata, H., Zhou, Q. & Baran, P. S. Strategic redox relay enables a scalable synthesis of ouabagenin, a bioactive cardenolide. *Science* **339**, 56-63 (2013).
30. Shigehisa, H., Koseki, N., Shimizu, N., Fujisawa, M., Niitsu, M. & Hiroya, K. Catalytic hydroamination of unactivated olefins using a Co catalyst for complex molecule synthesis. *J. Am. Chem. Soc.* **136**, 13534-13537 (2014).
31. Li, G., Han, A., Pulling, M. E., Estes, D. P. & Norton, J. R. Evidence for formation of a Co–H bond from $(\text{H}_2\text{O})_2\text{Co}(\text{dmgBF}_2)_2$ under H_2 : application to radical cyclizations. *J. Am. Chem. Soc.* **134**, 14662-14665 (2012).
32. Crossley, S. W. M., Barabé, F. & Shenvi, R. A. Simple, chemoselective, catalytic olefin isomerization. *J. Am. Chem. Soc.* **136**, 16788-16791 (2014).
33. King, S. M., Ma, X. & Herzon, S. B. A method for the selective hydrogenation of alkenyl halides to alkyl halides. *J. Am. Chem. Soc.* **136**, 6884-6887 (2014).
34. Zhang, W., Wang, F., McCann, S. D., Wang, D., Chen, P., Stahl, S. S. & Liu, G. Enantioselective cyanation of benzylic C–H bonds via copper-catalyzed radical relay. *Science* **353**, 1014-1018 (2016).
35. Wang, F., Chen, P. & Liu, G. Copper-catalyzed radical relay for asymmetric radical transformations. *Acc. Chem. Res.* **51**, 2036-2046 (2018).
36. Green, S. A., Matos, J. L. M., Yagi, A. & Shenvi, R. A. Branch-selective hydroarylation: iodoarene–olefin cross-coupling. *J. Am. Chem. Soc.* **138**, 12779-12782 (2016).
37. Shigehisa, H., Kikuchi, H. & Hiroya, K. Markovnikov-selective addition of fluororous solvents to unactivated olefins using a Co catalyst. *Chem. Pharm. Bull.* **64**, 371-374 (2016).

38. Eichhorn, E., Rieker, A. & Speiser, B. The electrochemical oxidation of $[\text{Co}^{\text{II}}(\text{salen})]$ in solvent mixtures—an example of a ladder scheme with coupled electron-transfer and solvent-exchange reactions. *Angew. Chem. Int. Ed.* **31**, 1215-1217 (1992).

39. Merchant, R. R., Oberg, K. M., Lin, Y., Novak, A. J. E., Felding, J. & Baran, P. S. Divergent synthesis of pyrone diterpenes via radical cross coupling. *J. Am. Chem. Soc.* **140**, 7462-7465 (2018).

40. Desimoni, G., Faita, G. & Jørgensen, K. A. Update 1 of: C_2 -Symmetric chiral bis(oxazoline) ligands in asymmetric catalysis. *Chem. Rev.* **111**, PR284-PR437 (2011).

41. Zhu, R. & Buchwald, S. L. Versatile enantioselective synthesis of functionalized lactones via copper-catalyzed radical oxyfunctionalization of alkenes. *J. Am. Chem. Soc.* **137**, 8069-8077 (2015).

42. Zhang, Z., Stateman, L. M. & Nagib, D. A. δ C–H (hetero)arylation via Cu-catalyzed radical relay. *Chem. Sci.* **10**, 1207-1211 (2019).

43. Perrotta, D., Wang, M.-M. & Waser, J. Lewis acid catalyzed enantioselective desymmetrization of donor–acceptor *meso*-diaminocyclopropanes. *Angew. Chem. Int. Ed.* **57**, 5120-5123 (2018).

44. Lu, F.-D., Liu, D., Zhu, L., Lu, L.-Q., Yang, Q., Zhou, Q.-Q., Wei, Y., Lan, Y. & Xiao, W.-J. Asymmetric propargylic radical cyanation enabled by dual organophotoredox and copper catalysis. *J. Am. Chem. Soc.* **141**, 6167-6172 (2019).

45. Zhu, N., Wang, F., Chen, P., Ye, J. & Liu, G. Copper-catalyzed intermolecular trifluoromethylazidation and trifluoromethylthiocyanation of allenes: efficient access to CF_3 -containing allyl azides and thiocyanates. *Org. Lett.* **17**, 3580-3583 (2015).

46. He, G., Zhao, Y., Zhang, S., Lu, C. & Chen, G. Highly efficient syntheses of azetidines, pyrrolidines, and indolines via palladium catalyzed intramolecular amination of $\text{C}(\text{sp}^3)\text{–H}$ and $\text{C}(\text{sp}^2)\text{–H}$ bonds at γ and δ positions. *J. Am. Chem. Soc.* **134**, 3-6 (2012).

47. Bloom, J. W. G., Raju, R. K. & Wheeler, S. E. Physical nature of substituent effects in XH/π interactions. *J. Chem. Theory Comput.* **8**, 3167-3174 (2012).

Acknowledgments: Financial support to S.L. was provided by NIGMS (R01GM130928) and Eli Lilly. Financial support to B.G.E. and R.A.D. was provided by start-up funding from Cornell University. This study made use of the NMR facility supported by the NSF (CHE-1531632) and resources of the National Energy Research Scientific Computing Center (NERSC), a U.S. Department of Energy Office of Science User Facility operated under Contract No. DE-AC02-05CH11231. We thank Anthony Condo for GC-MS analysis, Yifan Shen and Jinjian Liu for providing some of the substrates, and IKA for the donation of ElectraSyn 2.0.

