



QUARTZ

FORENSIC BLOOD TOXICOLOGY PROFICIENCY TESTING SCHEME

Scheme Description

LGC Standards Proficiency Testing

1 Chamberhall Business Park
Chamberhall Green
Bury
Lancashire
BL9 0AP
United Kingdom

Telephone: +44 (0) 161 762 2500
Fax: +44 (0) 161 762 2501
Email: ptcustomerservices@lgcgroup.com
Website: www.lgcstandards.com



QUARTZ Scheme Description

Record of issue status and modifications

ISSUE	ISSUE DATE	DETAILS	AUTHORISED BY
2	01/09/08	Updated with UKAS logo for single scope (0001) and removed reference to scheme year. Added additional sample (standard solution).	T.Noblett
3	15/09/08	Addition of 6MAM, Trazodone, Clonazepam Propranolol, Benzyl Piperazine (BZP) to the scheme drugs scope.	T.Noblett
4	03/07/09	Details of blood used, pre-screening performed on blood and classification of drugs added. Trial material details also included. Operational issues common to all schemes moved into General Protocol. List of abbreviations added. New PORTAL system added	M.Whetton
5	18/06/10	List of potential drugs updated in appendix B. Inclusion of sample 4 (alcohol in blood)	M.Whetton
6	Oct 2010	Document appendices reference updated	M.Whetton
7	23/03/11	Change Of Address to Page 1	N.Stephenson
8	08/06/11	List of potential drugs updated in appendix B. Trial references removed for sample 4	M.Whetton
9	22/07/11	Document updated for 2011/2012	M.Whetton
10	18/01/12	SDPA values updated for alcohol assessment. Appendix A updated.	W.Gaunt
11	July 2013	Removal of Appendix B and expansion of Scheme Aims and Sample Descriptions. Update of Sample 3. Change of scheme year.	K Morgan
12	Feb 2014	Clarification of traceability. Introduction of New Psychoactive Substance Sample and Alcohol Technical Defence. 'Methods' section added.	K Morgan
13	Dec 2014	New formulation of NPS (sample 6) for New scheme year.	K Morgan
14	Sept 2015	Addition of subcontracting information. Removal of hard copy report information.	K Morgan N Mason
15	Jan 2016	Addition of samples 7 (Synthetic cannabinoid sample) and 8 (Urine identification sample)	K Morgan

Notes:

Where this document has been translated, the English version shall remain the definitive version

Scheme Aims and Organisation

The primary aim of the Forensic blood toxicology proficiency testing scheme (QUARTZ) is to enable laboratories performing the analysis of drugs in post-mortem and other blood samples for toxicological purposes, particularly in a forensic context, to monitor their performance and compare it with that of their peers. Case-types that will be covered by the scheme will include deaths, drug facilitated sexual assaults, impaired driving and other forensic toxicology criminal cases. The QUARTZ scheme also aims to provide information to participants on technical issues and methodologies relating to forensic toxicology.

The QUARTZ scheme year operates from April to March. Further information about QUARTZ, including test material availability, round despatch dates and reporting deadlines, are available on the current QUARTZ application form.

Test Materials

Test material batches are tested for homogeneity for at least one test parameter where deemed appropriate. Details of homogeneity tests performed and results are given in the QUARTZ Scheme Reports.

Samples are generally prepared using pre-screened human blood.

The non-clinical blood material is obtained from the National blood service and prior to use in the QUARTZ scheme is screened for:

- HEP B antigen
- HEP C antigen
- Combo HIV 1 and 2
- Syphilis
- Human T-lymphotropic virus (HTLV)
- Cytomegalovirus (CMV)

Note: All test materials provided are intended for use as proficiency testing materials only and are not to be used for any other purposes.

Some aspects of the scheme, such as test material production, homogeneity testing and stability assessment, can from time to time be subcontracted. When subcontracting occurs, it is placed with a competent subcontractor and LGC is responsible for this work. The planning of the scheme, the evaluation of performance and the authorisation of the final report will never be subcontracted.

Sample 1 (Identification): Participants will be asked to identify up to 4 drugs relevant to forensic toxicology. The drugs chosen will include anaesthetics (e.g. ketamine), antidepressants (e.g. amitriptyline, dothiepin, citalopram), anticholinergics (e.g. procyclidine), anticonvulsants (e.g. carbamazepine, phenytoin, pregabalin), antihistamines (e.g. cyclizine, chlorpheniramine), antipsychotics (e.g. olanzapine, quetiapine), cardiovascular drugs (e.g. atenolol, flecainide), erectile dysfunction (e.g. sildenafil), opioid analgesics (e.g. morphine, methadone, oxycodone, tramadol), Non Steroidal Anti-Inflammatory Analgesics (e.g. salicylate, paracetamol, etoricoxib), benzodiazepines (e.g. diazepam, alprazolam, temazepam, flunitrazepam), hypnotic drugs (e.g. zaleplon, zolpidem, zopiclone), stimulants (e.g. cocaine, piperazines, amphetamines, APB, cathinones), cannabinoids, barbiturates, carboxyhaemoglobin. The sample will always contain at least one drug from the most commonly encountered drugs in forensic toxicology casework.

QUARTZ Scheme Description

Sample 2 (Quantification and Case Study): Participants will be told the identity, or generic classification, of the drug(s), and asked to quantify the concentration. They will also be asked to give an interpretation of the results in respect to a case study provided. The drugs chosen will be from the same drug classes as those described for Sample 1 above. The sample will always contain at least one drug from the most commonly encountered drugs in forensic toxicology casework.

Sample 3 (Known Drug Blood Quantification Sample): This sample consists of two samples which are alternated each quarter at different concentrations. One of the samples contains 4 common drugs of abuse and the other sample contains 4 prescription drugs that are commonly encountered in forensic casework.

Drugs of Abuse Sample

- 1 Morphine
- 2 Methadone
- 3 Amphetamine
- 4 Diazepam

Prescribed Drugs Sample

- 1 Citalopram
- 2 Codeine
- 3 Amitriptyline
- 4 Tramadol

Sample 4 (Alcohol in blood): A blood sample containing a known concentration of ethanol is provided. Participants are requested to determine and report the concentration of the alcohol. Participants are also asked to report the concentration of fluoride in the sample.

Sample 5 (Alcohol Technical Defence): A paper based exercise for blood and breath alcohol concentrations.

Sample 6 (Known Drug Blood Quantification Sample - New Psychoactive Substances): This sample is distributed twice per year containing different substances.

Sample 7 (Synthetic cannabinoid sample: Identification sample): A blood sample containing a synthetic cannabinoid. Participants are asked to screen for synthetic cannabinoids and identify one of the most common synthetic cannabinoids. This sample is distributed twice per year with a different synthetic cannabinoid included in each distribution.

Sample 8 (Identification): Participants are asked to identify up to 4 drugs or metabolites relevant to forensic toxicology. The drugs chosen will include anaesthetics (e.g. ketamine), antidepressants (e.g. amitriptyline, dothiepin, citalopram), anticholinergics (e.g. procyclidine), anticonvulsants (e.g. carbamazepine, phenytoin, pregabalin), antihistamines (e.g. cyclizine, chlorpheniramine), antipsychotics (e.g. olanzapine, quetiapine), cardiovascular drugs (e.g. atenolol, flecainide), erectile dysfunction (e.g. sildenafil), opioid analgesics (e.g. morphine, methadone, oxycodone, tramadol), Non Steroidal Anti-Inflammatory Analgesics (e.g. salicylate, paracetamol, etoricoxib), benzodiazepines (e.g. diazepam, alprazolam, temazepam, flunitrazepam), hypnotic drugs (e.g. zaleplon, zolpidem, zopiclone), stimulants (e.g. cocaine, piperazines, amphetamines, APB, cathinones), cannabinoids, barbiturates, carboxyhaemoglobin. The sample will always contain at least one drug from the most commonly encountered drugs in forensic toxicology casework. Urine specimen.

QUARTZ Scheme Description

Statistical Analysis

Information on the statistics used in QUARTZ can be found in the General Protocol and in the Scheme Report. Methods for determining assigned values and the values for SDPA used for individual samples are given in Appendix A.

Methods

Methods are listed in Appendix A and PORTAL. Please select the most appropriate method from the list. If none of the methods are appropriate, then please report your method as 'Other' and record a brief description in the Comments Section in PORTAL.

Results and Reports

QUARTZ results are returned through our electronic reporting software, PORTAL, full instructions for which are provided by email. However, participants may request result submission forms on which to report and return results if they are unable to report through electronic means. This will incur an additional charge.

QUARTZ reports will be available on the website within 10 working days of round closure. Participants will be emailed a link to the report when it is available.

APPENDIX A - Description of abbreviations used

Assigned Value (AV)

The assigned value may be derived in the following ways:

- From the robust mean (RMean). This is the median of participant results after the removal of test results that are inappropriate for statistical evaluation, e.g. miscalculations, transpositions and other gross errors. Generally, the assigned value will be set using results from all methods, unless the measurement is considered method-dependant, in which case the assigned value will be set by method as illustrated in the report tables.

For some analytes, where there is a recognised reference method for that type of measurement, this may be used as the assigned value for a particular analyte i.e. it would be applied to results obtained by any method.

Traceability: Assigned values which are derived from the participant results, or a subset of the results are not traceable to an international measurement standard. The uncertainty of assigned values derived in this way is estimated from the participant results, according to ISO 13528.

- From a formulation value (Formulation). This denotes the use of an assigned value derived from sample preparation details, where known and exact quantities of analyte have been used to prepare the sample.

Traceability: Assigned values calculated from the formulation of the test sample are traceable, via an unbroken metrological traceability chain, to an international measurement standard. The measurement uncertainty of the assigned value is calculated using the contributions from each calibration in the traceability chain.

- From a qualitative formulation (Qual Form). This applies to qualitative tests where the assigned value is simply based on the presence/absence of the analyte in the test material.

Traceability: Assigned values calculated from the qualitative formulation of the test sample are traceable to a certified reference standard or a microbiological reference strain.

- From expert labs (Expert). The assigned value for the analyte is provided by an 'expert' laboratory.

Traceability: Assigned values provided by an 'expert' laboratory may be traceable to an international measurement standard, according to the laboratory and the method used. The uncertainty of measurement for an assigned value produced in this way will be provided by the laboratory undertaking the analysis. Details of traceability and the associated uncertainty will be provided in the report for the scheme/round.

Range

This indicates the concentration range at which the analyte may be present in the test material.

QUARTZ Scheme Description

SDPA

SDPA represents the 'standard deviation for proficiency assessment' which is used to assess participant performance for the measurement of each analyte. This may be a fixed value (as stated), a percentage (%) of the assigned value or based on the robust standard deviation of the participant measurement results, either across all methods or by method depending on whether the measurement made is method dependent (see assigned value).

Units

This indicates the units used for the assessment of data. These are the units in which participants should report their results. For some analytes in some schemes participants may have a choice of which units to report their results, however, the units stipulated in this scheme description are the default units to which any results reported using allowable alternative results will be converted to.

DP

This indicates the number of decimal places to which participants should report their measurement results.

Sample 1 Identification of toxicology drugs

Participants will receive: 10ml blood matrix, issued quarterly.

Analyte	Method	Range	AV	SDPA	Units	DP
Drug identification	All	All	Qual Form	N/A	N/A	N/A

Sample 2 Quantification of toxicology drugs and Case Study

Participants will receive: 10ml blood matrix, issued quarterly.

Analyte	Method	Range	AV	SDPA	Units	DP
Drug quantification	All	<0.1	Formulation	25% of AV	mg/L	3 Sig Fig
		0.1 to 10		20% of AV		
		>10		15% of AV		

Sample 3 Known Drug Blood Quantification Sample

Participants will receive: 10ml blood matrix, issued quarterly alternate samples of Drugs and Abuse and Prescription.

Analyte	Method	Range	AV	SDPA	Units	DP
Drug quantification Alternate samples containing: Sample 1: Morphine, Methadone, Amphetamine and Diazepam. Sample 2: Citalopram, Codeine, Amitriptyline and Tramadol	All	<0.1	Formulation	25% of AV	mg/L	2
		0.1 to 10		20% of AV		
		>10		15% of AV		

QUARTZ Scheme Description

Sample 4 Analysis of alcohol in blood (driving impairment)

Participants will receive 10ml blood matrix

Method	Test	Range	AV	SDPA	Units	DP
Quantification of alcohol	All	≤ 100 >100	Formulation	3 mg/100ml 3% of AV	mg/100ml (mg%)	2
Quantification of fluoride	All	1.0 to 2.0	Formulation	RMean	%	2

Sample 5 Alcohol Technical Defence**

Participants will receive: Information and/or analytical results provided for paper exercise

Analyte	Method	Range	AV	SDPA	Units	DP	Expected outcome
Alcohol in Blood Alcohol technical defence (ATD) “back calculations”	Interpretation	N/A	Robust Mean	N/A	N/A	N/A	Advisory

**Please note that this sample is not currently within our UKAS scope of accreditation.

Sample 6**

New Psychoactive Substance (NPS) Blood Quantification Sample

Participants will receive: 10ml blood matrix, issued every six months (two rounds per year).

Analyte	Method	Range	AV	SDPA	Units	DP
Drug quantification of 4 New Psychoactive Substances. Round 1: Mephedrone Etizolam Methiopropamine (MPA) 6-(2-aminopropyl)benzofuran (6-APB) Round 2: Diclazepam Ethylphenidate 4-Methylethcathinone 1-(Benzofuran-5-yl)-N-methylpropan-2-amine (5-MAPB)	All	<0.1	Formulation	25% of AV	µg/L	2
		0.2 to 10		20% of AV		
		>10		15% of AV		

**Please note that this sample is not currently within our UKAS scope of accreditation.

Sample 7**

Synthetic Cannabinoid Blood Sample: Qualitative

Participants will receive: 10ml blood matrix, issued every six months.

Analyte	Method	Range	AV	SDPA	Units	DP
Drug identification	All	All	Qual Form	N/A	N/A	N/A

**Please note that this sample is not currently within our UKAS scope of accreditation.

Sample 8

Identification of toxicology drugs/metabolites – Urine specimen

Participants will receive: 10ml Urine, issued every six months.

Analyte	Method	Range	AV	SDPA	Units	DP
Drug identification	All	All	Qual Form	N/A	N/A	N/A