

Exclusion Procedure for Illness & Communicable Diseases

Disease/Illness	Recommended Exclusion Period	Comments
Chicken Pox	Until all spots have scabbed / healed over	See Vulnerable children
Conjunctivitis	None however for younger tactile children best practise to advise to keep at home until eyes are not weeping	
Diarrhoea & Vomiting	Must be clear for 48 hours after last episode	
Gastro-enteritis	Must be clear for 48 hours or until advised by relevant public health official	See Vulnerable children. Seek advice if over 4 children affected.
Hand, Foot & Mouth	None	Seek advice if over 4 children affected.
Headlice	No exclusion as long as parents are treating	
Impetigo	Until skin is healed over or 48 hours after starting antibiotic treatment	
Measles	4 days from onset of rash	
Mumps	Until the swelling has gone down – 5 days minimum	
Rash	Children with rashes should be considered infectious and assessed by their doctor.	
Ringworm	None as long as parents are treating	
Scabies	Child can return after first treatment	
Scarlet Fever	24 hours after commencing appropriate antibiotic treatment	If more than one child affected seek advice
Shingles	Until weeping has stopped and feeling well	
Slapped Cheek/ Fifth Disease/ Parvovirus B19	None once rash has developed	See Vulnerable children. Any pregnant contacts of case should seek advice from GP.
Temperature	If sent home, next 24 hours at home	
Threadworms	None	
TB	2 weeks following start of treatment or until declared free from illness by public health official	
Warts & Verrucae	No exclusion but should be treated and covered	