

**Supporting Information**  
**for**

Application of Continuous Flow-Flash Chemistry to Scale-up Synthesis of  
5-Cyano-2-formylbenzoic acid

*Masaki, Seto,\*‡Shinichi Masada,‡ Hirotsugu Usutani,\$, // David G. Cork,\$ # Koichiro  
Fukuda,‡,† and Tetsuji Kawamoto\*,‡,†*

‡Neuroscience Drug Discovery Unit, Research, Takeda Pharmaceutical Company  
Ltd., 26-1, Muraoka-Higashi 2-chome, Fujisawa, Kanagawa 251-8555, Japan

§Process Chemistry, Pharmaceutical Sciences, Takeda Pharmaceutical Company  
Ltd., 26-1, Muraoka-Higashi 2-chome, Fujisawa, Kanagawa 251-8555, Japan

**Present Address**



† Chemistry Research Division, Axcelead Drug Discovery Partners, Inc. 26-1,  
Muraoka-Higashi 2-chome, Fujisawa, Kanagawa 251-0012, Japan (K.F. and T.K.)

// Kyoto University Original Co., Ltd., Kyoto University, Yoshida-Hommachi,  
Sakyo-ku, Kyoto 606-8501, Japan (H.U.)



#2-4-56 Ueno Higashi, Toyonaka, Osaka, 560-0013, Japan (D.G.C.)

**Table of contents**



1. Copies of <sup>1</sup>H and <sup>13</sup>C NMR spectra of compounds **1**, **8b**, **8c**, and **8d** ..... S2

**5-Cyano-2-formylbenzoic acid (6-cyano-3-hydroxy-2-benzofuran-1(3H)-one, 1).**

## Ethyl 2-bromo-5-cyanobenzoate (8b).



## Isopropyl 2-bromo-5-cyanobenzoate (8c).



***tert*-Butyl 2-bromo-5-cyanobenzoate (8d).**