

Supplementary Materials

Discovery of triazolo-pyridazine/-pyrimidine derivatives bearing aromatic (heterocycle)-coupled azoles units as Class II c-Met inhibitors

Qian Zhang^{a,1}, Xiaobo Liu^{a,1}, Wenhui Gan^a, Jinjin Wu^a, Hualan Zhou^a, Zunhua Yang^b, Yiling Zhang^b, Min Liao^a, Ping Yuan^a, Shan Xu^a, Pengwu Zheng^{a*}, Wufu Zhu^{a*}

^a Jiangxi Provincial Key Laboratory of Drug Design and Evaluation, School of Pharmacy, Jiangxi Science & Technology Normal University, 605 Fenglin Road, Nanchang, Jiangxi 330013, China ^b College of Pharmacy, Jiangxi University of Traditional Chinese Medicine, Nanchang, 330004, China

* **Corresponding author.** Tel. & Fax: +86 791 8380-2393; E-mail: zhengpw @126.com (P. Zheng), zhuwufu-1122@163.com (W. Zhu)

^1H NMR Spectral

Figure S1: 12c. ^1H NMR (400 MHz). Solvent: $\text{DMSO-}d_6$




Figure S2: 12e. ¹H NMR (400 MHz). Solvent: DMSO-*d*₆




Figure S3: 13e. ^1H NMR (400 MHz). Solvent: $\text{DMSO-}d_6$




Figure S4: 18f. ^1H NMR (400 MHz). Solvent: $\text{DMSO-}d_6$





Figure S5: 19e. ^1H NMR (400 MHz). Solvent: $\text{DMSO-}d_6$



TOF MS analytical data

Figure S6: 13b Solvent: CH₃OH.

TOF MS ES+ (m/z): [M + H]⁺, calcd for C₂₃H₁₄ClF₄N₇O₂: 532.0912, found, 532.0919.




Figure S7: 13c Solvent: CH₃OH.

TOF MS ES+ (m/z): [M + H]⁺, calcd for C₂₃H₁₇FN₆O₂S: 461.1196, found, 461.1197.




Figure S8: 13g Solvent: CH₃OH.

TOF MS ES+ (m/z): [M + H]⁺, calcd for C₂₂H₁₆FN₇O₂S: 462.1148, found, 462.1147.

