

SUPPORTING INFORMATION

Catalytic Direct Cross-Coupling of Organolithium Compounds with Aryl chlorides

Valentín Hornillos, Massimo Giannerini, Carlos Vila, Martín Fañanás-Mastral* and Ben L. Feringa*

Correspondence to: b.l.feringa@rug.nl, m.fananas.mastral@rug.nl

Stratingh Institute for Chemistry, University of Groningen, Nijenborgh 4, 9747 AG,
Groningen, The Netherlands.

Table of Contents

Table of Contents	S2
General Methods	S3
Additional Data	S3
Table S1. Additional optimization data of the reaction between 1a and PhLi	S3
Table S2. Additional optimization data of the reaction between 1m and PhLi	S4
Scheme S1. Scope limitations with deactivated aryl chlorides	S4
General Procedures for the Cross-Coupling of Aryllithium Reagents	S5
Preparation of Organolithium Reagents	S5
Data of Compounds 2a-2s, 10	S6
¹H and ¹³C NMR spectra	S13


General methods:

All reactions were carried out under a nitrogen atmosphere using oven dried glassware and using standard Schlenk techniques. THF and toluene were dried and distilled over sodium. Pd₂(dba)₃, SPhos, XPhos, DavePhos, CPhos, Qphos, P(^tBu)₃, PCy₃ and Pd-PEPPSI-Ipent were purchased from Aldrich and used without further purification. ⁿBuLi (1.6 M solution in hexane) and PhLi (1.8 M solution in dibutylether) were purchased from Acros. ThienylLi (1.0 M in THF/hexane), ^{tert}BuLi (1.7 M in pentane) and the compounds used as precursor for the preparation of lithium reagents, namely furan, 1-bromo-2,3-dimethyl-benzene, 1-bromo-2-methoxybenzene and 1-bromo-2-(methoxymethoxy)benzene were purchased from Aldrich. All the chlorides were commercially available and were purchased from TCI Europe N.V. with the exception of 1-chloro-4-methoxybenzene (Aldrich). Organolithium reagents other than the aforementioned were prepared according to described procedures (see below).

Chromatography: Merck silica gel type 9385 230-400 mesh, TLC: Merck silica gel 60, 0.25 mm. Components were visualized by UV and cerium/molybdenum or potassium permanganate staining. Progress and conversion of the reaction were determined by GC-MS (GC, HP6890: MS HP5973) with an HP1 or HP5 column (Agilent Technologies, Palo Alto, CA). Mass spectra were recorded on an AEI-MS-902 mass spectrometer (EI+) or a LTQ Orbitrap XL (ESI+). ¹H- and ¹³C-NMR were recorded on a Varian AMX400 (400 and 100.59 MHz, respectively) using CDCl₃ as solvent. Chemical shift values are reported in ppm with the solvent resonance as the internal standard (CHCl₃: δ 7.26 for ¹H, δ 77.0 for ¹³C). Data are reported as follows: chemical shifts, multiplicity (s = singlet, d = doublet, t = triplet, q = quartet, br = broad, m = multiplet), coupling constants (Hz), and integration. Carbon assignments are based on APT ¹³C-NMR experiments.

Optimization Study: Additional Data


Table S1. Additional optimization data of the reaction between 1a and PhLi



entry ^a	[Pd]	Ligand	Conv. (%)	2a:3:4:5 ^b
1	Pd ₂ (dba) ₃	L4.SPhos	55	51:0:0:4
2	Pd ₂ (dba) ₃	L5.DavePhos	58	48:0:0:11
3	Pd ₂ (dba) ₃	L6.CPhos	79	68:0:0:11
4	Pd ₂ (dba) ₃	L7.QPhos	42	32:0:0:10

^aConditions: PhLi (0.45 mmol, 1.8 M solution in dibutyl ether diluted with THF to a final concentration of 0.6 M) was added to a solution of 2-chloronaphthalene (0.3 mmol) in toluene (2 mL unless otherwise noted). ^b2a:3:4:5 ratios determined by GC analysis. ^c7.5 mol% was used. ^dIn 1 mL of toluene. dba = dibenzylideneacetone.


Table S2. Additional optimization data of the reaction between 1m and PhLi



entry ^a	[Pd]	T °C	PhLi flow-rate (mL/h)	Conv. (%)	2m:6:7 ^b
1	Pd₂(dba)₃/L3	50	1	76	70:<1:6
2	Pd₂(dba)₃/L3	50	0.33	Full	88:<1:12
3	Pd₂(dba)₃/L3	35	0.33	64	61:<1:<3
4	Pd₂(dba)₃/L3	40	0.33	82	77:<1:5
5	Pd₂(dba)₃/L3	45	0.33	80	70:4:6
6 ^c	Pd₂(dba)₃/L3	40	0.33	96	86:2:8
7 ^{c,d}	Pd₂(dba)₃/L3	40	0.2	Full	89:3:8
8 ^c	Pd-PEPPSI-IPent	40	0.2	Full	90:4:6
9 ^c	Pd-PEPPSI-IPent	30	0.2	90	81:5:4

^aConditions: Phenyllithium (1.8 M solution in dibutyl ether diluted with THF to a final concentration of 0.6M) was added to a solution of 1-butyl-4-chlorobenzene (0.3 mmol) in toluene (2 mL unless otherwise noted). ^b2m:6:7 ratios determined by GC-analysis. ^cIn 1 mL of toluene. ^dPhenyllithium was diluted with toluene (instead of THF) to reach 0.6 M concentration. dba, dibenzylideneacetone.

Scheme S1. Pd-catalyzed cross-coupling of aryl lithium reagents with deactivated aryl chlorides: Scope Limitations



The Pd-catalyzed cross-coupling with aryl lithium reagents did not proceed efficiently when very deactivated aryl chlorides were used. The reaction between 4-(4-chlorophenyl)morpholine **1t** and PhLi led to low conversion and a mixture of products while the use of 6-chloroindole **1u** gave no reaction. Although it was used efficiently in the Pd-catalyzed cross-coupling with activated aryl chlorides, the less reactive thienyl lithium could not be coupled with deactivated and hindered aryl chlorides.

General procedures for the cross-coupling of aryl lithium reagents with aryl chlorides

Catalytic system A:

In a dry Schlenk flask Pd-PEPPSI-IPent (5 mol%, 0.015 mmol, 11.9 mg) and the substrate (0.3 mmol) were dissolved in 1 mL of dry toluene and the solution was stirred at the indicated temperature. The corresponding lithium reagent (1.5 equiv) was diluted with THF to reach the concentration of 0.6 M; this solution was slowly added with the indicated flow rate by the use of a syringe pump. After the addition was completed a saturated aqueous solution of NH₄Cl was added and the mixture was extracted with ether (3 x 5 mL). The organic phases were combined and dried with anhydrous Na₂SO₄. Evaporation of the solvent under reduced pressure afforded the crude product that was then purified by column chromatography.

Note: Thienyllithium (1.5 equiv) was diluted with toluene to reach a concentration of 0.6 M and TMEDA (1.5 equiv) was added. The resulting solution was slowly added with a flow rate of 0.5 mL/h to the reaction mixture warmed at 40 °C.

Catalytic system B:

In a dry Schlenk flask Pd₂(dba)₃ (2.5 mol%, 0.0075 mmol, 6.87 mg) and XPhos (10 mol%, 0.03 mmol, 14.3 mg) were dissolved in toluene, the substrate (0.3 mmol) was added and the solution stirred at the indicated temperature. The corresponding lithium reagent (1.5 equiv) was diluted with THF to reach a concentration of 0.6 M; this solution was slowly added with the indicated flow rate by the use of a syringe pump. After the addition was completed a saturated solution of aqueous NH₄Cl was added and the mixture was extracted with ether (3 x 5 mL). The organic phases were combined and dried with anhydrous Na₂SO₄. Evaporation of the solvent under reduced pressure afforded the crude product that was then purified by column chromatography.

Preparation of organolithium reagents:

A. 2,3-Dimethyl-phenyllithium and 2-Methoxy-phenyllithium

In a dry Schlenk flask the corresponding aryl bromide (1.8 mmol) was dissolved in dry THF (0.9 mL) and the solution was cooled down to -78 °C. ^tBuLi (2 equiv) was added slowly and the solution was stirred for 1 h. Then the solution was allowed to reach room temperature.

B. Furyllithium¹

Furan (9.0 mmol, 612.6 mg, 654.5 μl) was dissolved in THF (4.5 mL) and the solution was cooled down to -40 °C. ⁿBuLi (8.5 mmol) was added slowly. Then the solution was allowed to reach room temperature, stirred for 3 h and diluted with 4.4 mL of THF to reach a final concentration of 0.6M.

¹ Raczko, J.; Golebiowski, A.; Krajewski, J. W.; Gluzinski, P.; Jurczak, J. *Tetrahedron Lett.* **1990**, *31*, 3797.

C. 2-Methoxymethoxy-phenyllithium²


In a dry Schlenk flask (methoxymethoxy)benzene (1.0 mmol, 138 mg) was dissolved in dry THF (3 mL) and the solution was cooled down to -78 °C. ^tBuLi (1 equiv) was added slowly and the solution was stirred for 1 h. Then the solution was allowed to reach room temperature.

D. 2-Methoxy-1-naphthyllithium

In a dry Schlenk flask 1-bromo-2-methoxynaphthalene (1.8 mmol, 427 mg) was dissolved in dry THF (0.9 mL) and slowly added (flow rate: 8 ml/h) to a solution of ⁿBuLi (1.6 M in hexane, 1.05 equiv.) at -10 °C. The solution was then stirred for 30 min. and allowed to reach room temperature. The solution was diluted with 1 ml of toluene to reach a final concentration of 0.6M.

Data of Compounds 2a-2s, 10

Physical data for known compounds were identical in all respects to those previously reported (references are given).




2-Phenylnaphthalene (2a):³

CAS Registry Number: 612-94-2.

Synthesized using catalytic systems A and B with 2-chloronaphthalene (0.3 mmol, 49 mg) and 225 μ L of PhLi (1.8 M solution in dibutylether) diluted with 500 μ L of THF.

Catalytic system A: Reaction carried out at room temperature. Flow rate of PhLi solution = 0.5 mL/h. White solid obtained after column chromatography (SiO₂, *n*-pentane/EtOAc 100:1), 54 mg, 88% yield.

Catalytic system B: Reaction carried out at room temperature in 2 mL of toluene. Flow rate of PhLi solution = 1 mL/h. White solid obtained after column chromatography, 57 mg, 93% yield.



1-Phenylnaphthalene (2b):⁴

CAS Registry Number: 605-02-7

Synthesized using catalytic systems A and B with 1-chloronaphthalene (0.3 mmol, 49 mg) and 225 μ L of PhLi (1.8 M solution in dibutylether) diluted with 500 μ L of THF.

² Vo, C. T.; Mitchell, T. A.; Bode, J. W. *J. Am. Chem. Soc.* **2011**, *133*, 14082.


³ Giannerini, M.; Fañanás-Mastral, M.; Feringa, B. L. *Nature Chem.* **2013**, doi:10.1038/nchem.1678.

⁴ (a) Katakoa, N.; Shelby, Q.; Stambuli, J.P.; Hartwig, J.F. *J. Org. Chem.* **2002**, *67*, 5553.

Catalytic system A: Reaction carried out at room temperature. Flow rate of PhLi solution = 0.5 mL/h. White solid obtained after column chromatography (SiO₂, *n*-pentane/EtOAc 100:1), 56 mg, 91% yield.

Catalytic system B: Reaction carried out at room temperature in 2 mL of toluene. Flow rate of PhLi solution = 1 mL/h. White solid obtained after column chromatography, 60 mg, 98% yield.

The reaction performed with 10 mmol (1.63 g) of substrate in the presence of 1 mol% of Pd₂dba₃ and 4 mol% of **L3** afforded product **2b** in 92% yield. Toluene = 30 mL. PhLi (8.3 mL, 1.8 M solution in dibutylether) diluted with 16.6 mL of THF. Flow rate of PhLi solution = 3 mL/h.




4-(Trifluoromethyl)-1,1'-biphenyl (**2c**):⁵

CAS Registry Number: 398-36-7

Synthesized using catalytic systems A and B with 1-chloro-4-(trifluoromethyl)benzene (0.3 mmol, 54 mg) and 225 μL of PhLi (1.8 M solution in dibutylether) diluted with 500 μL of THF.

Catalytic system A: Reaction carried out at room temperature. Flow rate of PhLi solution = 0.5 mL/h. White solid obtained after column chromatography (SiO₂, *n*-pentane/Et₂O 100:1), 59 mg, 89% yield.


Catalytic system B: Reaction carried out at room temperature in 2 mL of toluene. Flow rate of PhLi solution = 1 mL/h. White solid obtained after column chromatography, 63 mg, 95% yield.



2-(Naphthalen-2-yl)thiophene (**2d**):⁶

CAS Registry Number: 16939-09-6

Synthesized using catalytic systems A with 2-chloronaphthalene (0.3 mmol, 49 mg) and 750 μL of thienyllithium (0.6M). White solid obtained after column chromatography (SiO₂, *n*-pentane/ EtOAc 100:1), 52 mg, 82% yield.



2-(Naphthalen-1-yl)thiophene (**2e**):⁷


⁵ Guha, N. R.; Reddy, C. B.; Aggarwal, N.; Sharma, D.; Shil, A. K.; Das, B. P. *Adv. Synth. Catal.* **2012**, 354, 2911.

⁶ Bolliger, J. L.; Frech, C. M. *Adv. Synth. Cat.* **2010**, 352, 1075.

⁷ Cahiez, G.; Moyeux, A.; Gager, O.; Poizat, M. *Adv. Synth. Cat.* **2013**, 355, 790

CAS Registry Number: 4632-51-3.


Synthesized using catalytic systems A with 1-chloronaphthalene (0.3 mmol, 49 mg) and 750 μL of thienyllithium (0.6M). Colorless oil obtained after column chromatography (SiO_2 , *n*-pentane/ EtOAc 100:1), 55 mg, 87% yield.



2-(4-(Trifluoromethyl)phenyl)thiophene (2f):⁸

CAS Registry Number: 115933-15-8.


Synthesized using catalytic systems A with 1-chloro-4-(trifluoromethyl)benzene (0.3 mmol, 54 mg) and 750 μL of thienyllithium (0.6M). White solid obtained after column chromatography (SiO_2 , *n*-pentane/ EtOAc 100:1), 57 mg, 83% yield.



2-(2,3-Dimethylphenyl)naphthalene (2g):⁹

CAS Registry Number: 382151-52-2.

Synthesized using catalytic systems B with 2-chloronaphthalene (0.3 mmol, 49 mg) and 725 μL of 2,3-dimethyl-phenyllithium (0.6M). Reaction carried out at room temperature in 2 mL of toluene. Flow rate of 2,3-dimethyl-phenyllithium solution = 1 mL/h. White solid obtained after column chromatography (SiO_2 , *n*-pentane/ EtOAc 100:1), 62 mg, 89% yield.



1-(2-Methoxyphenyl)naphthalene (2h):¹⁰


CAS Registry Number: 93321-11-0.

Synthesized using catalytic systems A with 1-chloronaphthalene (0.3 mmol, 49 mg) and 725 μL of 2-methoxy-phenyllithium (0.6M). Reaction carried out at room temperature. Flow rate of PhLi solution = 0.5 mL/h. White solid obtained after column chromatography (SiO_2 , *n*-pentane/ EtOAc 100:1), 59 mg, 84% yield.

⁸ Shi, S.; Zhang, Y.; *Green Chem.* **2008**, *10*, 868.

⁹ Yonehara, F.; Kido, Y.; Sugimoto, H.; Morita, S.; Yamaguchi, M. *J. Org. Chem.* **2003**, *68*, 6752.


¹⁰ Song, C.; Ma, Y.; Chai, Q.; Ma, C.; Jiang, W.; Andrus, M. B. *Tetrahedron* **2005**, *61*, 7438–7446.



2-Methoxy-1-(4-(trifluoromethyl)phenyl)naphthalene (2i):¹¹

CAS Registry Number: 922511-77-1.


Synthesized using catalytic systems A with 1-chloro-4-(trifluoromethyl)benzene (0.3 mmol, 54 mg) and 725 μ L of 2-methoxy-1-naphthyllithium (0.6M). Reaction carried out at room temperature. Flow rate of PhLi solution = 0.5 mL/h. White solid obtained after column chromatography (SiO_2 , *n*-pentane/ EtOAc 100:1), 70 mg, 77% yield.



1-(2-(methoxymethoxy)phenyl)naphthalene (2j):¹²

CAS Registry Number: 141362-07-4.


Synthesized using catalytic systems A with 1-chloronaphthalene (0.3 mmol, 49 mg) and 1500 μ L of 2-Methoxy-phenyllithium (0.3M). Reaction carried out at room temperature. Flow rate of PhLi solution = 0.5 mL/h. Waxy solid obtained after column chromatography (SiO_2 , *n*-pentane/ EtOAc 100:2), 66 mg, 83% yield.



2-(Naphthalen-1-yl)furan (2k):¹³

CAS Registry Number: 51792-32-6.

Synthesized using catalytic systems A with 1-chloronaphthalene (0.3 mmol, 49 mg) and 725 μ L of furyllithium (0.6M). Pale yellow oil obtained after column chromatography (SiO_2 , *n*-pentane/ EtOAc 100:1), 56 mg, 96% yield.



2-(4-(Trifluoromethyl)phenyl)furan (2l):¹⁴


¹¹ Li, H.; Wu, Y.; Yan, W. *J. Organomet. Chem.* **2006**, *691*, 5688.

¹² Wawrzyniak, P.; Heinicke, J. *Tetrahedron Lett.* **2006**, *47*, 8921.

¹³ Molander, G. A.; Beaumard, F. *Org. Lett.* **2010**, *12*, 4022.

CAS Registry Number: 214463-10-2.

Synthesized using catalytic systems A with 1-chloro-4-(trifluoromethyl)benzene (0.3 mmol, 54 mg) and 725 μL of furyllithium (0.6M). White solid obtained after column chromatography (SiO_2 , *n*-pentane/ EtOAc 100:1), 49 mg, 77% yield.




4-Butyl-1,1'-biphenyl (2m):¹⁵

CAS Registry Number: 37909-95-8.

Synthesized using catalytic systems A and B with 1-butyl-4-chlorobenzene (0.3 mmol, 51 mg) and 225 μL of PhLi (1.8 M solution in dibutylether) diluted with 500 μL of THF.

Catalytic system A: Reaction carried out at 35°C. Flow rate of PhLi solution = 0.2 mL/h. White solid obtained after column chromatography (SiO_2 , *n*-pentane/ EtOAc 100:1), 54 mg, 86% yield.

Catalytic system B: Reaction carried out at 40°C in 1 mL of toluene. Flow rate of PhLi solution = 0.2 mL/h. Colorless oil obtained after column chromatography, 53 mg, 84% yield.




2,4-Dimethyl-1,1'-biphenyl (2n):¹⁶

CAS Registry Number: 4433-10-7.

Synthesized using catalytic systems A and B with 1-chloro-2,4-dimethylbenzene (0.3 mmol, 42 mg) and 225 μL of PhLi (1.8 M solution in dibutylether) diluted with 500 μL of THF. **Catalytic system A:** Reaction carried out at 35°C. Flow rate of PhLi solution = 0.2 mL/h. White solid obtained after column chromatography (SiO_2 , *n*-pentane/ EtOAc 100:1), 44 mg, 80% yield.

Catalytic system B: Reaction carried out at 40°C in 1 mL of toluene. Flow rate of PhLi solution = 0.2 mL/h. Colorless oil obtained after column chromatography, 45 mg, 82% yield.



4-Methoxybiphenyl (2o):³

CAS Registry Number: 613-37-6.

Synthesized using catalytic systems A and B with 1-chloro-4-methoxybenzene (0.3 mmol, 43 mg) and 225 μL of PhLi (1.8 M solution in dibutylether) diluted with 500 μL of THF. **Catalytic system A:** Reaction carried out at 40°C. Flow rate of PhLi solution =


¹⁴ Denmark, S. E.; Baird, J. D. *Org. Lett.* **2006**, *8*, 793.

¹⁵ Littke, A. F.; Schwarz, L.; Fu, G. C. *J. Am. Chem. Soc.* **2002**, *124*, 6343.

¹⁶ Zim, D.; Gruber, A. S.; Ebeling, G.; Dupont, J.; Monteiro, A. L. *Org. Lett.* **2000**, *2*, 2881.

0.2 mL/h. White solid obtained after column chromatography (SiO₂, *n*-pentane/ EtOAc 100:1), 52 mg, 94% yield.


Catalytic system B: Reaction carried out at 40°C in 1 mL of toluene. Flow rate of PhLi solution = 0.2 mL/h. Colorless oil obtained after column chromatography, 34 mg, 62% yield.



3-Methoxy-1,1'-biphenyl (2p).¹⁷

CAS Registry Number: 2113-56-6.


Synthesized using catalytic systems A with 1-chloro-3-methoxybenzene (0.3 mmol, 43 mg) and 225 μ L of PhLi (1.8 M solution in dibutylether) diluted with 500 μ L of THF. Reaction carried out at 40°C. Flow rate of PhLi solution = 0.2 mL/h. Colorless oil obtained after column chromatography (SiO₂, *n*-pentane/ EtOAc 100:1), 42 mg, 76% yield.



2',5'-Dimethyl-*p*-terphenyl (2q).¹⁸

CAS Registry Number: 20260-22-4.

Synthesized using catalytic systems A with 1-chloro-4-methoxybenzene (0.3 mmol, 53 mg) and 500 μ L of PhLi (1.8 M solution in dibutylether) diluted with 1.5 mL of THF. Reaction carried out at 40°C. Flow rate of PhLi solution = 0.2 mL/h. Colorless oil obtained after column chromatography (SiO₂, *n*-pentane/ EtOAc 100:1), 62 mg, 80% yield.




2'-Methoxy-2,4-dimethyl-1,1'-biphenyl (2r).

Synthesized using catalytic systems A with 2,4-dimethylbenzene (0.3 mmol, 42 mg) and 725 μ L of 2-methoxy-phenyllithium (0.6M). Reaction carried out at 35°C. Flow rate of PhLi solution = 0.2 mL/h. Colorless oil obtained after column chromatography (SiO₂, *n*-pentane/ EtOAc 100:1), 55 mg, 86% yield. ¹H NMR (400 MHz, CDCl₃) δ 7.38 (td, *J* = 8.2, 1.8 Hz, 1H), 7.99 – 7.20 (m, 6H), 3.81 (s, 3H), 2.41 (s, 3H), 2.16 (s, 3H) ppm. ¹³C NMR (100 MHz, CDCl₃) δ 156.7, 136.9, 136.6, 135.7, 131.2, 130.8, 130.5, 130.0, 128.4, 126.3, 120.4, 110.6, 55.4, 21.2, 19.9 ppm. EI-MS *m/z* (%) 212 (100).

¹⁷ Mino, T.; Shirae, Y.; Sakamoto, M.; Fujita, T. *J. Org. Chem.* **2005**, *70*, 2191.


¹⁸ K.-Nezhad, A.; Panahi, F. *J. Organomet. Chem.* **2012**, *717*, 146.



2-(4-Methoxyphenyl)furan (2s):¹⁹

CAS Registry Number: 17113-31-4.

Synthesized using catalytic systems A with 1-chloro-4-methoxybenzene (0.3 mmol, 43 mg) and 725 μ L of furyllithium (0.6M). White solid obtained after column chromatography (SiO₂, *n*-pentane/ EtOAc 100:2), 51 mg, 80% yield.





2-([1,1'-Biphenyl]-4-yl)furan (10):²⁰



Synthesized using catalytic systems A with **9** (0.3 mmol, 57 mg) and 725 μ L of furyllithium (0.6M). White solid obtained after column chromatography (SiO₂, *n*-pentane), 59 mg, 90% yield. ¹H NMR (400 MHz, CDCl₃) δ 7.78 (d, *J* = 8.4 Hz, 2H), 7.69 – 7.62 (m, 4H), 7.52 (d, *J* = 1.7 Hz, 1H), 7.48 (t, *J* = 7.6 Hz, 2H), 7.38 (t, *J* = 7.3 Hz, 1H), 6.72 (d, *J* = 3.3 Hz, 1H), 6.52 (dd, *J* = 3.3, 1.7, Hz 1H) ppm. ¹³C NMR (101 MHz, CDCl₃) δ 153.8, 142.2, 140.6, 140.0, 129.9, 128.8, 127.4, 126.9, 124.2, 111.8, 105.2 ppm. HRMS (ESI+, *m/z*): calcd for C₁₆H₁₃O [M+H]⁺: 221.09609; found: 221.09511.



¹⁹ Naber, J. R.; Buchwald, S. L. *Adv. Synth. Cat.*, **2008**, 350, 957.



²⁰ Haner, J.; Jack, K.; Nagireddy, J.; Raheem, M. A.; Durham, R.; Tam, W. *Synthesis*, **2011**, 731.



^1H and ^{13}C NMR spectra



























^{19}F NMR

