Binding of hemopressin peptide with cannabinoid CB_1 receptor: structural insights

Mario Scrima,¹ Sara Di Marino,¹ Manuela Grimaldi,¹ Antonia Mastrogiacomo¹ Ettore Novellino², Maurizio Bifulco,¹ Anna Maria D'Ursi*¹

Contents

Binding of	of hemopressin peptide with cannabinoid CB ₁ receptor:	1
structural	insights	1
1 Supp	plementary results	2
1.1	Proton Chemical Shifts of Hemopressin in DPC/SDS micelles 90:10 M/M.	2
1.2	Proton Chemical Shifts of Hemopressin(1-6) in DPC/SDS micelles 90:10 M/M.	2
1.3	Plots of relative NOE cross peak intensities, versus the mixing time τ. NOE cross peak	
	intensities involving NH protons of Val, Lys, Asn, Phe, were respectively used to produc	ce
	the buildup curve.	3
1.4	Structural statistics for the final 20 structures of Hemopressin and Hemopressin(1-6) in	
	DPC/SDS micelles	4
1.5	Amide and fingerprint regions of NOESY spectra for Hemopressin and	
	Hemopressin(1-6) in DPC/SDS 90:10 M/M	5
1.6	Ramachandran Plots for the CB1 final structure from MODELLER calculation.	6
1.7	Molecular interactions of Rimonabant, Hemopressin and Hemopressin(1-6) into CB1	
	receptor and calculated values of inhibition constant.	7

¹Department of Pharmaceutical Sciences, University of Salerno, I-84084 Fisciano, Italy.

²Department of Pharmaceutical and Toxicological Chemistry, University of Naples Federico II,

^{*} Corresponding author

1 Supplementary results

1.1 Proton Chemical Shifts of Hemopressin in DPC/SDS micelles 90:10 M/M.

Table 1S. The 1 H chemical shifts of Hemopressin in water DPC/SDS 90:10 M/M and HFA/H $_{2}$ O at 300K and 600 MHz

Hemopressin in DPC/SDS

Tiemopressii iii Di C/5D5						
HN	$C^{\alpha}H$	$C^{\beta}H$	$\mathbf{C}^{\gamma}\mathbf{H}$	$C^{\delta}H$	C ^E H	Others
	4.368	2.397	1.891	3.290		
8.788	3.918	1.856	0.796/0.640			
8.323	4.618	2.724/2.620		7.594/6.780		
8.143	4.477	2.936/2.718		7.125	7.201	HZ7.040
8.397	4.078	1.728	1.283	1.562	2.883	QZ7.791
7.785	4.401	3.101/3.032		7.209	7.047	HZ7.140
7.456	4.219	1.539/1.474	1.1755	0.791/0.755		QQD0.780
7.814	4.307	3.749				
7.990	4.382	3.166/3.040		7.990	7.173	QB3.166
	8.788 8.323 8.143 8.397 7.785 7.456 7.814	4.368 8.788 3.918 8.323 4.618 8.143 4.477 8.397 4.078 7.785 4.401 7.456 4.219 7.814 4.307	HN $C^{\alpha}H$ $C^{\beta}H$ 4.3682.3978.7883.9181.8568.3234.6182.724/2.6208.1434.4772.936/2.7188.3974.0781.7287.7854.4013.101/3.0327.4564.2191.539/1.4747.8144.3073.749	HN C ^α H C ^β H C ^γ H 4.368 2.397 1.891 8.788 3.918 1.856 0.796/0.640 8.323 4.618 2.724/2.620 8.143 4.477 2.936/2.718 8.397 4.078 1.728 1.283 7.785 4.401 3.101/3.032 1.1755 7.814 4.307 3.749 1.1755	HN $C^{\alpha}H$ $C^{\beta}H$ $C^{\gamma}H$ $C^{\delta}H$ 4.3682.3971.8913.2908.7883.9181.8560.796/0.6408.3234.6182.724/2.6207.594/6.7808.1434.4772.936/2.7187.1258.3974.0781.7281.2831.5627.7854.4013.101/3.0327.2097.4564.2191.539/1.4741.17550.791/0.7557.8144.3073.749	HN $C^{\alpha}H$ $C^{\beta}H$ $C^{\gamma}H$ $C^{\delta}H$ $C^{\epsilon}H$ 4.3682.3971.8913.2908.7883.9181.8560.796/0.6408.3234.6182.724/2.6207.594/6.7808.1434.4772.936/2.7187.1257.2018.3974.0781.7281.2831.5622.8837.7854.4013.101/3.0327.2097.0477.4564.2191.539/1.4741.17550.791/0.7557.8144.3073.749


1.2 Proton Chemical Shifts of Hemopressin(1-6) in DPC/SDS micelles 90:10 M/M.


Table 2S. The ¹H chemical shifts of Hemopressin(1-6) in water DPC/SDS 90:10 M/M and HFA/H₂O at 300K and 600 MHz

Emopressina 6aa in DPC/SDS

Residue	HN	CαH	$\hat{\mathbf{C}}^{\beta}\mathbf{H}$	$\mathbf{C}^{\gamma}\mathbf{H}$	$C^{\delta}H$	CεH	Others
Pro1		4.400	2.425	1.965	3.345		
Val2		4.036	1.958	0.867/0.796			
Asn3	8.472	4.650	2.725/2.638		6.807/7.550		
Phe4	8.055	4.494	2.945/2.871		7.105	7.176	
Lys5	7.910	4.245	1.696	1.256	1.577	2.912	
Phe6	7.691	4.349	3.120/2.902		7.234		

1.3 Plots of relative NOE cross peak intensities, versus the mixing time τ . NOE cross peak intensities involving NH protons of Val, Lys, Asn, Phe, were respectively used to produce the buildup curve.






1.4 Structural statistics for the final 20 structures of Hemopressin and Hemopressin(1-6) in DPC/SDS micelles


Table 3S. Structural statistics for the final 20 structures of Hemopressin and Hemopressin(1-6) in DPC/SDS micelles

PARAMETER	Hemopressin DPS/SDS	Hemopressin(1-6) DPC/SDS		
Backbone RMSD from	Peptide region			
mean structure (Å)	2-8	2-6		
mean structure (A)	0.164	0.1991		
NOE distance				
restraints				
total	221	96		
Intra	102	57		
Short	66	25		
Medium	53	14		
Long	0	0		
Ramachandran Plot				
Residues in most	55.8%	20.3%		
favoured regions		20.3 //		
Residues in additional	7.5%	76.6%		
allowed regions	7.570	7 3.0 70		
Residues in generously	36.7%	1.6%		
allowed regions				

1.5 Amide and fingerprint regions of NOESY spectra for Hemopressin and Hemopressin(1-6) in DPC/SDS 90:10 M/M



1.6 Ramachandran Plots for the CB1 final structure from MODELLER calculation.



1.7 Molecular interactions of Rimonabant, Hemopressin and Hemopressin(1-6) into CB1 receptor and calculated values of inhibition constant.

Table 4S. Molecular interactions of Rimonabant, Hemopressin and Hemopressin(1-6) into CB1 receptor and calculated values of inhibition constant

vdW interaction residues (scaling factor = 1.50 Å)					
RIMONABANT					
Ligand	Ligand Protein				
phenyl portions	Phe200, Tyr275, Trp279, Trp356				
Pyrazole group	Val196, Met363, Leu387, Tyr275				
Amidic portion	Lys192,Asp366,Ser383,Tyr275				
piperidinyl moiety	Phe379, Phe177, Ile375, Leu387				
	HEMOPRESSI	N			
Pro1	,Asp176, Phe	180, Ile375, Arg182, I	His181		
Val2	Ser1	85,Arg182, His181			
Asn3	Phe189,	Lys192,Ser185, Arg1	82		
Phe4	Phe189 , Ser1	85, Leu190, Arg182,	<i>Trp255</i>		
Lys5	Asp272, Phe189, Met3	363, Tyr275, Asp366, I	Lys192, Leu193		
Phe6	Asp272, Val363, Phe368, Lys370, Gly369, Asp366				
Leu7	Leu7				
Ser8	<i>Trp255</i>				
His9	Asp272, Glu273, Thr274, I	Leu193, Phe189, Asn2	256, Tyr275, Trp255		
	HEMOPRESSIN(1-6)			
Pro1	Phe180, Ile3	75, Phe177, His181,A	sp176		
Val2	Ile375,Asp17	76, His181, Arg182, P.	he180		
Asn3	Asp176,Lys1	92, Ile375, Phe177, P.	he379		
Phe4					
Lys5	Met363, Leu193, Tyr2	275, Phe189, Asp366, 1	Lys192,Phe379		
Phe6	Gly369, Asp2	72, Val367, Phe368, I	Lys370,		
Values of inhibition constant ^a and energies ^b					
	LIGAND				
	Rimonabant	Hemopressin	Hemopressin(1-6)		
Inhibition Constant, Ki	20.84pM	1.64mM	1.44uM		
Free Energy of Binding	-14.57	-3.80	-7.97		
Torsional Free Energy	1.19	10.44	6.86		
^a the inhibition constant is measured at 298.15 K.					
^b energies are expressed in Kcal/mol.					