

Phenoxazinones Revisited – Structural Merger with an Epoxyquinone Antibiotic

Patrícia Bezerra Gomes,^a Markus Nett,^a Hans-Martin Dahse,^a and Christian Hertweck^{*a,b}

^a Leibniz Institute for Natural Product Research and Infection Biology, HKI, Beutenbergstr. 11a, D-07745 Jena, Germany. Fax: + 49 3641 5320804; Tel: + 49 3641 5321100; E-mail:

Christian.Hertweck@hki-jena.de

^b Friedrich Schiller University, Jena, Germany




Figure S1. ¹H NMR (500 MHz, DMSO-*d*₆) spectrum of **1**




Figure S2. ^{13}C NMR (125 MHz, $\text{DMSO-}d_6$) spectrum of **1**




Figure S3. HSQC (500 MHz, $\text{DMSO-}d_6$) spectrum of **1**




Figure S4. ^1H - ^1H COSY (500 MHz, $\text{DMSO}-d_6$) spectrum of **1**




Figure S5. HMBC (500 MHz, $\text{DMSO}-d_6$) spectrum of **1**




Figure S6. ^1H NMR (600 MHz, $\text{DMSO-}d_6$) spectrum of **2**




Figure S7. ^{13}C NMR (150 MHz, $\text{DMSO-}d_6$) spectrum of **2**




Figure S8. HSQC (600 MHz, DMSO- d_6) spectrum of **2**




Figure S9. ^1H - ^1H COSY (600 MHz, DMSO- d_6) spectrum of **2**




Figure S10. HMBC (600 MHz, DMSO-*d*₆) spectrum of **2**




Figure S11. NOESY (600 MHz, DMSO-*d*₆) spectrum of **2**