

Supporting Information

Synthesis and antiproliferative evaluation of certain indeno[1,2-c]quinoline derivatives. Part 2

**Chih-Hua Tseng,^a Cherng-Chyi Tzeng,^a Chiao-Li Yang,^a Pei-Jung Lu,^b Hui-Ling Chen,^b
Hao-Yi Li,^c You-Chung Chuang,^c Chia-Ning Yang,^{*,c} and Yeh-Long Chen,^{*,a}**

^a*Department of Medicinal and Applied Chemistry, College of Life Science, Kaohsiung Medical University, Kaohsiung City 807, Taiwan*

^b*Institute of Clinical Medicine, National Cheng-Kung University, School of Medicine, 138 Sheng-Li Road, Tainan 704, Taiwan*

^c*Institute of Biotechnology, National University of Kaohsiung, 700 Kaohsiung University Rd, Kaohsiung, Taiwan.*

Contents:

S1: Title Page

S2: Detailed parameters of **5a** was determined by the X-ray crystallography

Detailed parameters of 5a was determined by the X-ray crystallography.

Compound **5a** was unambiguously confirmed by its X-ray structural analysis. The structure was solved and refined by direct methods Shelx 97^{ref} suite of programs. Red single crystal (0.25 x 0.20 x 0.15 mm³) of **5a** was obtained by slow evaporation from ethanol/CH₂Cl₂ (30/70) solution: monoclinic, space group P 2(1)/n, $a = 14.5476(3) \text{ \AA}$, $b = 8.4212(2) \text{ \AA}$, $c = 14.9701(3) \text{ \AA}$, $\alpha = 90.00^\circ$, $\beta = 103.7400(10)^\circ$, $\gamma = 90.00^\circ$, $V = 1781.48(7) \text{ \AA}^3$, $Z = 4$, δ (calcd) = 1.295 Mg.m⁻³, FW = 347.41 for C₂₁H₂₁N₃O₂, $F(000) = 736$. Complete crystallographic data for the structural analysis have been deposited with the Cambridge Crystallographic Data Centre, CCDC no.760627 for compound **5a**. Copies of this information may be obtained free of charge from the Director, Cambridge Crystallographic Data Centre, 12 Union Road, Cambridge, CB2 1EZ, UK. (fax: +44-1223-336033, e-mail: deposit@ccdc.cam.ac.uk or via www.ccdc.cam.ac.uk).

(Ref: Sheldrick, G. M., SHELXS 97, Program Crystal Structure Solution, **1997**, University of Göttingen, Germany.)