

Supplemental Table 5. Adjusted hazard ratios (95% CI) of early menopause (menopause <45 yrs) vs. normal menopause (menopause ≥45 yrs) and cardiovascular disease risk in women from ‘Other’ ethnicities

	CHD	Stroke	ASCVD	HF
All	0.996 (0.848 1.169)	1.000 (0.852 1.172)	1.008 (0.857 1.187)	0.979 (0.834 1.149)
<i>P-interaction</i>	0.07	0.23	0.12	0.25
DM	1.205 (0.800 1.815)	1.150 (0.777 1.704)	1.217 (0.804 1.843)	1.109 (0.733 1.678)
Non-DM	0.943 (0.791 1.125)	0.972 (0.816 1.159)	0.961 (0.803 1.149)	0.958 (0.804 1.142)

T2DM: type 2 diabetes; White: non-Hispanic white; Black: non-Hispanic black. CHD: coronary heart disease; ASCVD: Atherosclerotic cardiovascular disease (CHD or stroke); HF: heart failure

HRs were adjusted for age, race, cohort indicator, education, smoking status, alcohol drinking status, BMI, mean blood pressure, mean total cholesterol, antihypertensive medication, lipid-lowering medication, ever used hormone therapy, pregnancy history, and diabetes.