

Supplemental Material S1. Apraxia measures for a subsample of 65 participants from Fridriksson et al. (2018).

Characteristic	Typical BDNF genotype (Val/Val) <i>n</i> = 36		Atypical BDNF genotype (Val/Met, Met/Met) <i>n</i> = 29		95% CI of the difference	Two-sided <i>p</i> -value
	<i>M</i>	<i>SD</i>	<i>M</i>	<i>SD</i>		
ASRS	2.89	2.00	2.90	2.06	(-1.02, 1.00)	.988
ABA limb apraxia	46.64	4.09	41.14	10.72	(-1.63, 9.37)	.006
ABA oral apraxia	43.31	7.77	37.93	9.75	(1.04, 9.71)	.016

Note. BDNF = brain-derived neurotrophic factor; CI = confidence interval; ASRS = Apraxia of Speech Rating Scale (Strand, Duffy, Clark, & Josephs, 2014); ABA = Apraxia Battery for Adults-2nd Edition (Dabul, 2000).

Reference

- Dabul, B. (2000). *Apraxia Battery for Adults-Second Edition (ABA-2)*. Austin, TX: Pro-Ed.
- Fridriksson, J., Elm, J., Stark, B. C., Basilakos, A., Rorden, C., Sen, S., . . . Bonilha, L. (2018). BDNF genotype and tDCS interaction in aphasia treatment. *Brain Stimulation, 11*(6), 1276-1281. <https://doi.org/10.1016/j.brs.2018.08.009>
- Strand, E. A., Duffy, J. R., Clark, H. M., & Josephs, K. (2014). The Apraxia of Speech Rating Scale: A tool for diagnosis and description of apraxia of speech. *Journal of Communication Disorders, 51*, 43-50. <https://doi.org/10.1016/j.jcomdis.2014.06.008>