

Supplemental Material S6. Mixed models for whispering, coughing, and throat clearing before and during voice rest.

Behavior	Effect	Estimate ± SE	<i>p</i> value
Whispering	Intercept	21.889 ± 4.131	< .0001*
	Group		.0118*
	Therapy	-14.117 ± 5.225	
	Counseling	-12.583 ± 5.386	
	Evaluation	Reference	
	Time		.6974
	Baseline	Reference	
Voice rest	-4.937 ± 5.616		
	Group × Time		.6759
Coughing	Intercept	46.222 ± 5.353	< .0001*
	Group		.0053*
	Therapy	-26.994 ± 6.772	
	Counseling	-16.883 ± 6.980	
	Evaluation	Reference	
	Time		< .0001*
	Baseline	Reference	
Voice rest	-16.080 ± 4.547		
	Group × Time		.0639
Throat clearing	Intercept	51.508 ± 5.380	< .0001*
	Group		.0175*
	Therapy	-24.794 ± 6.805	
	Counseling	-12.749 ± 7.064	
	Evaluation	Reference	
	Time		< .0001*
	Baseline	Reference	
Voice rest	-24.461 ± 5.819		
	Group × Time		.0746

**p* < .05 for Type 3 effect in mixed models with voice behavior as the response variable, Group, Time, and Group × Time interaction as fixed effects and participant as repeated measures.