

**Intra-genomic variability and pseudogenes in ribosomal
ITS regions of *Paraplagusia blochii* (Pleuronectiformes:
Cynoglossidae)**

**Li Gong^{1,2,*}, Xiaoyu Kong^{3,4,*}, Hairong Luo^{3,4}, Shixi Chen^{3,4}, Wei Shi^{3,4} and Min
Yang^{3,4}**

¹ National Engineering Laboratory of Marine Germplasm Resources Exploration and Utilization,
Zhejiang Ocean University, 316022, Zhoushan, China

² National Engineering Research Center for Facilitated Marine Aquaculture, Marine Science and
Technology College, Zhejiang Ocean University, 316022, Zhoushan, China

³ Key Laboratory of Tropical Marine Bio-Resources and Ecology, Guangdong Provincial Key
Laboratory of Applied Marine Biology, South China Sea Institute of Oceanology, Chinese Academy of
Sciences, Guangzhou 510000, China

⁴ South China Sea Bio-Resource Exploitation and Utilization Collaborative Innovation Center,
Guangzhou 510000, China

Submitted: January 31, 2019. Final revision received: July 15, 2019. Accepted: July 21, 2019

Supplementary material

* Corresponding authors; e-mail: xykong@scsio.ac.cn (X. Kong); gongli1027@163.com (Li Gong)




Figure S1. Inferred secondary structures of the two types of rDNA sequences of *Paraplagusia blochii*.