

Rooting of cuttings of *Passiflora suberosa*, a medicinal passion fruit species: characterization and modulation by external biochemical factors

Johnatan Vilasboa, Cibele Tesser da Costa, Hélio Nitta Matsuura and Arthur Germano Fett-Neto

Plant Physiology Laboratory, Center for Biotechnology and Department of Botany, Federal University of Rio Grande do Sul (UFRGS), CP 15005, Porto Alegre, RS, 91501-970, Brazil

ARTICLE HISTORY

Received 1 July 2019

Accepted 12 September 2019

CONTACT Arthur Germano Fett-Neto E-mail: fettneto@cbiot.ufrgs.br

Supplementary material

Table S1. Rooting percentage of *Passiflora suberosa* cuttings under three different IBA concentrations. DAE = days after exposure.

IBA concentration (ppm)	Rooting (%)			
	9 DAE	12 DAE	15 DAE	18 DAE
0	56	83	100	100
0.1	78	94	100	100
1	61	94	94	89

Table S2. Rooting percentage of *Passiflora suberosa* cuttings under four modulation treatments. DAE = days after exposure.

Condition	Rooting (%)	
	11 DAE	13 DAE
T1 (no IBA, unmodulated)	42	59
T2 (IBA, unmodulated)	67	95
T3 (no IBA, up-modulated)	40	53
T4 (IBA, down-modulated)	0	0