

Age-Related Effects on Cross-Modal Duration Perception

Alexandra N. Scurry^{1,*}, Dustin Dutcher¹, John S. Werner² and Fang Jiang¹

¹ Department of Psychology, University of Nevada, Reno, NV 89557, USA

² Department of Ophthalmology and Vision Science, University of California, Davis Medical Center,
Sacramento, CA 95817, USA

Received 15 March 2019; accepted 30 August 2019

* To whom correspondence should be addressed. E-mail: ascurry@unr.edu

Supplementary Material




Figure S1. Individual variability in perceptual compression and expansion independent of trial order. The percentage of the test V stimulus reported as longer than the standard V stimulus is plotted for each participant from the older (left panel) and younger (right panel) group. Data are from both standard-first and test trials combined. Individuals are color-coded.




Figure S2. Individual variability in perceptual compression and expansion for standard-first trials. The percentage of the test V stimulus reported as longer than the standard V stimulus is plotted for each participant from the older (left panel) and younger (right panel) group for standard-first trials. Individuals are color-coded.




Figure S3. Individual variability in perceptual compression and expansion for test first trials.

The percentage of the test V stimulus reported as longer than the standard V stimulus is plotted for each participant from the older (left panel) and younger (right panel) group for standard-first trials. Individuals are color-coded.