

# Macro- and microhabitat preferences of Eastern Hermann's tortoise

*(Testudo hermanni boettgeri)*

Marko Nikolić<sup>1,\*</sup>, Jovana Cvetković<sup>1</sup>, Dragana Stojadinović<sup>1</sup> and Jelka Crnobrnja-Isailović<sup>1,2</sup>

1 - Department of Biology and Ecology, Faculty of Sciences and Mathematics, University of Niš, Niš, Serbia

2 - Department of Evolutionary Biology, Institute for Biological Research "Siniša Stanković" - Institute of National Importance for Republic of Serbia, University of Belgrade, Beograd, Serbia

\*Corresponding author; e-mail: zerocool.axl@gmail.com

## **Supplementary Material**

**Table S1.** Habitat types used to characterize the landscape based on manual photo interpretation of aerial orthoimages. Cover categories are: 0 (0%), 1 (1-10%), 2 (10-25%), 3 (25-50%), 4 (50-75%), and 5 (75-100%).

Vegetation types	Cover categories		
	Tree	Shrub	Grass
Barren land	0-1-2	0-1-2	0-1-2
Grassland	0-1-2	0-1-2	3-4-5
Open shrubland	0-1-2	3	3-4-5
Dense shrubland	0-1-2	3-4-5	0-1-2-3
Open woodland	3	0-1-2	0-1-2
Wooded grassland	3	0-1-2	3-4-5
Wooded grassland-shrubland	3	3-4-5	3-4-5
Open forest	4	0-1-2-3-4-5	0-1-2-3-4-5
Dense forest	5	0-1-2-3-4-5	0-1-2-3-4-5