

**DNA-barcode and endophallus morphology delimit congruent species
in a systematic revision of the oxyporine rove beetles of Russia
(Coleoptera: Staphylinidae: Oxyporinae)**

Alexandra S. Tokareva

Zoological Institute, Russian Academy of Sciences, Universitetskaya nab. 1, St. Petersburg
199034, Russia

Fedor V. Konstantinov

Zoological Institute, Russian Academy of Sciences, Universitetskaya nab. 1, St. Petersburg
199034, Russia

Department of Entomology, Faculty of Biology, St. Petersburg State University,
Universitetskaya nab. 7/9, St. Petersburg 199034, Russia

Adam J. Brunke

Canadian National Collection of Insects, Agriculture and Agri-Food Canada, Ottawa, ON,
K1A 0C6, Canada

Alexey Solodovnikov

Natural History Museum of Denmark, Zoological Museum, Universitetsparken 15,
Copenhagen 2100, Denmark

Zoological Institute, Russian Academy of Sciences, Universitetskaya nab. 1, St. Petersburg
199034, Russia

Supplementary material

- MATERIAL S1 Pairwise nucleotide distance matrix of the barcoding CO1 region for *Oxyporus* species. Genetic distances in terms of the number of base substitutions per site between sequences.
- MATERIAL S2 Morphological matrix for *Oxyporus* in nexus format.
- MATERIAL S3 COI sequences matrix, aligned with MAFFT v. 7.450, in fasta format.
- MATERIAL S4-S24 Reconstructed ancestral states for characters 1-6, 9-14, 16-19, 21, 23-25, 27, 29 in the Extensible TreeGraph format.