

# The SIR dynamic model of infectious disease transmission and its analogy with chemical kinetics

Cory M. Simon<sup>1</sup>

<sup>1</sup>School of Chemical, Biological, and Environmental Engineering, Oregon State University.

Cory.Simon@oregonstate.edu

May 12, 2020

## Abstract

Mathematical models of the dynamics of infectious disease transmission are used to forecast epidemics and assess mitigation strategies. We reveal that the classic Susceptible-Infectious-Recovered (SIR) epidemic model resembles a dynamic model of a batch reactor carrying out an autocatalytic reaction with catalyst deactivation. This analogy between disease transmission and chemical reactions allows chemists and chemical engineers to peer into dynamic models of infectious disease transmission and gives insights into the kinetics of autocatalytic reactions.

## 1 introduction

Mathematical models of the dynamics of infectious disease transmission [1, 2] are useful for forecasting epidemics, assessing intervention strategies, and inferring properties of diseases.

In compartmental epidemic models [3], each member of the population is categorized based on their disease status in addition to, possibly, their attributes and the treatment they received. The dynamics of disease transmission are then typically modeled with differential equations that describe the flow of individuals to and from the compartments as the population mixes, the disease is spread/contracted, and infected individuals progress through the stages of the disease. Differential equations are a natural choice because we can make reasonable assumptions about the *rates* at which people are infected and progress through the stages of the disease. In this article, we highlight the analogy between compartmental dynamic models of disease transmission and chemical reaction kinetics.

## 2 the SIR model

In the classic SIR model of an epidemic [2, 4–6], each member of the population belongs to one of three compartments: Susceptible, Infectious, or Recovered.

## 2.1 the reactions





Figure 1: The SIR model. The boxes represent the set of Susceptible, Infectious, and Recovered individuals. The arrows represent flow from one compartment to another and are annotated with per capita flow rates.

Susceptible folks can contract the disease if they come into contact with an infectious individual. Once infected, they move into the infectious compartment, assuming zero delay between infection and the ability to transmit the disease. This is analogous to an irreversible autocatalytic chemical reaction [7, 8] between a reactant, S, and catalyst, I:



Infectious individuals eventually recover or die from the disease, entering the recovered compartment, and then cannot transmit the disease or contract it again. This is analogous to a reaction where the catalyst, I, irreversibly degrades or converts to a deactivated form, R:



The R category is sometimes called the Removed category [9] instead, emphasizing its inclusion of disease-induced deaths.

We assume that recovery from the disease confers permanent immunity to reinfection, thus neglecting the possibility of an  $R \longrightarrow S$  reaction.

So, the SIR model of an epidemic is analogous to an autocatalytic reaction (rxn. {1}) with catalyst deactivation (rxn. {2}). An infectious individual (the catalyst, I) (i) converts susceptibles (the reactant, S) into more infectious individuals (more catalyst) and (ii) recovers (deactivates) with time.

Fig. 1 depicts the flow of individuals through compartments under the SIR model, induced by rxns. {1} and {2}. Note the absence of flow to/from external populations; as in a closed batch reactor, we neglect immigration and emigration. Moreover, we take births and deaths (from causes other than the disease) to be negligible over the time scale of the epidemic.

## 2.2 the dynamic mathematical model

Mathematically, the SIR model [4–6, 10] is equivalent to a dynamic model of a well-mixed, isothermal batch reactor carrying out the two homogeneous, elementary rxns. {1} and {2}.

Let  $[S](t)$ ,  $[I](t)$ , and  $[R](t)$  be the fraction of the population that is susceptible, infectious, and recovered, respectively, at time  $t$ . Considering a large population, we treat  $[S]$ ,  $[I]$ , and  $[R]$  as continuous variables.

**The incidence rate.** Assuming their spatial mixing is uniform [11, 12], we invoke the law of mass action to model the rate at which susceptible and infectious individuals “react” via bimolecular, autocatalytic rxn. {1}. The incidence rate of the disease, i.e. the number of new infections per unit time [10], is then  $\beta[S][I]$  (per capita). A symmetric, bilinear function of  $[S]$  and  $[I]$ , intuitively, the incidence rate

doubles if  $[I]$  ( $[S]$ ) doubles while  $[S]$  ( $[I]$ ) is fixed. The second-order transmission rate constant  $\beta > 0$  is the product of the average frequency of contacts of a person and the transmissibility of the disease (the probability of transmission conditioned upon contact).

**The recovery rate.** We posit that infectious individuals “decay” (recover) via rxn. {2} with first-order kinetics, i.e., with rate  $\gamma[I]$  (per capita). The inverse of the first-order recovery rate constant  $\gamma > 0$  is the average time period that an infected individual is infectious<sup>1</sup> [13].

With the assumptions above (see Fig. 1 for flows), we arrive at the following set of nonlinear, coupled differential equations that comprise the SIR dynamic model of infectious disease transmission:

$$\frac{d[S]}{dt} = -\beta[S][I] \quad (1)$$

$$\frac{d[I]}{dt} = \beta[S][I] - \gamma[I] \quad (2)$$

$$\frac{d[R]}{dt} = \gamma[I]. \quad (3)$$

The only two parameters in the SIR model are the transmission and recovery rate constants,  $\beta$  and  $\gamma$ , respectively. While  $\gamma$  could be estimated independently from studies on the duration of infectiousness,  $\beta$  could be identified by fitting differential eqns. 1-3 to epidemic time series data (case counts) [10, 12], much like identifying a reaction rate constant from concentration time series [14]. Adding eqns. 1-3 confirms that the SIR model considers a closed system and neglects demography, i.e.,  $[S](t) + [I](t) + [R](t) = 1, \forall t \geq 0$ . As a consequence, eqns. 1 and 2 fully determine the SIR model dynamics, and  $[R](t)$  follows from  $[R](t) = 1 - [S](t) - [I](t)$ .

### 2.3 the replacement and basic reproduction numbers

Two important numbers aid our characterization and understanding of SIR model dynamics: the replacement number,  $r$ , and the basic reproduction number,  $\mathcal{R}_0$ .

The *replacement number*,  $r = r(t)$ , is the expected number of folks (directly) infected by a typical infectious individual, mixing in the population, over the course of their infectiousness [2]. Because the concentration of susceptible folks  $[S] = [S](t)$  influences the frequency that a typical infectious individual contacts a susceptible individual,  $r$  changes over time. In the SIR model, a typical infectious individual is expected to be infectious for a time period of  $\gamma^{-1}$  and, during this time, produce  $\beta[S](t)$  new infections per unit time (incidence rate per infectious individual). The replacement number is therefore:

$$r = r(t) = \frac{\beta}{\gamma}[S](t). \quad (4)$$

The *basic reproduction number*,  $\mathcal{R}_0$ , is defined as the initial replacement number when *one* infectious individual is introduced into an *all-susceptible* population [2, 6]. Because the entire population is

---

<sup>1</sup>The probabilistic implication of first-order decay for a single infected individual is that their time period of infectiousness is an exponentially distributed random variable with mean  $\gamma^{-1}$  [3].

susceptible in this context,  $\mathcal{R}_0$  in the SIR model is the replacement number in eqn. 4 when  $[S] \approx 1$ :

$$\mathcal{R}_0 = \frac{\beta}{\gamma}. \quad (5)$$

i.e.,  $\mathcal{R}_0$  is the expected number of infections directly caused by a single infectious individual introduced into an entirely susceptible population over the course of their infectiousness [2].

The replacement number,  $r$ , and basic reproduction number,  $\mathcal{R}_0$ , are both dimensionless and are properties of both the disease and the population [15]. While  $r = r(t)$  changes with time,  $\mathcal{R}_0$  is constant and defined only at the initial stage when one infectious individual is introduced to an all-susceptible population. Notably, the two numbers are related via  $r(t) = \mathcal{R}_0[S](t)$ .

If the basic reproduction number  $\mathcal{R}_0$  is large (small), the infectious are infectious for a long (short) period of time, the disease is (not) easily transmitted, and/or the mixing of susceptibles and infectious is (not) vigorous. Under the analogy with chemical kinetics, since the activity and longevity of the catalyst,  $l$ , are embedded in  $\beta$  and  $\gamma$ , respectively:  $\mathcal{R}_0$  is large (small) if the catalyst has a high (low) activity and/or remains active for a long (short) time. Because  $r = \mathcal{R}_0[S]$ , these remarks hold for the replacement number,  $r$ , as well. Though,  $r$  decreases as the concentration of the reactant,  $[S]$ , decreases, owing to the reduced frequency that any given catalyst particle encounters a reactant particle to catalyze its conversion into another catalyst particle by rxn. {1}.

### 3 SIR model dynamics

In the SIR model, what happens if we introduce a small number of infectious individuals into a large population of susceptible individuals? This is akin to injecting our deactivating auto-catalyst,  $l$ , into a well-mixed batch reactor containing pure  $S$ . The corresponding initial conditions are:

$$[S](0) = [S]_0 \quad (6)$$

$$[I](0) = [I]_0 \quad (7)$$

$$[R](0) = 0, \quad (8)$$

with  $[S]_0 + [I]_0 = 1$ ,  $[S]_0, [I]_0 > 0$ , and  $[I]_0 \ll 1$ . We consider  $[R](0) = 0$  for the interesting case where the population is exposed to a novel pathogen to which it has no immunity.

#### 3.1 the replacement number determines $[I]'(t)$

The replacement number  $r(t)$  in eqn. 4 is key to understanding SIR model dynamics. By inspection of eqn. 2,  $[I](t)$  is increasing at time  $t$  if the replacement number  $r(t) > 1$  and decreasing if  $r(t) < 1$ . This is intuitive: if the typical infectious person mixing in the population is expected to infect less than one susceptible person before they recover, they are not expected to replace themselves with a new infectious individual to propagate the disease, and  $[I](t)$  decreases. Note  $r(t) < 1 \iff [S](t) < \mathcal{R}_0^{-1}$ .

### 3.2 does an epidemic ensue?

We first address a qualitative question: given the initial conditions in eqn. 6-8, does the disease invade the population? The outcome depends on the initial replacement number  $r_0 := r(0) = \mathcal{R}_0[S]_0$  in a threshold manner. If  $r_0 > 1$ ,  $[I](t)$  initially increases. i.e., the disease spreads; an epidemic ensues. If  $r_0 < 1$ ,  $[I](t)$  initially decreases and, since  $[S](t)$  is a decreasing function of time (see eqn. 1), decays to zero. i.e., the disease dies out; an epidemic does not ensue.

Under the chemical reaction analogy, if  $r_0 < 1$  ( $r_0 > 1$ ), the injected catalyst particles deactivate via rxn. {2} faster (slower) than they catalyze rxn. {1} to convert the reactant, S, into more catalyst, I, to propagate autocatalytic rxn. {1}.

For our remaining analysis, we take  $r_0 > 1$  and further analyze the dynamics of an SIR epidemic.

### 3.3 initial exponential growth

Early in the epidemic, the number of infectious folks grows, approximately, exponentially with growth rate  $(r_0 - 1)\gamma$ :

$$[I](t) \approx [I]_0 e^{(r_0 - 1)\gamma t}. \quad (9)$$

This follows from eqn. 2 if we approximate  $[S](t) \approx [S]_0$ , valid only in the initial stage of the epidemic; as the disease spreads,  $[S]$  decreases and diminishes the replacement number. Eqn. 9 is thus an overestimate.

Since eqn. 9 is also a valid initial approximation for  $r_0 < 1$ , it reinforces that an epidemic will not ensue if  $r_0 < 1$ , since  $[I](t)$  would then decay approximately exponentially.

### 3.4 a simulation of an SIR model epidemic

Fig. 2 shows a simulation of SIR model dynamics for  $\mathcal{R}_0 = 2$  and  $[I]_0 = 10^{-5}$  (see Sec. A1.1 for code). Initially, the concentration of infectious folks,  $[I]$ , grows approximately exponentially (see Fig. A8 for a comparison of  $[I](t)$  with eqn. 9). The concentration of susceptible folks,  $[S]$ , decreases monotonically as the disease invades. As a result, the frequency with which any given infectious individual comes into contact with a susceptible individual decreases. In conjunction with the infectious folks recovering, this eventually causes the (net) growth rate of  $[I]$  to diminish and, when the replacement number  $r$  in eqn. 4 drops below one, causes  $[I]$  to decay. The epidemic dies out ( $\lim_{t \rightarrow \infty} [I](t) = 0$ ). The R




Figure 2: Numerical approximation of the solution to the SIR model in eqns. 1-3 ( $\mathcal{R}_0 = 2$ ) with initial conditions in eqns 6-8 ( $[I]_0 = 10^{-5}$ ).

category accumulates the folks that have been infected by and have recovered from the disease,  $[R](t) = \gamma \int_0^t [I](\tau) d\tau$  (see eqn. 3 with initial condition 8). Notably, the disease does not infect the entire population, even after an infinite amount of time. i.e., the epidemic dies out not because the population is depleted of susceptible folks, but rather because it is depleted of infectious folks [5, 12].

Under the chemical kinetics analogy, the autocatalytic rxn. {1} begins to slow down after the concentration of the reactant,  $[S]$ , decreases sufficiently. Once the replacement number  $r(t)$  drops below one, the catalyst,  $I$ , is deactivating via rxn. {2} faster than it is converting the reactant,  $S$ , into more catalyst to replenish itself via rxn. {1}. Consequently, the catalyst concentration,  $[I]$ , begins to drop and decay to zero. Owing to catalyst deactivation (rxn. {2}), not all reactant,  $S$ , is consumed, even after an infinite amount of time.

### 3.5 solution in the $([S], [I])$ phase space




Figure 3: The trajectory of the solution to the SIR model ( $\mathcal{R}_0 = 2$ , initial conditions in eqns 6-8 with  $[I]_0 = 10^{-5}$ ) in the  $([S], [I])$  phase plane, given by eqn. 10. The trajectory is colored according to time. The initial conditions, peak prevalence of infectious folks, and final conditions are marked. Since  $[S](t) + [I](t) \leq 1$ , solutions in the gray region are infeasible.

We can analytically find the trajectory of the solution to eqns. 1-2 in the  $([S], [I])$  phase plane. Dividing eqn. 2 by eqn. 1 takes us into the phase plane by giving a differential equation with a  $\frac{d[I]}{d[S]}$  derivative, with the view of  $[I]$  as a function of  $[S]$ . Separating, integrating, and applying the initial conditions in eqns. 6 and 7, we arrive at the solution path [3]:

$$[I](t) = 1 - [S](t) + \frac{1}{\mathcal{R}_0} \log \left( \frac{[S](t)}{[S]_0} \right). \quad (10)$$

Fig. 3 shows the trajectory given in eqn. 10 ( $\mathcal{R}_0 = 2$ ,  $[I]_0 = 10^{-5}$ ) and reinforces that  $[S](t)$  decreases monotonically with time, that  $[I](t)$  increases, peaks, then diminishes to zero, and that a fraction of the population remains susceptible after the epidemic dies out.

### 3.6 final size

Given the epidemic (autocatalytic reaction) dies out before all of the susceptible folks (reactant) are infected, what fraction of the population (reactant) remains uninfected (unreacted) after the epidemic ends?

We find an implicit equation for  $[S]_\infty := \lim_{t \rightarrow \infty} [S](t)$  by taking the limit  $t \rightarrow \infty$  in eqn. 10:

$$0 = 1 - [S]_\infty + \frac{1}{\mathcal{R}_0} \log \left( \frac{[S]_\infty}{[S]_0} \right). \quad (11)$$




Figure 4: The fraction of the population that (a) remains susceptible (never infected) after the SIR epidemic runs its course,  $[S]_\infty$ , via eqn. 11 and (b) is infected at peak prevalence,  $\max_t [I](t)$ , via eqn. 12, both as a function of  $\mathcal{R}_0$ . Corresponds to initial conditions in eqns. 6-8 with  $[I]_0 = 10^{-5}$ .

We used the fact that the epidemic eventually dies out,  $\lim_{t \rightarrow \infty} [I](t) = 0$ .  $[S]_\infty$  is the unique root of eqn. 11 in  $(0, \mathcal{R}_0^{-1})$  [2]. The fraction of the population infected over the course of the epidemic is  $\lim_{t \rightarrow \infty} [R](t) =: [R]_\infty = 1 - [S]_\infty$  since the R category accumulates those that have recovered from the disease. Fig. 4a shows that, as  $\mathcal{R}_0$  increases from one, more of the population will be infected over the course of the epidemic.

### 3.7 peak prevalence of infectious folks

The peak prevalence of infectious folks,  $\max_t [I](t)$ , is an important quantity since it determines the maximum burden on healthcare. Both eqn. 10 and eqn. 2 show that the maximum in  $[I](t)$  (look for  $\frac{d[I]}{d[S]} = 0$  or  $\frac{d[I]}{dt} = 0$ , respectively) occurs when  $[S](t) = \mathcal{R}_0^{-1}$  (i.e., when replacement number  $r(t) = 1$ ). Via eqn. 10, then:

$$\max_t [I](t) = 1 - \frac{1}{\mathcal{R}_0} [1 + \log(\mathcal{R}_0 [S]_0)]. \quad (12)$$

Fig. 4b shows how the peak prevalence of infectious folks increases with  $\mathcal{R}_0$ . Before the maximum, the replacement number  $r(t) > 1$  and  $[I](t)$  is increasing; after the peak,  $r(t) < 1$  and  $[I](t)$  is decreasing to zero.

### 3.8 herd immunity

A population is conferred *herd immunity* when a sufficient fraction is immune to the disease so as to indirectly protect it from invasion of the disease upon the introduction of an infectious individual.

Notably, such immunity could be acquired by either previous infection or by vaccination. [2]

What fraction  $v$  of a population must be immune to achieve herd immunity?  $[I](t)$  will decrease upon introducing an infectious individual if the replacement number  $r(t) < 1$ , i.e. if  $[S] < \mathcal{R}_0^{-1}$ . Thus, a fraction  $v > 1 - \mathcal{R}_0^{-1}$  of the population must be immune to achieve herd immunity. Fig. 5 shows the region  $v > 1 - \mathcal{R}_0^{-1}$  (gray) and illustrates that, if  $\mathcal{R}_0$  is large (small), more (less) of the population must be immune to prevent an epidemic via herd immunity.

In the chemical kinetics analogy, herd immunity results from reducing the concentration of the reactant,  $[S]$ , so that a catalyst particle,  $I$ , fed to the reactor is expected to deactivate before it encounters an  $S$  particle and auto-catalyzes rxn. {1} to replace itself. i.e., to achieve herd immunity,  $[S]$  must be reduced sufficiently to make the replacement number  $r$  less than one.

To account for vaccination dynamics in the midst of an epidemic, the SIR model can be adapted to allow flow in Fig. 1 from the  $S$  category directly to the  $R$  category, representing the administration of a vaccine to susceptible folks that confers complete immunity [10,16,17].

An interesting comparison is when herd immunity is achieved through a vaccine ( $S \rightarrow R$ ) versus through infection and recovery ( $S \rightarrow I \rightarrow R$ ). One obvious benefit of the path  $S \rightarrow R$  over the path  $S \rightarrow I \rightarrow R$  is avoiding disease-induced suffering and, possibly, death. Anyway, administering an effective vaccine to a fraction  $v = 1 - \mathcal{R}_0^{-1}$  of an all-susceptible population will suffice to achieve herd immunity. However, this does not imply that the same fraction will be infected if herd immunity is instead achieved through infection and recovery, by introducing infectious individuals and allowing the epidemic to run its course. The (orange) curve in Fig. 5 shows  $[R]_\infty$ , the fraction of the population that will be infected by the disease over the course of an SIR epidemic (without any vaccination). A significantly greater fraction of the population will be infected by the disease if we let it run its course than would have needed vaccination to achieve herd immunity. When  $[S](t)$  reaches  $\mathcal{R}_0^{-1}$  in an SIR epidemic (without vaccination),  $[I](t)$  is at its maximum (see eqn. 10 and Fig. 3); these infectious individuals will infect more susceptible folks before they all recover, even though on *average* each is expected to recover before they infect a susceptible individual ( $r < 1$ ).




Figure 5: Herd immunity. Gray region shows the fraction  $v > 1 - \mathcal{R}_0^{-1}$  of a population that must be immune to achieve herd immunity. Such immunity could be conferred by vaccination or by previous infection. For comparison, orange line shows  $[R]_\infty$ , the number of folks that will be infected over the course of an SIR epidemic without any vaccination ( $[I]_0 = 10^{-5}$ ).

### 3.9 conclusion: $\mathcal{R}_0$ is a useful tool

The basic reproduction number  $\mathcal{R}_0$  is a useful property of the SIR model that influences if the disease invades the population, the initial exponential growth rate of infectious folks, the fraction of the population infected over the course of the epidemic, the peak prevalence of infectious folks, and the fraction of the population that must be vaccinated to achieve herd immunity. This result is intuitive since the average frequency of contacts of an individual in the population, the transmissibility of the disease, and the duration of infectiousness are all embedded in the basic reproduction number  $\mathcal{R}_0$ .

## 4 extensions to the SIR model

The SIR model is a very simple epidemic model, but we can extend it to model other epidemiological factors and prevention/control measures by introducing:

**additional compartments.** For example, the SEQIJR model [18] introduces three additional compartments: Exposed, Quarantined, and Isolated (J). Usually, the E compartment is included to model the latent period of a disease; exposed individuals have been exposed to and infected by the disease but are not yet infectious [3, 19]. The Q and J compartments are included to model control measures: both quarantining individuals exposed to the disease and isolating infectious individuals. Members of the Q and J compartments contact susceptible folks with a reduced frequency [3]. See Fig. 6. As another example, to distinguish between Asymptomatic and symptomatic infectious individuals [20], which may have different




Figure 6: Introducing additional compartments to the SIR model can account for other epidemiological factors and control measures. Exposed individuals have been exposed to and infected by the disease, but cannot yet transmit the disease, due to a latent period. Owing to control measures, Quarantined and isolated (J) individuals contact susceptible folks with a reduced frequency.

recovery rates, frequencies of contacts, and transmissibilities, we can introduce an A compartment. Finally, to account for different mixing patterns, infectiousness, and susceptibility among different age groups [3], age-structured compartmental models partition the S and I compartments into age groups.

Additional compartments complicate the derivation of the basic reproduction number  $\mathcal{R}_0$  from the model [21]. Still, the peak prevalence of infectious folks and final size of the epidemic increase with  $\mathcal{R}_0$  in most models [22]. Further, a more precise definition of  $\mathcal{R}_0$  is needed if compartments are introduced to account for heterogeneity in the population [23].

**time-varying parameters.** For example, if we allow the transmission rate constant  $\beta$  to vary with time,  $\beta = \beta(t)$ , we can model (a) members of the population changing/adapting their behavior, e.g., reducing their frequency of social contacts and practicing social distancing [24, 25] and (b) seasonality

of an infectious disease [26, 27]. Reducing the transmission rate constant  $\beta$ , by members of the population reducing their frequency of social contacts and/or by reducing the transmissibility of the disease, reduces the replacement number  $r$  (see eqn. 4); if  $r$  is reduced below one, the prevalence of infectious folks will decrease.

**stochasticity.** As opposed to the deterministic differential eqns. 1-3, we can introduce randomness into the SIR model to account for the stochastic and uncertain nature of human interaction and disease transmission [28, 29]. Stochastic epidemic models aim to describe the probabilistic distribution of outcomes, e.g., the distribution of  $[S]_{\infty}$  [28, 29]. Stochasticity can be particularly important for small populations and in the early stage of an epidemic when there are small numbers of infectious individuals [29]. Analogously, stochastic models of chemical reaction dynamics are necessary when the reactants are not abundant, such as in a biological cell [30, 31].

**spatial heterogeneity.** Finally, we can model spatial heterogeneity in an epidemic in a discrete [32, 33] or continuous [10] manner and allow for more detail/structure in the social contacts/mixing within a population or between populations [34, 35]. Modeling the spatial movement of susceptible and infectious individuals as a diffusive process results in reaction-diffusion equations, also familiar to chemists and chemical engineers [10]. Compartmental, metapopulation epidemic models with travel between spatially segregated regions resemble models of multiple batch reactors connected with pipes that allow flow between them [32, 36].

## 5 conclusions

Mathematical models of the dynamics of disease transmission are used to forecast epidemics and assess mitigation strategies. The analogy between disease transmission and an autocatalytic reaction allows chemists and chemical engineers to peer into epidemic models. Moreover, the analogy illustrates how concepts in one field can transfer, aid understanding, and generate insights to/in another field.

## References

- [1] Fred Brauer. Mathematical epidemiology: Past, present, and future. *Infectious Disease Modelling*, 2(2):113–127, 2017.
- [2] Herbert W Hethcote. The mathematics of infectious diseases. *SIAM Review*, 42(4):599–653, 2000.
- [3] Fred Brauer. Compartmental models in epidemiology. In *Mathematical Epidemiology*, pages 19–79. Springer, 2008.
- [4] William Ogilvy Kermack and Anderson G McKendrick. A contribution to the mathematical theory of epidemics. *Proceedings of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 115(772):700–721, 1927.
- [5] James D Murray. Epidemic models and the dynamics of infectious diseases. In *Mathematical Biology*, pages 610–650. Springer, 1993.

- [6] Leah Edelstein-Keshet. *Mathematical Models in Biology*. SIAM, 1988.
- [7] Peter Schuster. What is special about autocatalysis? *Monatshefte für Chemie-Chemical Monthly*, pages 1–13, 2019.
- [8] SK Scott. Reversible autocatalytic reactions in an isothermal CSTR: multiplicity, stability and relaxation times. *Chemical Engineering Science*, 38(10):1701–1708, 1983.
- [9] James C Frauenthal. *Mathematical modeling in epidemiology*. Springer Science & Business Media, 2012.
- [10] Maia Martcheva. *An Introduction to Mathematical Epidemiology*, volume 61. Springer, 2015.
- [11] Edwin B Wilson and Jane Worcester. The law of mass action in epidemiology. *Proceedings of the National Academy of Sciences*, 31(1):24, 1945.
- [12] Howard Howie Weiss. The SIR model and the foundations of public health. *MATerials MATemáticas*, pages 0001–17, 2013.
- [13] Matt J Keeling and Pejman Rohani. *Modeling infectious diseases in humans and animals*. Princeton University Press, 2011.
- [14] H Scott Fogler. *Essentials of Chemical Reaction Engineering*. Pearson Education, 2010.
- [15] Paul L Delamater, Erica J Street, Timothy F Leslie, Y Tony Yang, and Kathryn H Jacobsen. Complexity of the basic reproduction number ( $R_0$ ). *Emerging Infectious Diseases*, 25(1):1, 2019.
- [16] Tunde Tajudeen Yusuf and Francis Benyah. Optimal control of vaccination and treatment for an SIR epidemiological model. *World Journal of Modelling and Simulation*, 8(3):194–204, 2012.
- [17] Boris Shulgin, Lewi Stone, and Zvia Agur. Pulse vaccination strategy in the SIR epidemic model. *Bulletin of Mathematical Biology*, 60(6):1123–1148, 1998.
- [18] Abba B Gumel, Shigui Ruan, Troy Day, James Watmough, Fred Brauer, P Van den Driessche, Dave Gabrielson, Chris Bowman, Murray E Alexander, Sten Ardal, et al. Modelling strategies for controlling sars outbreaks. *Proceedings of the Royal Society of London. Series B: Biological Sciences*, 271(1554):2223–2232, 2004.
- [19] Maia Martcheva. Techniques for computing  $\mathcal{R}_0$ . In *An Introduction to Mathematical Epidemiology*, pages 91–121. Springer, 2015.
- [20] Ana Luz Vivas-Barber, Carlos Castillo-Chavez, and Ernest Barany. Dynamics of an SAIQR Influenza Model. *Biomath*, 3(2):1409251, 2014.
- [21] Pauline van den Driessche. Reproduction numbers of infectious disease models. *Infectious Disease Modelling*, 2(3):288–303, 2017.
- [22] P Van den Driessche and James Watmough. Further notes on the basic reproduction number. In *Mathematical Epidemiology*, pages 159–178. Springer, 2008.

- [23] Odo Diekmann, Johan Andre Peter Heesterbeek, and Johan AJ Metz. On the definition and the computation of the basic reproduction ratio  $R_0$  in models for infectious diseases in heterogeneous populations. *Journal of Mathematical Biology*, 28(4):365–382, 1990.
- [24] Eli P Fenichel, Carlos Castillo-Chavez, M Graziano Ceddia, Gerardo Chowell, Paula A Gonzalez Parra, Graham J Hickling, Garth Holloway, Richard Horan, Benjamin Morin, Charles Perrings, et al. Adaptive human behavior in epidemiological models. *Proceedings of the National Academy of Sciences*, 108(15):6306–6311, 2011.
- [25] Timothy C Reluga. Game theory of social distancing in response to an epidemic. *PLoS Computational Biology*, 6(5), 2010.
- [26] David N Fisman. Seasonality of infectious diseases. *Annual Review of Public Health*, 28:127–143, 2007.
- [27] Joan L Aron and Ira B Schwartz. Seasonality and period-doubling bifurcations in an epidemic model. *Journal of Theoretical Biology*, 110(4):665–679, 1984.
- [28] Linda JS Allen. An introduction to stochastic epidemic models. In *Mathematical epidemiology*, pages 81–130. Springer, 2008.
- [29] Tom Britton. Stochastic epidemic models: a survey. *Mathematical Biosciences*, 225(1):24–35, 2010.
- [30] Radek Erban, Jonathan Chapman, and Philip Maini. A practical guide to stochastic simulations of reaction-diffusion processes. *arXiv preprint arXiv:0704.1908*, 2007.
- [31] Daniel T Gillespie, Andreas Hellander, and Linda R Petzold. Perspective: Stochastic algorithms for chemical kinetics. *The Journal of Chemical Physics*, 138(17):05B201\_1, 2013.
- [32] P Van den Driessche. Spatial structure: Patch models. In *Mathematical epidemiology*, pages 179–189. Springer, 2008.
- [33] Matt J Keeling and Ken TD Eames. Networks and epidemic models. *Journal of the Royal Society Interface*, 2(4):295–307, 2005.
- [34] Duygu Balcan, Bruno Gonçalves, Hao Hu, José J Ramasco, Vittoria Colizza, and Alessandro Vespignani. Modeling the spatial spread of infectious diseases: The global epidemic and mobility computational model. *Journal of Computational Science*, 1(3):132–145, 2010.
- [35] Phillip Stroud, Sara Del Valle, Stephen Sydoriak, Jane Riese, and Susan Mniszewski. Spatial dynamics of pandemic influenza in a massive artificial society. *Journal of Artificial Societies and Social Simulation*, 10(4):9, 2007.
- [36] Julien Arino, Jonathan R Davis, David Hartley, Richard Jordan, Joy M Miller, and P Van Den Driessche. A multi-species epidemic model with spatial dynamics. *Mathematical Medicine and Biology*, 22(2):129–142, 2005.
- [37] Christopher Rackauckas and Qing Nie. Differentialequations.jl a performant and feature-rich ecosystem for solving differential equations in Julia. *Journal of Open Research Software*, 5(1), 2017.

## A1 the appendix

### A1.1 simulating the SIR model in Julia

Code in the Julia language to numerically approximate the solution to the SIR model, via the DifferentialEquations.jl [37] package, is below.

```
using DifferentialEquations

R0 = 2.0    # basic reproduction number

# right-hand side of the ODE, viewed as:
#   du/dt = f = f(u, p, t)
#   where u := [S, I, R]
function update_f!(f, u, p, t)
    # for clarity, unpack vector u
    s = u[1]
    i = u[2]
    r = u[3]
    # update f
    f[1] = -R0 * s * i
    f[2] =  R0 * s * i - i
    f[3] = i
end

# initial condition
eps = 10^-5 # initial fraction infectious
u0 = [1-eps; eps; 0.0]

# define the ODE problem
time_span = (0.0, 25.0)
prob = ODEProblem(update_f!, u0, time_span)

# numerically solve ODE
sol = solve(prob)

# obtain numerical approximation to the solution at e.g. t = 10
sol(10.0)
```

### A1.2 an alternative visualization of SIR model dynamics

Fig. A7 presents an alternative visualization of SIR model dynamics that emphasizes  $[S](t) + [I](t) + [R](t) = 1$  for all  $t > 0$ .




Figure A7: Numerical approximation of the solution to the SIR model in eqns. 1-3 for  $\mathcal{R}_0 = 2$  and initial conditions in eqns 6-8 ( $[I]_0 = 1e - 5$ ). At any given time  $t$ , the proportion of the panel colored green, red, and blue, respectively, represents the fraction of the population in the susceptible, infectious, and recovered compartment.

### A1.3 early exponential growth of $[I]$

See Fig. A8 to see how the approximation of  $[I](t)$  in eqn. 9 fares.




Figure A8: The  $[I](t)$  curve from Fig. 2 along with eqn. 9 (gray, dashed line) to show that  $[I](t)$  exhibits, approximately, exponential growth in the early stage of the epidemic. The fraction of the population that is infectious,  $[I](t)$ , is shown on both a (a) linear and (b) log scale.