

Supplementary Material

Antifungal activity and mechanism of action of the Co(III) coordination complexes with diamine chelate ligands against reference and clinical strains of *Candida* spp.

Katarzyna Turecka^{1*}, Agnieszka Chylewska², Anna Kawiak³, Krzysztof Waleron¹

* Correspondence: Katarzyna Turecka: tureckak@gumed.edu.pl

Supplementary Tables

TABLE S2. MFC values in $\mu\text{g/mL}$ of $[\text{CoCl}_2(\text{dap})_2]\text{Cl}$ (**1**), $[\text{CoCl}_2(\text{en})_2]\text{Cl}$ (**2**) stored in darkness and light against *Candida* strains evaluated after 14 and 30 days. Data shown are mean \pm SD.

Strain	Compound	Darkness		Light	
		14 d*	30 d*	14 d*	30 d*
<i>C. albicans</i> ATCC 10231	$[\text{CoCl}_2(\text{dap})_2]\text{Cl}$ (1)	125 \pm 9.19	125 \pm 6.94	125 \pm 10.41	125 \pm 6.94
	$[\text{CoCl}_2(\text{en})_2]\text{Cl}$ (2)	125 \pm 9.89	125 \pm 9.19	125 \pm 6.94	125 \pm 9.89
<i>C. glabrata</i> ATCC 2001	$[\text{CoCl}_2(\text{dap})_2]\text{Cl}$ (1)	62.5 \pm 5.21	62.5 \pm 4.04	62.5 \pm 4.95	62.5 \pm 4.04
	$[\text{CoCl}_2(\text{en})_2]\text{Cl}$ (2)	62.5 \pm 4.95	62.5 \pm 4.95	62.5 \pm 5.21	62.5 \pm 8.47

*d – day