

Supplementary Material

Virulence Of *Mycobacterium Tuberculosis* Clinical Isolates Is Associated With Sputum Pre-treatment Bacterial Load, Lineage, Survival In Macrophages, And Cytokine Response

Trinh T. B. Tram, Hoang N. Nhung, Srinivasan Vijay, Hoang T. Hai, Do D. A. Thu, Vu T. N. Ha, Tran D. Dinh, Philip M. Ashton, Nguyen T. Hanh, Nguyen H. Phu, Guy E. Thwaites, Nguyen T. T. Thuong

* **Correspondence:** Nguyen T. T. Thuong, thuongntt@oucru.org

1 Supplementary Tables

Supplementary Table 1. Percentage of cell lysis of THP-1 infected with 153 *Mtb* clinical isolates

Cell lysis (%) Day (N, %)	0 - 5	≤5 - 10	≤10 - 30	≤30 - 50	≤50 - 70	≤70 - 90	≤90 - 100
1	153 (100.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
2	151 (98.7)	2 (1.3)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
3	127 (83.0)	9 (5.9)	12 (7.8)	5 (3.3)	0 (0.0)	0 (0.0)	0 (0.0)
4	80 (52.3)	9 (5.9)	30 (19.6)	14 (9.2)	8 (5.2)	4 (2.6)	8 (5.2)
5	61 (39.9)	4 (2.6)	17 (11.1)	19 (12.4)	15 (9.8)	10 (6.5)	27 (17.6)
6	34 (22.2)	3 (2.1)	20 (13.1)	15 (9.8)	27 (17.6)	8 (5.2)	46 (30.1)

N: number of isolates