

TABLE 1. The *in vitro* antifungal activities of the target compounds **4a–4q** at 50 $\mu\text{g/mL}$.

Compounds	Inhibition rate (%)				
	<i>G. zeae</i>	<i>B. dothidea</i>	<i>P. infestans</i>	<i>Phomopsis</i> sp.	<i>T. cucumeris</i>
4a	58.6 \pm 2.2	58.1 \pm 1.6	44.4 \pm 1.5	21.0 \pm 2.4	17.1 \pm 1.2
4b	62.2 \pm 1.4	54.8 \pm 0.7	28.5 \pm 2.0	38.7 \pm 1.3	29.0 \pm 1.2
4c	65.7 \pm 1.3	60.1 \pm 1.1	19.8 \pm 0.6	43.0 \pm 2.9	56.9 \pm 2.4
4d	58.9 \pm 1.1	52.0 \pm 1.2	40.9 \pm 1.4	50.0 \pm 1.3	44.5 \pm 1.5
4e	53.6 \pm 0.7	40.7 \pm 1.1	29.4 \pm 0.7	26.7 \pm 0.4	32.0 \pm 1.4
4f	51.7 \pm 1.1	43.3 \pm 0.1	35.0 \pm 1.9	30.8 \pm 2.3	42.2 \pm 2.0
4g	58.4 \pm 1.2	60.7 \pm 1.2	77.3 \pm 2.1	56.7 \pm 2.1	62.0 \pm 1.0
4h	35.6 \pm 0.6	33.5 \pm 0.8	73.0 \pm 1.0	30.8 \pm 1.0	22.2 \pm 2.2
4i	48.9 \pm 1.7	58.1 \pm 1.5	83.5 \pm 0.6	55.2 \pm 2.1	64.3 \pm 1.5
4j	58.3 \pm 1.6	51.1 \pm 0.9	30.1 \pm 2.6	58.4 \pm 1.7	44.7 \pm 1.6
4k	55.2 \pm 2.2	55.2 \pm 1.2	61.9 \pm 2.0	43.7 \pm 2.0	37.0 \pm 1.8
4l	58.0 \pm 2.3	49.2 \pm 1.3	70.0 \pm 1.2	31.5 \pm 0.9	59.8 \pm 0.9
4m	73.1 \pm 1.0	41.0 \pm 1.6	63.6 \pm 1.3	48.4 \pm 1.1	44.3 \pm 1.6
4n	70.3 \pm 1.1	45.6 \pm 1.1	73.1 \pm 1.8	33.7 \pm 0.8	58.5 \pm 1.8
4o	45.0 \pm 2.2	22.1 \pm 0.9	75.9 \pm 1.2	40.0 \pm 2.3	54.3 \pm 1.7
4p	53.4 \pm 1.9	61.3 \pm 1.1	79.0 \pm 1.1	64.0 \pm 1.3	62.8 \pm 0.7
4q	56.8 \pm 1.5	62.0 \pm 2.0	81.1 \pm 0.3	63.1 \pm 1.2	65.1 \pm 1.3
Dimethomorph	74.3 \pm 2.0	72.3 \pm 1.6	78.2 \pm 1.1	69.3 \pm 1.6	68.3 \pm 1.6