Erratum

In the article by Zhang et al., entitled 'Aspirin plus Clopidogrel as Secondary Prevention after Stroke or Transient Ischemic Attack: A Systematic Review and Meta-Analysis' [Cerebrovasc Dis 2015;39:13–22, DOI: 10.1159/000369778] there are several errors in the published version. The corrections are as follows:

1. In the original article there is a mix-up of figures. The correct figs. 2–6 with the corresponding legends are printed below.
2. In fig. 5 'Comparison of aspirin and clopidogrel versus aspirin or clopidogrel alone on stroke, MI, or vascular death', 19th line, '(p = 0.057)' should be '(p = 0.37)'.
3. On p. 18, left column, 12th line, 'p = 0.15' should be 'p = 0.16'.
4. On p. 18, right column, 13th line, '(fig. 5)' should be '(fig. 6)'.
5. On p. 18, right column, 18th line, '0.84–1.03' should be'0.84–1.01'.

Study or Subgroup	A+C Events	A+C Total	A/C Events	A/C Total	Weight	Risk ratio M-H, Random, 95% CI
Short-term combination						
Bal Dit Sollier 2009	0	10	1	12	0.3%	0.39 [0.02, 8.73]
CARESS 2005	0	51	4	56	0.3%	0.12 [0.01, 2.21]
CHANCE 2013	212	2,584	303	2,586	26.2%	0.70 [0.59, 0.83]
CLAIR 2010	0	46	2	52	0.3%	0.23 [0.01, 4.58]
FASTER 2007	14	198	21	194	5.0%	0.65 [0.34, 1.25]
Subtotal (95% CI)		2,889		2,900	32.0%	0.69 [0.59, 0.81]
Total events	226		331			
Heterogeneity: $Tau^2 = 0$; $Chi^2 = 2.10$, d.f. = 4, (p = 0.72); $I^2 = 0\%$						
Test for overall effect: Z = 4.52 (p < 0.00001)						
Long-term combination						
CHARISMAsub 2011	105	2,157	131	2,163	19.1%	0.80 [0.63, 1.03]
MATCH 2004	339	3,797	347	3,802	28.5%	0.98 [0.85, 1.13]
SPS3 2012	125	1,517	138	1,503	20.5%	0.90 [0.71, 1.13]
Subtotal (95% CI)		7,471		7,468	68.0%	0.92 [0.83, 1.03]
Total events	569		616			
Heterogeneity: $Tau^2 = 0.0$; $Chi^2 = 1.87$, d.f. = 2, (p = 0.39); $I^2 = 0\%$						
Test for overall effect: Z = 1.41 (p = 0.16)						
Total (95% CI)		10,360		10,368	100.0%	0.82 [0.70, 0.96]
Total events	795		947			
Heterogeneity: $Tau^2 = 0.02$; $Chi^2 = 12.63$, d.f. = 7, (p = 0.08); $I^2 = 45\%$						
Test for overall effect: Z = 2.48 (p = 0.01)						
Test for subgroup differences: $Chi^2 = 8.67$; d.f. = 1, (p = 0.03); $I^2 = 88.5\%$						



Fig. 2. Comparison of aspirin and clopidogrel versus aspirin or clopidogrel alone on stroke.

Study or subgroup	A+C Events	A+C Total	A/C Events	A/C Total	Weight	Risk ratio M-H, Fixed, 95% CI
Short-term combination						
Bal Dit Sollier 2009	0	10	1	12	0.2%	0.39 [0.02, 8.73]
CARESS 2005	0	51	4	56	0.5%	0.12 [0.01, 2.21]
CHANCE 2013	204	2,584	295	2,586	32.9%	0.69 [0.58, 0.82]
CLAIR 2010	0	46	2	52	0.3%	0.23 [0.01, 4.58]
FASTER 2007	12	198	21	194	2.4%	0.56 [0.28, 1.11]
Subtotal (95% CI)		2,889		2,900	36.2%	0.67 [0.57, 0.79]
Total events	216		323			
Heterogeneity: $Chi^2 = 2.35$, d.f. = 4 (p = 0.67); $I^2 = 0\%$						
Test for overall effect: Z = 4.77 (p < 0.00001)						
Long-term combination						
CHARISMAsub 2011	91	2,157	114	2,163	12.7%	0.80 [0.61, 1.05]
MATCH 2004	309	3,797	333	3,802	63.8%	0.93 [0.80, 1.08]
SPS3 2012	100	1,517	124	1,503	13.9%	0.80 [0.62, 1.03]
Subtotal (95% CI)		7,471		7,468	63.8%	0.88 [0.78, 0.98]
Total events	500		571			
Heterogeneity: $Chi^2 = 1.54$, d.f. = 2 (p = 0.46); $I^2 = 0\%$						
Test for overall effect: Z = 2.26 (p = 0.02)						
Total (95% CI)		10,360		10,368	100.0%	0.80 [0.73, 0.88]
Total events	716		894			
Heterogeneity: $Chi^2 = 10.26$, d.f. = 7 (p = 0.17); $I^2 = 32\%$						
Test for overall effect: Z = 4.61 (p < 0.00001)						
Test for subgroup differences: $Chi^2 = 6.73$; d.f. = 1 (p = 0.009), $I^2 = 85.1\%$						



Fig. 3. Comparison of aspirin and clopidogrel versus aspirin or clopidogrel alone on ischemic stroke.

Study or Subgroup	A+C Events	A+C Total	A/C Events	A/C Total	Weight	Risk ratio M-H, Fixed, 95% CI
Short-term combination						
Bal Dit Sollier 2009	0	10	0	12		Not estimable
CARESS 2005	0	51	0	56		Not estimable
CHANCE 2013	8	2,584	8	2,586	18.8%	1.00 [0.38, 2.66]
CLAIR 2010	0	46	0	52		Not estimable
FASTER 2007	2	198	0	194	1.2%	4.90 [0.24, 101.40]
Subtotal (95% CI)		2,889		2,900	20.0%	1.23 [0.50, 3.04]
Total events	10		8			
Heterogeneity: $Chi^2 = 0.97$, d.f. = 1, (p = 0.32); $I^2 = 0\%$						
Test for overall effect: Z = 0.45 (p = 0.65)						
Long-term combination						
CHARISMAsub 2011	10	2,157	9	2,163	21.1%	1.11 [0.45, 2.74]
MATCH 2004	32	3,797	17	3,802	10.0%	1.88 [1.05, 3.39]
SPS3 2012	15	1,517	8	1,503	18.9%	1.86 [0.79, 4.37]
Subtotal (95% CI)		7,471		7,468	80.0%	1.67 [1.10, 2.56]
Total events	57		34			
Heterogeneity: $Chi^2 = 1.00$, d.f. = 2, (p = 0.61); $I^2 = 0\%$						
Test for overall effect: Z = 2.39 (p = 0.02)						
Total (95% CI)		10,360		10,368	100.0%	1.59 [1.08, 2.33]
Total events	67		42			
Heterogeneity: $Chi^2 = 2.44$, d.f. = 4, (p = 0.66); $I^2 = 0\%$						
Test for overall effect: Z = 2.36 (p = 0.02)						
Test for subgroup differences: $Chi^2 = 0.36$, d.f. = 1, (p = 0.55); $I^2 = 0\%$						



Fig. 4. Comparison of aspirin and clopidogrel versus aspirin or clopidogrel alone on hemorrhagic stroke.

Study or Subgroup	A+C Events	A+C Total	A/C Events	A/C Total	Weight	Risk ratio M-H, Random, 95% CI
Short-term combination						
Bal Dit Sollier 2009	0	10	1	12	0.2%	0.39 [0.02, 8.73]
CARESS 2005	1	51	4	56	0.4%	0.27 [0.03, 2.38]
CHANCE 2013	216	2,584	307	2,586	24.1%	0.70 [0.60, 0.83]
CLAIR 2010	0	46	2	52	0.2%	0.23 [0.01, 4.58]
FASTER 2007	17	198	23	194	4.5%	0.72 [0.40, 1.31]
Subtotal (95% CI)		2,889		2,900	29.3%	0.70 [0.60, 0.82]
Total events	234		337			
Heterogeneity: Tau^2 = 0; Chi^2 = 1.42, d.f. = 4, (p = 0.84); I^2 = 0%						
Test for overall effect: Z = 4.44 (p < 0.00001)						
Long-term combination						
CHARISMAsub 2011	174	2,157	207	2,163	21.1%	0.84 [0.70, 1.02]
MATCH 2004	477	3,797	490	3,802	29.3%	0.97 [0.87, 1.10]
SPS3 2012	153	1,517	174	1,503	20.1%	0.87 [0.71, 1.07]
Subtotal (95% CI)		7,471		7,468	70.7%	0.92 [0.84, 1.01]
Total events	804		871			
Heterogeneity: Tau^2 = 0.00; Chi^2 = 1.98, d.f. = 2, (p = 0.37); I^2 = 0%						
Test for overall effect: Z = 1.72 (p = 0.09)						
Total (95% CI)		10,360		10,368	100.0%	0.84 [0.73, 0.96]
Total events	1,038		1,208			
Heterogeneity: Tau^2 = 0.01; Chi^2 = 12.43, d.f. = 7, (p = 0.09); I^2 = 44%						
Test for overall effect: Z = 2.63 (p = 0.008)						
Test for subgroup differences: Chi^2 = 9.02, d.f. = 1, (p = 0.003); I^2 = 88.9%						



Fig. 5. Comparison of aspirin and clopidogrel versus aspirin or clopidogrel alone on stroke, MI, or vascular death.

Study or subgroup	A+C Events	A+C Total	A/C Events	A/C Total	Weight	Risk ratio M-H, Random, 95% CI
Short-term combination						
Bal Dit Sollier 2009	0	10	0	12		Not estimable
CARESS 2005	0	51	0	56		Not estimable
CHANCE 2013	7	2,584	8	2,586	7.0%	0.88 [0.32, 2.41]
CLAIR 2010	0	46	0	52		Not estimable
FASTER 2007	5	198	0	194	1.0%	10.78 [0.60, 193.62]
Subtotal (95% CI)		2,889		2,900	8.0%	2.17 [0.18, 25.71]
Total events	12		8			
Heterogeneity: Tau^2 = 2.22; Chi^2 = 2.82, d.f. = 1, (p = 0.09); I^2 = 65%						
Test for overall effect: Z = 0.61 (p = 0.54)						
Long-term combination						
CHARISMAsub 2011	92	2,157	61	2,163	29.6%	1.51 [1.10, 2.08]
MATCH 2004	169	3,797	71	3,802	32.7%	2.38 [1.81, 3.13]
SPS3 2012	105	1,517	56	1,503	29.7%	1.86 [1.35, 2.55]
Subtotal (95% CI)		7,471		7,468	92.0%	1.90 [1.46, 2.48]
Total events	366		188			
Heterogeneity: Tau^2 = 0.03; Chi^2 = 4.60, d.f. = 2, (p = 0.10); I^2 = 57%						
Test for overall effect: Z = 4.77 (p < 0.00001)						
Total (95% CI)		10,360		10,368	100.0%	1.83 [1.37, 2.45]
Total events	378		196			
Heterogeneity: Tau^2 = 0.05; Chi^2 = 8.29, d.f. = 4, (p = 0.08); I^2 = 52%						
Test for overall effect: Z = 4.08 (p < 0.0001)						
Test for subgroup differences: Chi^2 = 0.01, d.f. = 1, (p = 0.92); I^2 = 0%						



Fig. 6. Comparison of aspirin and clopidogrel versus aspirin or clopidogrel alone on major bleeding.