

1 **Full title: Associations of sedentary time with fat distribution in a high risk population**

2

3 **Authors and Affiliations**

4

5 **Joseph Henson¹, Charlotte L Edwardson¹, Bruno Morgan², Mark A Horsfield³, Danielle**
6 **H Bodicoat¹, Stuart JH Biddle⁴, Trish Gorely⁵, Myra A Nimmo⁶, Gerry P McCann⁷,**
7 **Kamlesh Khunti⁸, Melanie J Davies¹, Thomas Yates¹**

8

9 ¹ NIHR Leicester-Loughborough Diet, Lifestyle and Physical Activity Biomedical Research
10 Unit, UK and Diabetes Research Centre, College of Medicine, Biological Sciences and
11 Psychology, University of Leicester, UK.

12

13 ²Department of Cancer Studies and Molecular Medicine, College of Medicine, Biological
14 Sciences and Psychology, University of Leicester, UK.

15

16 ³Department of Cardiovascular Sciences, College of Medicine, Biological Sciences and
17 Psychology, University of Leicester, UK.

18

19 ⁴ NIHR Leicester-Loughborough Diet, Lifestyle and Physical Activity Biomedical Research
20 Unit, UK and School of Sport, Exercise and Health Sciences, Loughborough University, UK.

21

22 ⁵School of Sport, University of Stirling, UK.

23

24 ⁶College of Life & Environmental Sciences, University of Birmingham, UK.

25

26 ⁷Department of Cardiovascular Sciences, University of Leicester and the NIHR Leicester
27 Cardiovascular Biomedical Research Unit, Leicester, UK.

28

29 ⁸NIHR Collaborations for Leadership in Applied Health Research and Care (CLAHRC) East
30 Midlands, UK and Diabetes Research Centre, College of Medicine, Biological Sciences and
31 Psychology, University of Leicester, UK.

32

33 Main text word count = 3433; Number of Tables = 4; Number of Figures = 1; Number of
34 references =39

35

36 **Corresponding Author:**

37 Joseph Henson

38 Leicester Diabetes Centre

39 Leicester General Hospital

40 Leicester

41 LE5 4PW

42 UK

43

44 **Email address:**jjh18@le.ac.uk

45 **Tel:** +44 116 258 8599.

46 **Fax:** +44116 258 4053.

47

48

49

50

51 **Abstract**

52

53 *Purpose*

54 The effect of sedentary behaviour on regional fat deposition, independent of physical activity
55 remains equivocal. We examined the cross-sectional associations between objectively
56 measured sedentary time and markers of regional fat distribution (heart, liver, visceral,
57 subcutaneous and total body fat) in a population at a high risk of type 2 diabetes mellitus
58 (T2DM).

59

60 *Methods*

61 Participants were recruited from primary care to two diabetes prevention programmes.
62 Sedentary time (<25 counts per 15 seconds) was measured using Actigraph GT3X
63 accelerometers. Heart, liver, visceral, subcutaneous and total body fat were quantified using
64 magnetic resonance images (MRI). Fat volumes were calculated by multiplying the cross-
65 sectional areas of the fat-containing pixels by the slice thickness. The liver fat percentage was
66 measured using a representative region of interest created in the right lobe of the liver
67 avoiding the main portal veins. Linear regression models examined the association of
68 sedentary time with markers of regional fat deposition.

69

70 *Results*

71 Sixty-six participants (age=47.9±16.2years;male=50.0%) were included. Following
72 adjustment for several covariates, including glycaemia, whole body fat and moderate-to-
73 vigorous physical activity (MVPA), each 30 minutes of sedentary time was associated with
74 15.7cm³ higher heart fat ($p=0.008$), 1.2% higher liver fat ($p=0.026$) and 183.7cm³ higher
75 visceral fat ($p=0.039$).

76 Conclusion

77 This study provides new evidence suggesting that objectively measured sedentary behaviour
78 may have an independent association upon heart, liver and visceral fat in individuals at a high
79 risk of T2DM.

80

81

82 **Keywords;** Type 2 diabetes, Sedentary behaviour, High risk, Fat distribution, MRI, Primary
83 care

84

85

86

87

88

89

90

91

92

93

94

95

96

97

98

99

100

101 **Introduction**

102

103 Abdominal obesity is known to predispose individuals to cardiovascular disease (CVD) and
104 type 2 diabetes (T2DM), with regional fat deposits being postulated to be of greater
105 importance than overall adiposity in causing metabolic and cardiovascular disturbance (5,
106 31). Several studies have implicated pericardial and liver fat as particular pathogenic risk
107 factors (23,26), with excess visceral adiposity also being associated with dyslipidemia,
108 systemic inflammation, insulin resistance, T2DM and all-cause mortality (1,7,15,18).

109

110 Despite the well-documented positive effects of moderate-to-vigorous physical activity
111 (MVPA) on regional fat deposition (16), the associative role of sedentary behaviour,
112 independent of physical activity, is less well understood and the available literature
113 equivocal.

114

115 Over the past decade there has been an accumulation of epidemiological evidence from both
116 cross-sectional and prospective observational studies indicating that sedentary behaviour
117 (best conceptualised as any non-exercise sitting time (30)) may be independently associated
118 with several deleterious health outcomes, including T2DM, obesity, the metabolic syndrome,
119 cardiovascular disease and cardiovascular mortality (8,33,36,37). However, previous cross-
120 sectional and longitudinal studies conducted in the general population have shown no
121 association between sedentary behaviour and visceral fat accumulation in adults (20,25,29).
122 Although associations have previously been observed between objectively measured
123 sedentary time and pericardial fat (11,20), the relationships were either attenuated after
124 adjustment for MVPA (11) or MVPA was quantified using self report (20), thus raising
125 issues regarding response bias and poor levels of validity (27). It therefore remains unclear

126 whether objectively measured sedentary behaviour is associated with regional fat deposition,
127 independent of MVPA or total physical activity. Moreover, to our knowledge, there are
128 currently no reports examining the association between sedentary behaviour and liver fat.

129

130 It is also necessary to establish the association between sedentary behaviour and fat
131 distribution in those at high risk of chronic disease. Both national and international
132 recommendations and policies specify that chronic-disease prevention strategies should
133 include targeted interventions aimed at the identification and management of high risk
134 individuals (2). Moreover, sedentary time has been shown to be more strongly and adversely
135 associated with cardio-metabolic variables (including markers of adiposity) in high risk
136 individuals, (14) and those with established T2DM (3,4) after adjustment for MVPA and
137 other important confounders. Given that associations between sedentary time and markers of
138 adiposity (body mass index (BMI) and waist circumference) were weaker compared to other
139 cardio-metabolic variables (14), the association of sedentary behaviour may extend beyond
140 traditional measures of adiposity and may lie in the location of fat deposition. In particular,
141 within cells of non-adipose tissue that normally contain only small amounts of fat (ectopic
142 fat). Such ectopic depositions result in excess lipids being driven into alternative, non-
143 oxidative pathways, which in turn promotes metabolically relevant cellular dysfunction
144 (lipotoxicity).

145

146 The aim of this study, therefore, was to examine the association between objectively
147 measured sedentary time and heart, liver, visceral, subcutaneous and total body fat,
148 independent of MVPA and whole body fat in a population at high risk of T2DM.

149

150

151 **Methods**

152

153 **Subjects**

154

155 The present study reports a baseline convenience subsample (n=66) from the Walking Away
156 from Type 2 Diabetes Study (WA) and Project STAND (Sedentary Time **And** Diabetes).
157 When combined, the full cohort for both studies included 1,026 participants (WA=833,
158 Project STAND=193). Both of these diabetes prevention studies were conducted by the same
159 research group within the same geographical area (Leicestershire and South East Midlands,
160 United Kingdom (UK)) and baseline data collection was undertaken during 2010. All
161 measurements were performed by the same team of trained staff who followed identical
162 standard operating procedures. A detailed description of both trial methods have been
163 published elsewhere (38,39).

164

165 **Walking Away**

166

167 Participants (aged 30-74 years) were recruited from 10 primary care practices within the
168 Leicestershire region (city and county), UK. Individuals at high risk of impaired glucose
169 regulation (IGR) (composite of impaired glucose tolerance (IGT) and/or impaired fasting
170 glycaemia (IFG)) or T2DM were identified using a modified version of the automated
171 Leicester Risk Score, specifically designed to be administered in primary care (10). The
172 Morbidity, Information Query and Export Syntax (MIQUEST) programme was used to assess
173 medical records and rank individuals for diabetes risk using predefined weighted variables
174 commonly held on practice databases (age, gender, BMI, family history of T2DM and use of
175 antihypertensive medication). Those scoring within the 90th percentile in each practice were

176 invited to take part in the study. This approach has been shown to have good sensitivity and
177 specificity for identifying participants at a high risk of IGR (10).

178

179 Project STAND

180

181 Young adults who were at risk of developing T2DM were recruited from primary care
182 practices located across Leicestershire and the South East Midlands region. Practice
183 databases were searched for participants meeting the following inclusion criteria: a) aged 18-
184 40 years with a BMI in the obese range ($\geq 30\text{kg/m}^2$; $\geq 27.5\text{kg/m}^2$ for south Asians) or b) aged
185 18-40 years with a BMI in the overweight range $\geq 25\text{kg/m}^2$ ($\geq 23\text{kg/m}^2$ for south Asians) plus
186 one additional risk factor: a family history of T2DM or CVD, previous gestational diabetes,
187 polycystic ovarian syndrome, HbA1c $\geq 5.8\%$ or IGR (38).

188

189 Individuals were excluded from both studies if they were taking steroids or had previously
190 diagnosed T2DM. Written Informed consent was obtained from all eligible participants and
191 both studies gained full ethical and governance approval.

192

193 Covariates

194

195 Information on current smoking status, family history of T2DM, medication status and
196 ethnicity (coded according to census criteria) was obtained following an interview-
197 administered questionnaire with a health care professional. Waist circumference was
198 measured over light clothing between the lower rib margin and the iliac crest. Height and
199 weight (Tanita TBE 611, Tanita, West Drayton, UK) were obtained by trained staff according
200 to standard operating procedures. The subsequent values were used to compute BMI (kg/m^2).

201 Systolic and diastolic blood pressure (mmHg) were taken three times in succession and the
202 mean of the last two used for analysis.

203

204 Social deprivation was determined by assigning an Index of Multiple Deprivation (IMD)
205 score to the participant's resident area (based on postcode) (32). IMD scores are publically
206 available continuous measures of compound social and material deprivation. Areas are
207 ranked from least deprived to most deprived based upon several dimensions linked to health
208 outcomes (including; income, employment, education, living environment and health).

209

210 Venous blood samples were obtained following an overnight fast, and all assays were
211 measured in the same laboratory. Analysis was conducted by individuals blinded to the
212 patients' identity, using stable methods, standardised to external quality assurance values.
213 HbA1c was analysed using the Bio-Rad Variant II HPLC system (Bio-Rad Clinical
214 Diagnostics, Hemel Hempstead, UK) and total cholesterol was measured using standard
215 enzymatic techniques.

216

217 Quantification of sedentary time

218

219 All eligible participants were asked to wear a tri-axial accelerometer at the baseline visit
220 (Actigraph GT3X, Pensacola, FL, USA), for a minimum of seven consecutive days during
221 waking hours. These accelerometers translate raw accelerations into activity counts. Cut-
222 points modified by Troiano et al. were used to categorise an epoch as sedentary (<25 counts
223 per 15 seconds) or MVPA (≥ 505 counts per 15 seconds). These intensity thresholds were
224 calculated as a weighted average determined from previous treadmill or track walking studies
225 (34). Total physical activity volume represented the summation of counts within each epoch.

226 Non-wear time was defined as a minimum of 60 minutes of continuous zero counts and days
227 with at least 600 minutes of wear time were considered valid (13, 14). In order to be included
228 in the analysis, participants were required to have a minimum of four valid days (35).

229

230 A data analysis tool (KineSoft version 3.3.75, Kinesoft, New Brunswick, Canada;
231 www.kinesoft.org) was used to process the accelerometer data.

232

233 Measure of adiposity

234

235 Magnetic Resonance Imaging (MRI) was performed at Glenfield Hospital, Leicester, UK,
236 where heart, liver, visceral, subcutaneous and total body fat (includes liver, intra-abdominal,
237 subcutaneous and visceral fat) was quantified. MRI is a reliable modality for the assessment
238 of adipose tissue and is capable of measuring fat distribution with a high spatial resolution
239 (22).

240

241 Scanning was performed using either a 1.5 Tesla Avanto (WA) or a 3.0 Tesla Skyra system
242 (STAND) (Siemens Medical, Erlangen, Germany). Flexible body array coils were applied to
243 the thorax and abdomen for signal reception. For lipid volume quantification, a 2-point Dixon
244 gradient-echo pulse sequence was used to separate tissue water signal from lipid signal and to
245 create two separate image sets with signal intensity showing ‘fat’ and ‘water’ content (21). 3-
246 D images were acquired axially with 5 mm slice thickness and in-plane resolution of
247 1.56 mm, interpolated to 0.78 mm. The field of view was 500 mm (left-right) by 375 mm
248 (anterior-posterior). Images were acquired in three contiguous blocks, covering the thoracic,
249 abdominal and pelvic regions, with each block acquired in a breath-hold at full inspiration to
250 minimise motion-related artefacts and to negate changes in slice position. The acquisition

251 time for each block was 18s. All scans were performed by the same team of trained staff
252 according to standardised procedures.

253

254 Analysis of the MR images was performed using image analysis software produced in-house
255 (Java Image Manipulation, Version 7). All analysis was undertaken by the same researcher
256 who was blinded to the clinical, anthropometric and physical activity data.

257

258 For analysis, the 'fat' and 'water' images were mathematically combined to create a 'fat
259 percentage' image. Fat-containing pixels were then defined as those with a pixel intensity
260 between 51 and 99% (100% being due to image artefact). The images were reconstructed into
261 15 mm thick contiguous slices, from the top of the pulmonary trunk extending to the bottom
262 of the symphysis pubis. Volumes of interest for the whole body and heart were created by
263 outlining the perimeter of the body and heart respectively on each relevant slice using a
264 mouse-controlled pointer and excluding those pixels outside the structures. The region of
265 interest surrounding the heart included myocardial, epicardial (pericardial) and immediate
266 extra-pericardial (thoracic) fat.

267

268 The visceral (and retroperitoneal) fat was further separated, by outlining the abdominal and
269 chest wall muscles and excluding the pixels for the subcutaneous fat. The fat volume was
270 calculated automatically by multiplying the cross-sectional areas of the fat-containing pixels,
271 summed over all slices on which the tissue was outlined, by the slice thickness. This created
272 three fat volumes: total body fat, visceral fat from the top of the pulmonary trunk to the
273 bottom of the symphysis pubis, and the heart fat volume. The liver fat percentage was also
274 measured using a representative region of interest created in the right lobe of liver avoiding

275 the main portal veins. Subcutaneous fat was calculated by subtracting visceral fat from total
276 body fat.

277

278 Statistical Analysis

279

280 IBM SPSS Statistics v20.0 (Chicago, IL, USA) was used to conduct all statistical analyses.

281 Linear regression analysis was used on the combined study cohorts to examine the
282 independent association of sedentary time (independent variable), with various markers of
283 regional fat deposition (dependent variable). We display results per 30 minutes of sedentary
284 time for ease of interpretation.

285

286 Model 1 was adjusted for age (continuous), gender, ethnicity (white European/south
287 Asian/other), social deprivation (continuous), family history of T2DM (yes/no), smoking
288 status (current/ex/never smoked), total cholesterol, HbA1c, systolic blood pressure, blood
289 pressure medication (ACE inhibitors (yes/no)), beta-blockers (yes/no), lipid lowering
290 medication (yes/no), time accelerometer worn (average number of minutes per day) and
291 MVPA. We also undertook the same model, but adjusted for total physical activity volume
292 (counts per day) rather than MVPA given that others have suggested this mediates significant
293 associations between sedentary behaviour and metabolic health (24). In order to examine the
294 extent to which total adiposity attenuated these relationships, model 2 was further adjusted
295 for whole body fat. Models were assessed for normality and multi co-linearity was assessed
296 through the variance inflation factor (VIF). To further represent the strength of sedentary time
297 with markers of adiposity, variables were also examined as tertiles using analysis of
298 covariance procedures.

299

300 Significant observations were followed up with interaction terms to assess associations
301 between sedentary time and study, sex, level of MVPA, whole body fat and HbA1c. All
302 interactions were adjusted for the covariates listed in model 1.

303

304 Two-tailed p values of 0.05 or less were considered statistically significant for main effects.
305 $p < 0.1$ was considered significant for interactions. To allow for direct comparisons across fat
306 deposition markers, results of the generalised linear regression analysis are also presented as
307 the standardised beta co-efficient (β) \pm standard error(SE).

308

309 **Results**

310

311 Table 1 displays the demographic, anthropometric, MRI-derived and accelerometer
312 characteristics of included participants. In total, 32 participants from Project STAND
313 (age=33.1 \pm 6.0 years; male=34.4%) and 34 participants from WA (age=61.9 \pm 8.0 years;
314 male=64.7%) had valid measures of objective activity and MRI data.

315

316 There were no statistical differences ($p > 0.05$) in anthropometric, metabolic, and social
317 deprivation measures between participants who were included in this analysis vs. those not
318 included (did not undergo an MRI scan).

319

320 Model 1 illustrates the linear relationship between each 30 minute block of sedentary time
321 and markers of regional fat deposition. Following adjustment for various confounders,
322 including HbA1c, and MVPA, 30 minutes of sedentary time was associated with 20.5cm³
323 higher heart fat ((95% CI) 5.4, 35.6), 1.4% higher liver fat (0.3, 2.5) and a 409.2cm³ higher
324 visceral fat (127.6, 690.8). All significant associations seen in Model 1 persisted after further

325 adjustment for whole body fat in Model 2 (15.7cm³ higher heart fat ((95% CI) 0.5, 30.8),
326 1.2% higher liver fat (0.3, 2.3) and a 191.3cm³ higher visceral fat (2.7, 368.8).

327

328 No significant associations were observed for whole body and subcutaneous fat (Table 2).
329 Supplementary Table 1 also displays the overall associations (presented as standardised $\beta \pm$
330 SE) in the combined cohort for total sedentary time with MRI-derived markers of regional fat
331 deposition.

332

333 In order to provide visual representation of reported associations, figure 1 illustrates the
334 associations between total sedentary time and heart fat, liver fat and visceral fat when
335 examined as tertiles, after adjustment for the covariates listed above. Compared to those in
336 the lowest tertile of sedentary time, those in the highest tertile had, on average, 13.2cm³
337 higher heart fat ($p < 0.001$), 1.6% higher liver fat ($p < 0.001$) and a 556.3cm³ higher visceral fat
338 ($p < 0.001$).

339

340 Interaction analyses indicated a significant effect for study group with the older cohort (WA)
341 demonstrating stronger associations of sedentary time with visceral fat (presented as
342 unstandardised β (95% CI)) (WA = 800.0 (345.3, 1255.9) vs. STAND = 69.4 (-297.8, 436.6)
343 (p for interaction=0.010). Sex interactions also indicated that sedentary time had a larger
344 impact on visceral fat in males (male = 779.1 (171.4, 1386.9) vs. female = 133.4 (-269.0,
345 544.8) (p for interaction=0.049) (Table 3). No other significant interactions for associations
346 with measures of ectopic fat were observed for study group, sex, whole body fat, MVPA or
347 HbA1c level ($p > 0.1$).

348

349 The findings above were unaffected if waist circumference or BMI rather than whole body fat
350 was used in Model 2 (data not shown).

351

352 **Discussion**

353

354 This study conducted in individuals at high risk of T2DM, demonstrated that sedentary time
355 was associated with heart, liver and visceral fat, independent of measured confounders,
356 including glycaemia, whole body fat and MVPA. The findings from this study extend
357 previous cross-sectional results observed in the general population, by demonstrating the
358 association of objectively measured sedentary behaviour with markers of regional fat
359 deposition. To our knowledge, this is the first study to show associations between sedentary
360 time and liver, heart and visceral fat in a population with a high risk of chronic disease.

361

362 The observation that sedentary time is associated with liver fat, independent of adiposity, is a
363 novel finding and may suggest an independent association between sedentary time and liver
364 fat accumulation. Nevertheless, the associations observed between sedentary time and heart
365 and visceral fat are in contrast to the majority of (11,25,29), but not all (20) previous
366 literature, which has tended to show either weak or no associations. The discrepancy in
367 findings between studies may be partially explained by the fact that sedentary time has
368 previously been quantified using self-report (29), which has high measurement error (27), or
369 undertaken in generally healthy, low risk populations compared to the present analysis, which
370 specifically targeted individuals with a high risk of chronic disease and underlying metabolic
371 dysfunction.

372

373 Visceral, hepatic, and cardiac adiposity, rather than obesity *per se*, have all been causally
374 associated with glucose, insulin metabolism and subsequent metabolic dysfunction (6).
375 These mechanisms may induce multiple autocrine, paracrine and endocrine influences, which
376 include the pro-inflammatory cytokine response (28). Therefore, the associations observed for
377 regional and ectopic fat in the present study may help to partially explain the relatively strong
378 association between sedentary time and glucose metabolism consistently reported in those
379 with a high risk of, or diagnosed, T2DM (3,4,14). Although a causal link between sedentary
380 behaviour and differential regional and ectopic fat distribution has not been directly
381 elucidated, there is some supporting evidence. As this analysis and others have found only
382 relatively weak associations between sedentary behaviour and markers of overall adiposity
383 (4,13,14), it is likely that potential mechanisms are beyond total energy balance. One possible
384 candidate could be through the actions of lipoprotein lipase (LPL). Research using animal
385 models of sedentary behaviour have shown that muscle inactivity causes rapid and dramatic
386 reductions in LPL activity (12). In turn, it has been suggested that reductions in LPL mass
387 and activity may directly promote intra-abdominal visceral fat accumulation (17). Therefore,
388 if generalisable to humans, it may be plausible that muscle inactivity induced by
389 prolonged/chronic sitting related sedentary behaviour causes reductions in postural muscle
390 LPL activity. This in turn may help to promote the deposition of triglycerides into cells of
391 non-adipose tissue, fuelling the detrimental phenomenon of ectopic over-accumulation (31).
392 However, this potential mechanism lacks confirmation in human research and thus remains
393 suggestive rather than definitive. Our study supports the need for further experimental
394 research in humans focusing on lipid metabolism and distribution.

395

396 Sedentary time in the current study was shown to have a stronger association with visceral fat
397 in older, compared to younger adults and in males compared to females. Although visceral fat

398 is known to increase with age, clear sex dimorphisms also exist, largely due to anatomical
399 differences in adipose tissue deposition (6). For example, even after correcting for total body
400 fat mass, women have been shown to have a lower ratio of visceral adipose tissue to total
401 body fat mass compared to men (19). The underlying mechanisms driving these observations
402 are largely unknown; it is likely to be a complex phenotype that includes sex hormones and
403 adipose tissue storage dysfunction in several sites, including the heart and liver (6).
404 Therefore, the preliminary findings from this study further highlight the importance of
405 carefully considering the population under investigation in future experimental and
406 epidemiologic investigations.

407

408 The present study has several strengths: most notably the use of objective methodologies to
409 estimate exposures and outcomes in a high risk of T2DM population recruited through
410 primary care. This is particularly important as our population is representative of those who
411 are likely to be identified as being at high risk of type 2 diabetes mellitus within routine care
412 and referred on to available prevention programmes. Furthermore, all participants were from
413 the same geographical location, with similar risk, metabolic and physical activity profiles. All
414 measurements (including MRI scans) were also performed by the same team of trained staff,
415 following identical standard operating procedures.

416

417 However, the following limitations should be considered. Firstly, given the high risk nature
418 of the cohort, the results may have limited generalisability and the small sample size may
419 restrict the external validity of our findings. Secondly, the cross-sectional design limits
420 inference about the direction of causality between the sedentary variables and MRI markers;
421 reverse causality remains a possibility, particularly as the relationship between adiposity and
422 sedentary time may be bi-directional (9). It is also plausible that unmeasured lifestyle

423 variables (e.g. snacking, alcohol consumption) and pre-existing co-morbidities may have
424 confounded the observed relationships. Thirdly, cardiac images were un-gated and we were
425 unable to distinguish between pericardial, epicardial and pericoronary fat. However, it could
426 be argued that measuring whole heart fat reduces any potential bias, particularly related to
427 measurement in leaner individuals. Fourthly, accelerometers rely on categorising movement
428 (acceleration), as opposed to distinguishing between specific postures (sitting, lying and
429 standing behaviours), which may lead to an under-estimation of the true association between
430 sedentary time and markers of adiposity.

431

432 In conclusion, the present study provides new evidence suggesting that objectively measured
433 sedentary behaviour is associated with heart, liver and visceral fat in individuals at a high risk
434 of T2DM. Interestingly, since the associations remained after adjustment for whole body fat
435 and MVPA, it may suggest that sedentary behaviour is linked to selective depositions of fat
436 which cannot be fully explained by an increase in overall adiposity and may act via an
437 independent mechanism. However, given the limitations, more research is needed to
438 determine the distinct pathological effects of each type of fat and how these endpoints might
439 be associated with different behaviours, in particular sitting-related sedentary time.

440

441 **Acknowledgements**

442

443 The research was supported by The National Institute for Health Research Collaboration for
444 Leadership in Applied Health Research and Care - Leicestershire, Northamptonshire and
445 Rutland (NIHR CLAHRC – LNR), the University of Leicester Clinical Trials Unit and the
446 NIHR Leicester-Loughborough Diet, Lifestyle and Physical Activity Biomedical Research
447 Unit which is a partnership between University Hospitals of Leicester NHS Trust,

448 Loughborough University and the University of Leicester. The views expressed are those of
449 the author(s) and not necessarily those of the NHS, the NIHR or the Department of Health.
450 MRI scans (for the WA cohort only) were funded by Unilever Discover, UK. Project STAND
451 was funded by the Medical Research Council and National Prevention Research Initiative
452 funding partners (MRC Project no.91409). Dr G McCann is funded by a post-doctoral NIHR
453 fellowship.

454

455 **Conflict of Interest**

456

457 The authors declare no conflict of interest. The results of the present study do not constitute
458 endorsement by ACSM.

459

460 **References**

461

462 1. Calabro P, Yeh ET. Intra-abdominal adiposity, inflammation, and cardiovascular risk:
463 new insight into global cardiometabolic risk. *Curr Hypertens Rep.* 2008; 10(1):32-8.

464

465 2. Chatterton H, Younger T, Fischer A, Khunti K, Programme Development Group.
466 Risk identification and interventions to prevent type 2 diabetes in adults at high risk:
467 summary of NICE guidance. *BMJ.* 2012; 345:e4624.

468

469 3. Cooper AJ, Brage S, Ekelund U, Wareham NJ, Griffin SJ, Simmons RK. Association
470 between objectively assessed sedentary time and physical activity with metabolic risk factors
471 among people with recently diagnosed type 2 diabetes. *Diabetologia.* 2014; 57(1):73-82.

472

- 473 4. Cooper AR, Sebire S, Montgomery AA et al. Sedentary time, breaks in sedentary time
474 and metabolic variables in people with newly diagnosed type 2 diabetes. *Diabetologia*. 2012;
475 55(3):589-99.
- 476
- 477 5. Despres JP. Excess visceral adipose tissue/ectopic fat the missing link in the obesity
478 paradox? *J Am Coll Cardiol*. 2011; 57(19):1887-9.
- 479
- 480 6. Despres JP, Lemieux I. Abdominal obesity and metabolic syndrome. *Nature*. 2006;
481 444(7121):881-7.
- 482
- 483 7. Ebbert JO, Jensen MD. Fat depots, free fatty acids, and dyslipidemia. *Nutrients*. 2013;
484 5(2):498-508.
- 485
- 486 8. Edwardson CL, Gorely T, Davies MJ et al. Association of sedentary behaviour with
487 metabolic syndrome: a meta-analysis. *PLoS One*. 2012; 7(4):e34916.
- 488
- 489 9. Golubic R, Wijndaele K, Sharp SJ et al. Physical activity, sedentary time and gain in
490 overall and central body fat: 7-year follow-up of the ProActive trial cohort. *Int J Obes*
491 *(Lond)*. 2014.
- 492
- 493 10. Gray LJ, Davies MJ, Hiles S et al. Detection of impaired glucose regulation and/or
494 type 2 diabetes mellitus, using primary care electronic data, in a multiethnic UK community
495 setting. *Diabetologia*. 2012; 55(4):959-66.
- 496

- 497 11. Hamer M, Venuraju SM, Urbanova L, Lahiri A, Steptoe A. Physical activity,
498 sedentary time, and pericardial fat in healthy older adults. *Obesity (Silver Spring)*. 2012;
499 20(10):2113-7.
500
- 501 12. Hamilton MT, Hamilton DG, Zderic TW. Exercise physiology versus inactivity
502 physiology: an essential concept for understanding lipoprotein lipase regulation. *Exerc Sport*
503 *Sci Rev*. 2004; 32(4):161-6.
504
- 505 13. Healy GN, Matthews CE, Dunstan DW, Winkler EA, Owen N. Sedentary time and
506 cardio-metabolic biomarkers in US adults: NHANES 2003-06. *Eur Heart J*. 2011; 32(5):590-
507 7.
508
- 509 14. Henson J, Yates T, Biddle SJ et al. Associations of objectively measured sedentary
510 behaviour and physical activity with markers of cardiometabolic health. *Diabetologia*. 2013;
511 56(5):1012-20.
512
- 513 15. Hocking S, Samocha-Bonet D, Milner KL, Greenfield JR, Chisholm DJ. Adiposity
514 and insulin resistance in humans: the role of the different tissue and cellular lipid depots.
515 *Endocr Rev*. 2013; 34(4):463-500.
516
- 517 16. Kay SJ, Fiatarone Singh MA. The influence of physical activity on abdominal fat: a
518 systematic review of the literature. *Obes Rev*. 2006; 7(2):183-200.
519

- 520 17. Kobayashi J, Tashiro J, Murano S, Morisaki N, Saito Y. Lipoprotein lipase mass and
521 activity in post-heparin plasma from subjects with intra-abdominal visceral fat accumulation.
522 *Clin Endocrinol (Oxf)*. 1998; 48(4):515-20.
523
- 524 18. Kuk JL, Katzmarzyk PT, Nichaman MZ, Church TS, Blair SN, Ross R. Visceral fat is
525 an independent predictor of all-cause mortality in men. *Obesity (Silver Spring)*. 2006;
526 14(2):336-41.
527
- 528 19. Kuk JL, Lee S, Heymsfield SB, Ross R. Waist circumference and abdominal adipose
529 tissue distribution: influence of age and sex. *Am J Clin Nutr*. 2005; 81(6):1330-4.
530
- 531 20. Larsen BA, Allison MA, Kang E, Saad S, Laughlin GA, Araneta MR, Barrett-Connor
532 E, Wassel CL. Associations of Physical Activity and Sedentary Behavior with Regional Fat
533 Deposition. *Med Sci Sports Exerc*. 2013.
534
- 535 21. Le-Petross H, Kundra V, Szklaruk J, Wei W, Hortobagyi GN, Ma J. Fast three-
536 dimensional dual echo dixon technique improves fat suppression in breast MRI. *J Magn*
537 *Reson Imaging*. 2010; 31(4):889-94.
538
- 539 22. Machann J, Thamer C, Stefan N et al. Follow-up whole-body assessment of adipose
540 tissue compartments during a lifestyle intervention in a large cohort at increased risk for type
541 2 diabetes. *Radiology*. 2010; 257(2):353-63.
542

- 543 23. Mahabadi AA, Massaro JM, Rosito GA et al. Association of pericardial fat,
544 intrathoracic fat, and visceral abdominal fat with cardiovascular disease burden: the
545 Framingham Heart Study. *Eur Heart J*. 2009; 30(7):850-6.
546
- 547 24. Maher C, Olds T, Mire E, Katzmarzyk PT. Reconsidering the sedentary behaviour
548 paradigm. *PLoS One*. 2014; 9(1):e86403.
549
- 550 25. McGuire KA, Ross R. Incidental physical activity and sedentary behavior are not
551 associated with abdominal adipose tissue in inactive adults. *Obesity (Silver Spring)*. 2012;
552 20(3):576-82.
553
- 554 26. Nazare JA, Smith JD, Borel AL et al. Ethnic influences on the relations between
555 abdominal subcutaneous and visceral adiposity, liver fat, and cardiometabolic risk profile: the
556 International Study of Prediction of Intra-Abdominal Adiposity and Its Relationship With
557 Cardiometabolic Risk/Intra-Abdominal Adiposity. *Am J Clin Nutr*. 2012; 96(4):714-26.
558
- 559 27. Neilson HK, Robson PJ, Friedenreich CM, Csizmadi I. Estimating activity energy
560 expenditure: how valid are physical activity questionnaires? *Am J Clin Nutr*. 2008;
561 87(2):279-91.
562
- 563 28. Richardson VR, Smith KA, Carter AM. Adipose tissue inflammation: feeding the
564 development of type 2 diabetes mellitus. *Immunobiology*. 2013; 218(12):1497-504.
565

- 566 29. Saunders TJ, Tremblay MS, Despres JP, Bouchard C, Tremblay A, Chaput JP.
567 Sedentary behaviour, visceral fat accumulation and cardiometabolic risk in adults: a 6-year
568 longitudinal study from the Quebec Family Study. *PLoS One*. 2013; 8(1):e54225.
569
- 570 30. Sedentary Behaviour Research N. Letter to the editor: standardized use of the terms
571 "sedentary" and "sedentary behaviours". *Appl Physiol Nutr Metab*. 2012; 37(3):540-2.
572
- 573 31. Snel M, Jonker JT, Schoones J et al. Ectopic fat and insulin resistance:
574 pathophysiology and effect of diet and lifestyle interventions. *Int J Endocrinol*. 2012;
575 2012:983814.
576
- 577 32. The English Indices of Deprivation: Summary. *Available at:*
578 [http://webarchive.nationalarchives.gov.uk/20120919132719/http://www.communities.gov.uk/](http://webarchive.nationalarchives.gov.uk/20120919132719/http://www.communities.gov.uk/documents/communities/pdf/576659.pdf)
579 [documents/communities/pdf/576659.pdf](http://webarchive.nationalarchives.gov.uk/20120919132719/http://www.communities.gov.uk/documents/communities/pdf/576659.pdf). Accessed January 11, 2012. 2007
580
- 581 33. Thorp AA, Owen N, Neuhaus M, Dunstan DW. Sedentary behaviors and subsequent
582 health outcomes in adults a systematic review of longitudinal studies, 1996-2011. *Am J Prev*
583 *Med*. 2011; 41(2):207-15.
584
- 585 34. Troiano RP, Berrigan D, Dodd KW, Masse LC, Tilert T, McDowell M. Physical
586 activity in the United States measured by accelerometer. *Med Sci Sports Exerc*. 2008;
587 40(1):181-8.
588
- 589 35. Trost SG, McIver KL, Pate RR. Conducting accelerometer-based activity assessments
590 in field-based research. *Med Sci Sports Exerc*. 2005; 37(11 Suppl):S531-43.

591 36. Wijndaele K, O'Rowe G, Ekelund U et al. Increasing objectively measured sedentary
592 time increases clustered cardiometabolic risk: a 6 year analysis of the ProActive study.
593 *Diabetologia*. 2014; 57(2):305-12.

594

595 37. Wilmot EG, Edwardson CL, Achana FA et al. Sedentary time in adults and the
596 association with diabetes, cardiovascular disease and death: systematic review and meta-
597 analysis. *Diabetologia*. 2012; 55(11):2895-905.

598

599 38. Wilmot EG, Edwardson CL, Biddle SJ et al. Prevalence of diabetes and impaired
600 glucose metabolism in younger 'at risk' UK adults: insights from the STAND programme of
601 research. *Diabet Med*. 2013; 30(6):671-5.

602

603 39. Yates T, Davies MJ, Henson J et al. Walking away from type 2 diabetes: trial protocol
604 of a cluster randomised controlled trial evaluating a structured education programme in those
605 at high risk of developing type 2 diabetes. *BMC Fam Pract*. 2012; 13:46,2296-13-46.

606

607

608

609

610

611

612

613 **Legends to figures**

614 Figure 1A. Tertiles of sedentary time with heart fat.

615 Figure 1B. Tertiles of sedentary time with visceral fat.

616 Figure 1C. Tertiles of sedentary time with liver fat.

617 Tertiles of sedentary time with heart fat (Figure 1A), visceral fat (Figure 1B) and liver fat
618 (Figure 1C). Estimated marginal means are adjusted for age, gender, smoking status, family
619 history of T2DM, ethnicity, social deprivation, ACE inhibitors, beta blockers, lipid lowering
620 medication, systolic blood pressure, cholesterol, HbA1c, MVPA, time accelerometer worn
621 and whole body fat. Tertile cut-points for sedentary time were 9.6h and 10.9h per day.
622 Medians and ranges for tertile 1=8.8 h (7.7–9.6); tertile 2=10.3 h (9.6–10.8); tertile 3=11.8 h
623 (10.9–14.0). $p < 0.001$ for trend (**Figure 1A, Figure 1B, Figure 1C**). Bars represent mean and
624 error bars are 95% confidence intervals.

625 **Supplementary Tables**

626 Supplementary Table 1. Associations of total sedentary time with markers of MRI-derived
627 regional fat distribution when adjusted for either MVPA or total physical activity volume.