

| | Healthy control subjects | Patients with mild asthma (BTS step 1) | Patients with moderate asthma (BTS steps 2 and 3) | Patients with severe asthma (BTS steps 4 and 5) | <i>P</i> value† |
|--|--------------------------|--|---|---|--------------------|
| Number | 10 | 3 | 6 | 10 | - |
| Age (y), mean ± SEM | 33.8 ± 5.6 | 47.7 ± 9.1 | 28.8 ± 4.9 | 42.3 ± 3.7 | 0.134 |
| Sex (M/F) | 5/5 | 2/1 | 2/4 | 8/4 | 0.560 |
| Asthma duration (y), mean ± SEM | N/A | 33.0 ± 6.4 | 19.7 ± 2.6 | 24.6 ± 3.8 | 0.215 |
| Inhaled corticosteroid dose (µg), beclomethasone equivalents ‡, mean ± SEM | 0 | 0 | 1067.0 ± 133.3* | 1655.0 ± 202.0* | < 0.0001 |
| No. at BTS step 5 | 0 | 0 | 0 | 3 | 0.154 |
| No. taking long-acting β-agonist | 0 | 0 | 6 | 12 | < 0.0001 |
| Exacerbations in last year, median (IQR) | N/A | 0 | 0 (0 - 2.5) | 3.5 (1.3 - 6.8) | 0.010 |
| Mean daytime symptom score, median (IQR) | N/A | 0.1 (0 - 0.3) | 0.2 (0 - 0.5) | 1.2 (0.1 - 1.8) | 0.074 |
| Mean daily night-time symptom score, median (IQR) | N/A | 0 (0 - 1) | 0 (0 - 0.4) | 1.1 (0 - 1.9) | 0.156 |
| Reliever use per week, median (IQR) | N/A | 2.0 (0 - 34.5) | 3.0 (0 - 19.9) | 35 (19.8 - 49.4) | 0.017 |
| Sputum eosinophil count (%), median (IQR) | 0 (0 - 0.8) | 4.1 (3.3 - 9.8) | 3.1 (1.2 - 10.9) | 2.5 (0.5 - 14.2)* | 0.007 |
| PEF amplitude (% of mean), mean ± SEM | N/A | 35.2 ± 12.6 | 21.7 ± 7.9 | 41.3 ± 9.3 | 0.410 |
| FEV ₁ (% predicted), mean ± SEM | 99.7 ± 4.1 | 88.7 ± 8.7 | 93.0 ± 5.9 | 75.3 ± 4.9 | 0.013 |
| FEV ₁ /FVC ratio (%), mean ± SEM | 79.7 ± 3.1 | 72.0 ± 2.5 | 75.8 ± 4.1 | 66.7 ± 3.4 | 0.058 |
| PC ₂₀ methacholine (mg/ml), median (IQR) | >16 | 0 (0 - 0.2)* | 0.6 (0.1 - 6.5) | 0.1 (0 - 0.9)* | 0.001 |
| Serum IgE (kU/L), median (IQR) | 38.4 (21.3 - 114.5) | 76.4 (74.8 - 806.0) | 206.0 (118.0 - 1417.0) | 86.4 (39.2 - 552.5) | 0.081 |
| No. with positive skin prick test response | 10 | 3 | 6 | 10 | 0.651 |
| No. with positive skin prick test response to <i>Aspergillus fumigatus</i> | 0 | 1 | 2 | 4 | 0.230 |

Table 6-1: Clinical characteristics of donors recruited for collection of bronchial biopsy specimens.

BTS, British Thoracic Society; N/A, not applicable; PEF, peak expiratory flow; U/L, units per litre. ‡ Ratio for budesonide Turbuhaler calculated as 1.5; † Statistical significance across asthma groups unless otherwise stated; * Statistical significance in comparison with healthy control group; Bold text highlights statistically significant *P* values. Data are presented as mean ± SEM or median (IQR, interquartile range).