


5-Amino-1-phenyl-1H-pyrazole-4-carboxylic acid

Muhammad Zia-ur-Rehman, Mark R. J. Elsegood, Nosheen Akbar and
Rahman Shah Zaib Saleem

Acta Cryst. (2008). **E64**, o1312–o1313

This article is distributed under the terms of the Creative Commons Attribution Licence <http://creativecommons.org/licenses/by/2.0/uk/legalcode>, which permits unrestricted use, distribution, and reproduction in any medium, provided the original authors and source are cited.



Acta Crystallographica Section E: Structure Reports Online is the IUCr's highly popular open-access structural journal. It provides a simple and easily accessible publication mechanism for the growing number of inorganic, metal-organic and organic crystal structure determinations. The electronic submission, validation, refereeing and publication facilities of the journal ensure very rapid and high-quality publication, whilst key indicators and validation reports provide measures of structural reliability. In 2007, the journal published over 5000 structures. The average publication time is less than one month.

Crystallography Journals **Online** is available from journals.iucr.org

5-Amino-1-phenyl-1*H*-pyrazole-4-carboxylic acid

Muhammad Zia-ur-Rehman,^{a*} Mark R. J. Elsegood,^b Nosheen Akbar^c and Rahman Shah Zaib Saleem^d

^aApplied Chemistry Research Centre, PCSIR Laboratories Complex, Lahore 54600, Pakistan, ^bChemistry Department, Loughborough University, Loughborough LE11 3TU, England, ^cCentre for High Energy Physics, University of the Punjab, Lahore 54590, Pakistan, and ^dDepartment of Chemistry, Michigan State University, East Lansing, Michigan 48824, USA
Correspondence e-mail: rehman_pcsir@hotmail.com


Received 5 June 2008; accepted 17 June 2008

Key indicators: single-crystal X-ray study; $T = 150$ K; mean $\sigma(\text{C}-\text{C}) = 0.002$ Å; R factor = 0.043; wR factor = 0.117; data-to-parameter ratio = 19.3.

In the molecule of the title compound, $\text{C}_{10}\text{H}_9\text{N}_3\text{O}_2$, the pyrazole ring is approximately coplanar with the amino and carboxyl groups. The phenyl group is twisted by 48.13 (3)° relative to this plane. An intramolecular $\text{N}-\text{H}\cdots\text{O}$ hydrogen bond stabilizes the planar conformation of the molecule. The molecules are linked into two-dimensional sheets by two strong intermolecular $\text{N}-\text{H}\cdots\text{N}$ and $\text{O}-\text{H}\cdots\text{O}$ hydrogen bonds. The latter forms the classic carboxylic acid dimer motif.

Related literature

For related literature, see: Baroni & Kovyrzina (1961); Baraldi *et al.* (1998); Bruno *et al.* (1990); Chen & Li (1998); Cottineau *et al.* (2002); Dardari *et al.* (2006); Jin *et al.* (2004); Li *et al.* (2006); Londershausen (1996); Mishra *et al.* (1998); Neunhoeffer *et al.* (1959); Siddiqui *et al.* (2007); Smith *et al.* (2001); Zhong *et al.* (2006); Zia-ur-Rehman *et al.* (2005, 2006).



Experimental

Crystal data

$\text{C}_{10}\text{H}_9\text{N}_3\text{O}_2$
 $M_r = 203.20$
Monoclinic, $P2_1/n$
 $a = 3.7937$ (5) Å

$b = 21.613$ (3) Å
 $c = 11.1580$ (16) Å
 $\beta = 92.170$ (2)°
 $V = 914.2$ (2) Å³

$Z = 4$
Mo $K\alpha$ radiation
 $\mu = 0.11$ mm⁻¹

$T = 150$ (2) K
 $0.28 \times 0.10 \times 0.07$ mm

Data collection

Bruker APEXII CCD diffractometer
Absorption correction: multi-scan (SADABS; Sheldrick, 2007)
 $T_{\min} = 0.971$, $T_{\max} = 0.993$

10482 measured reflections
2800 independent reflections
1967 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.034$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.043$
 $wR(F^2) = 0.116$
 $S = 1.02$
2800 reflections
145 parameters

H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\text{max}} = 0.34$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.27$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{N4}-\text{H4A}\cdots\text{O3}$	0.903 (18)	2.136 (18)	2.8233 (16)	132.3 (14)
$\text{N4}-\text{H4B}\cdots\text{N3}^i$	0.876 (18)	2.239 (18)	3.0087 (17)	146.5 (15)
$\text{O4}-\text{H4}\cdots\text{O3}^{ii}$	0.92 (2)	1.70 (2)	2.6189 (14)	178.4 (19)

Symmetry codes: (i) $x - \frac{1}{2}, -y + \frac{1}{2}, z + \frac{1}{2}$; (ii) $-x, -y, -z$.

Data collection: APEX2 (Bruker, 2006); cell refinement: SAINT (Bruker, 2006); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: SHELXTL (Sheldrick, 2008); software used to prepare material for publication: SHELXTL and local programs.

The authors are grateful to the Pakistan Council of Scientific and Industrial Research Laboratories Complex, Lahore, for the provision of necessary chemicals.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BT2722).

References

- Baraldi, P. G., Manfredini, S., Romagnoli, R., Stevanato, L., Zaid, A. N. & Manservigi, R. (1998). *Nucleosides Nucleotides*, **17**, 2165–2171.
Baroni, E. E. & Kovyrzina, K. A. (1961). *Zh. Obshch. Khim.* **31**, 1641–1645.
Bruker (2006). APEX2 and SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.
Bruno, O., Bondavalli, F., Ranise, A., Schenone, P., Losasso, C., Cilenti, L., Matera, C. & Marmo, E. (1990). *Il Farmaco*, **45**, 147–66.
Chen, H. S. & Li, Z. M. (1998). *Chem. J. Chin. Univ.* **19**, 572–576.
Cottineau, B., Toto, P., Marot, C., Pipaud, A. & Chenault, J. (2002). *Bioorg. Med. Chem.* **12**, 2105–2108.
Dardari, Z., Lemrani, M., Sebban, A., Bahloul, A., Hassair, M., Kitane, S., Berrada, M. & Boudouma, M. (2006). *Arch. Pharm.* **339**, 291–298.
Jin, Z.-M., Li, L., Li, M.-C., Hu, M.-L. & Shen, L. (2004). *Acta Cryst.* **C60**, o642–o643.
Li, S.-Y., Zhong, P., Hu, M.-L., Luo, Y. & Li, J.-H. (2006). *Acta Cryst.* **E62**, o3821–o3822.
Londershausen, M. (1996). *Pestic. Sci.* **48**, 269–274.
Mishra, P. D., Wahidullah, S. & Kamat, S. Y. (1998). *Indian J. Chem. Sect. B*, **37**, 199.
Neunhoeffer, O., Alsdorf, G. & Ulrich, H. (1959). *Chem. Ber.* **92**, 252–256.
Sheldrick, G. M. (2007). *SADABS*. University of Göttingen, Germany.
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
Siddiqui, H. L., Zia-ur-Rehman, M., Ahmad, N., Weaver, G. W. & Lucas, P. D. (2007). *Chem. Pharm. Bull.* **55**, 1014–1017.

Smith, S. R., Denhardt, G. & Terminelli, C. (2001). *Eur. J. Pharmacol.* **432**, 107–119.
Zhong, P., Zhang, X.-H., Xiao, H.-P. & Hu, M.-L. (2006). *Acta Cryst.* **E62**, o513–o515.

Zia-ur-Rehman, M., Choudary, J. A. & Ahmad, S. (2005). *Bull. Korean Chem. Soc.* **26**, 1771–1175.
Zia-ur-Rehman, M., Choudary, J. A., Ahmad, S. & Siddiqui, H. L. (2006). *Chem. Pharm. Bull.* **54**, 1175–1178.

supplementary materials

Acta Cryst. (2008). E64, o1312-o1313 [doi:10.1107/S1600536808018394]

5-Amino-1-phenyl-1*H*-pyrazole-4-carboxylic acid

M. Zia-ur-Rehman, M. R. J. Elsegood, N. Akbar and R. Shah Zaib Saleem

Comment

Pyrazole and its derivatives are known as heterocyclic compounds, having a wide range of biological activities. Some pyrazoles have been reported to possess significant antiarrhythmic and sedative (Bruno *et al.*, 1990), hypoglycemic (Cot-tineau *et al.*, 2002), antiviral (Baraldi *et al.*, 1998), and pesticidal (Londershausen, 1996) activities. Some of their derivatives have also been successfully tested for their antifungal (Chen & Li, 1998), antihistaminic (Mishra *et al.*, 1998) and anti-inflammatory (Smith *et al.*, 2001) activities. In addition, they have also been used as ligands to investigate the structure–activity relationship of the active site of metalloproteins (Dardari *et al.*, 2006) and for the preparation of some commercially important dyestuffs (Baroni & Kovyrzina, 1961; Neunhoeffer *et al.*, 1959).

As part of our ongoing research on the synthesis and biological evaluation of heterocyclic compounds (Zia-ur-Rehman *et al.*, 2005, 2006; Siddiqui *et al.*, 2007), the crystal structure of the title compound, (I), was determined. In (I), the pyrazole ring is approximately co-planar with the amino and carboxylic acid groups. The C—N bond lengths in the pyrazole ring are 1.3146 (18) and 1.3530 (16) Å, which are shorter than a typical C—N single bond length of 1.443 Å, but longer than a typical C—N bond length of 1.269 Å (Jin *et al.*, 2004), indicating electron delocalization. Most of the bond lengths and angles in *N*-phenylpyrazole group are in consistent with those in similar molecules (Li *et al.*, 2006; Zhong *et al.*, 2006). Each molecule exhibits an intramolecular N—H···O hydrogen bond which stabilizes the planar conformation and is linked to an adjacent one through head-to-tail pairs of O—H···O intermolecular interactions giving rise to dimeric motifs typical for carboxylic acids. Neighbouring dimers are further arranged into two-dimensional sheets in the (101) plane through N—H···N interactions (Fig.2).

Experimental

A mixture of 5-amino-1-phenyl-1*H*-pyrazole-4-carboxylic acid, ethyl ester (2.312 g; 10.0 mmoles), potassium hydroxide (1.12 g; 20 mmoles) and ethanol (25 ml) was refluxed for two hours. The reaction mixture was poured into ice cooled water and acidified with dilute hydrochloric acid to Congo Red. The precipitated solids were collected by filtration, washed and dried. Crystals suitable for single-crystal X-ray diffraction were grown by slow evaporation of solution of the title compound in a mixture of ethanol and water (85:15); m.p. 460 K; yield: 68%.

Refinement

H atoms bound to C were placed in geometric positions (C—H distance = 0.95 Å) using a riding model. H atoms on N and O had coordinates freely refined. U_{iso} values were set to $1.2U_{\text{eq}}$ ($1.5U_{\text{eq}}$ for OH).

Figures




Fig. 1. The asymmetric unit of the title compound. Displacement ellipsoids are drawn at the 50% probability level.




Fig. 2. Perspective view of the crystal packing showing hydrogen-bond interactions (dashed lines). H atoms not involved in hydrogen bonding have been omitted for clarity.

5-Amino-1-phenyl-1H-pyrazole-4-carboxylic acid

Crystal data

$C_{10}H_9N_3O_2$

$M_r = 203.20$

Monoclinic, $P2_1/n$

Hall symbol: $-P\ 2_1/n$

$a = 3.7937\ (5)\ \text{\AA}$

$b = 21.613\ (3)\ \text{\AA}$

$c = 11.1580\ (16)\ \text{\AA}$

$\beta = 92.170\ (2)^\circ$

$V = 914.2\ (2)\ \text{\AA}^3$

$Z = 4$

$F_{000} = 424$

$D_x = 1.476\ \text{Mg m}^{-3}$

Melting point: 460 K

Mo $K\alpha$ radiation

$\lambda = 0.71073\ \text{\AA}$

Cell parameters from 2299 reflections

$\theta = 3.4\text{--}29.6^\circ$

$\mu = 0.11\ \text{mm}^{-1}$

$T = 150\ (2)\ \text{K}$

Block, colourless

$0.28 \times 0.10 \times 0.07\ \text{mm}$

Data collection

Bruker APEXII CCD
diffractometer

Radiation source: fine-focus sealed tube

Monochromator: graphite

$T = 150\ (2)\ \text{K}$

ω rotation with narrow frames scans

Absorption correction: multi-scan
(SADABS; Sheldrick, 2007)

$T_{\min} = 0.971$, $T_{\max} = 0.993$

10482 measured reflections

2800 independent reflections

1967 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.034$

$\theta_{\text{max}} = 30.6^\circ$

$\theta_{\text{min}} = 1.9^\circ$

$h = -5 \rightarrow 5$

$k = -30 \rightarrow 30$

$l = -15 \rightarrow 15$

Refinement

Refinement on F^2

Least-squares matrix: full

Secondary atom site location: difference Fourier map

Hydrogen site location: geom except NH & OH co-ords freely refined

$$R[F^2 > 2\sigma(F^2)] = 0.043$$

$$wR(F^2) = 0.116$$

$$S = 1.02$$

2800 reflections

145 parameters

Primary atom site location: structure-invariant direct methods

H atoms treated by a mixture of independent and constrained refinement

$$w = 1/[\sigma^2(F_o^2) + (0.0521P)^2 + 0.3077P]$$

$$\text{where } P = (F_o^2 + 2F_c^2)/3$$

$$(\Delta/\sigma)_{\max} < 0.001$$

$$\Delta\rho_{\max} = 0.34 \text{ e } \text{\AA}^{-3}$$

$$\Delta\rho_{\min} = -0.27 \text{ e } \text{\AA}^{-3}$$

Extinction correction: none

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.0569 (4)	0.34024 (6)	-0.20989 (12)	0.0196 (3)
H1	-0.0345	0.3206	-0.2807	0.023*
C2	0.0428 (4)	0.40410 (7)	-0.19887 (13)	0.0242 (3)
H2	-0.0548	0.4284	-0.2628	0.029*
C3	0.1711 (4)	0.43247 (7)	-0.09461 (14)	0.0262 (3)
H3	0.1623	0.4762	-0.0873	0.031*
C4	0.3122 (4)	0.39693 (7)	-0.00110 (13)	0.0235 (3)
H4C	0.3958	0.4165	0.0707	0.028*
C5	0.3323 (4)	0.33300 (6)	-0.01161 (12)	0.0196 (3)
H5	0.4313	0.3087	0.0522	0.024*
C6	0.2054 (3)	0.30513 (6)	-0.11682 (11)	0.0168 (3)
N2	0.2312 (3)	0.23989 (5)	-0.13188 (9)	0.0172 (2)
N3	0.3587 (3)	0.21589 (5)	-0.23810 (10)	0.0205 (3)
C7	0.3374 (4)	0.15556 (6)	-0.22523 (12)	0.0200 (3)
H7	0.4084	0.1268	-0.2839	0.024*
C8	0.1976 (4)	0.13829 (6)	-0.11488 (11)	0.0175 (3)
C9	0.1300 (3)	0.19438 (6)	-0.05734 (11)	0.0163 (3)
N4	-0.0209 (3)	0.20277 (6)	0.04849 (10)	0.0221 (3)
H4A	-0.068 (5)	0.1668 (8)	0.0857 (15)	0.027*
H4B	-0.028 (5)	0.2377 (8)	0.0881 (16)	0.027*
C10	0.1212 (4)	0.07860 (6)	-0.06630 (12)	0.0198 (3)
O3	-0.0099 (3)	0.07239 (4)	0.03332 (9)	0.0247 (2)
O4	0.1962 (3)	0.03107 (5)	-0.13546 (9)	0.0289 (3)

supplementary materials

H4 0.132 (5) -0.0050 (10) -0.0983 (17) 0.043*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0207 (7)	0.0229 (7)	0.0153 (6)	0.0004 (5)	0.0039 (5)	0.0023 (5)
C2	0.0263 (7)	0.0229 (7)	0.0239 (7)	0.0054 (6)	0.0068 (6)	0.0071 (5)
C3	0.0298 (8)	0.0172 (6)	0.0324 (8)	-0.0002 (6)	0.0113 (6)	-0.0003 (5)
C4	0.0257 (7)	0.0229 (7)	0.0222 (7)	-0.0043 (5)	0.0052 (5)	-0.0049 (5)
C5	0.0210 (7)	0.0208 (6)	0.0171 (6)	-0.0009 (5)	0.0011 (5)	-0.0005 (5)
C6	0.0183 (6)	0.0163 (6)	0.0161 (6)	-0.0002 (5)	0.0046 (5)	0.0009 (5)
N2	0.0238 (6)	0.0162 (5)	0.0119 (5)	-0.0004 (4)	0.0040 (4)	0.0001 (4)
N3	0.0285 (6)	0.0208 (6)	0.0126 (5)	0.0010 (5)	0.0069 (4)	-0.0007 (4)
C7	0.0266 (7)	0.0192 (6)	0.0144 (6)	0.0006 (5)	0.0042 (5)	-0.0011 (5)
C8	0.0227 (6)	0.0164 (6)	0.0136 (6)	-0.0001 (5)	0.0029 (5)	0.0000 (4)
C9	0.0195 (6)	0.0164 (6)	0.0133 (6)	-0.0004 (5)	0.0013 (5)	0.0009 (4)
N4	0.0348 (7)	0.0168 (5)	0.0153 (5)	-0.0022 (5)	0.0095 (5)	-0.0009 (4)
C10	0.0256 (7)	0.0173 (6)	0.0165 (6)	-0.0001 (5)	0.0034 (5)	-0.0009 (5)
O3	0.0384 (6)	0.0179 (5)	0.0185 (5)	-0.0016 (4)	0.0096 (4)	0.0006 (4)
O4	0.0493 (7)	0.0158 (5)	0.0228 (5)	-0.0021 (5)	0.0157 (5)	-0.0022 (4)

Geometric parameters (\AA , $^\circ$)

C1—C2	1.387 (2)	N2—N3	1.3968 (15)
C1—C6	1.3883 (18)	N3—C7	1.3146 (18)
C1—H1	0.9500	C7—C8	1.4092 (17)
C2—C3	1.387 (2)	C7—H7	0.9500
C2—H2	0.9500	C8—C9	1.4001 (17)
C3—C4	1.387 (2)	C8—C10	1.4331 (18)
C3—H3	0.9500	C9—N4	1.3438 (16)
C4—C5	1.3891 (19)	N4—H4A	0.903 (18)
C4—H4C	0.9500	N4—H4B	0.876 (18)
C5—C6	1.3891 (18)	C10—O3	1.2423 (16)
C5—H5	0.9500	C10—O4	1.3221 (16)
C6—N2	1.4239 (16)	O4—H4	0.92 (2)
N2—C9	1.3530 (16)		
C2—C1—C6	119.58 (13)	C9—N2—C6	128.69 (11)
C2—C1—H1	120.2	N3—N2—C6	119.69 (10)
C6—C1—H1	120.2	C7—N3—N2	104.53 (10)
C3—C2—C1	120.06 (13)	N3—C7—C8	112.64 (12)
C3—C2—H2	120.0	N3—C7—H7	123.7
C1—C2—H2	120.0	C8—C7—H7	123.7
C4—C3—C2	119.97 (13)	C9—C8—C7	104.64 (11)
C4—C3—H3	120.0	C9—C8—C10	124.25 (12)
C2—C3—H3	120.0	C7—C8—C10	131.08 (12)
C3—C4—C5	120.55 (13)	N4—C9—N2	125.61 (12)
C3—C4—H4C	119.7	N4—C9—C8	127.68 (12)
C5—C4—H4C	119.7	N2—C9—C8	106.64 (11)

C4—C5—C6	118.97 (13)	C9—N4—H4A	112.7 (11)
C4—C5—H5	120.5	C9—N4—H4B	125.6 (11)
C6—C5—H5	120.5	H4A—N4—H4B	120.0 (16)
C1—C6—C5	120.86 (13)	O3—C10—O4	122.72 (12)
C1—C6—N2	118.73 (12)	O3—C10—C8	121.96 (12)
C5—C6—N2	120.40 (12)	O4—C10—C8	115.31 (12)
C9—N2—N3	111.54 (10)	C10—O4—H4	109.2 (12)
C6—C1—C2—C3	-1.1 (2)	N3—C7—C8—C9	0.05 (16)
C1—C2—C3—C4	-0.3 (2)	N3—C7—C8—C10	178.11 (14)
C2—C3—C4—C5	1.2 (2)	N3—N2—C9—N4	-176.22 (12)
C3—C4—C5—C6	-0.7 (2)	C6—N2—C9—N4	0.3 (2)
C2—C1—C6—C5	1.6 (2)	N3—N2—C9—C8	0.98 (15)
C2—C1—C6—N2	-177.34 (12)	C6—N2—C9—C8	177.55 (13)
C4—C5—C6—C1	-0.7 (2)	C7—C8—C9—N4	176.51 (14)
C4—C5—C6—N2	178.20 (12)	C10—C8—C9—N4	-1.7 (2)
C1—C6—N2—C9	-130.01 (14)	C7—C8—C9—N2	-0.62 (15)
C5—C6—N2—C9	51.1 (2)	C10—C8—C9—N2	-178.85 (13)
C1—C6—N2—N3	46.33 (17)	C9—C8—C10—O3	-0.8 (2)
C5—C6—N2—N3	-132.62 (13)	C7—C8—C10—O3	-178.56 (14)
C9—N2—N3—C7	-0.93 (15)	C9—C8—C10—O4	178.38 (13)
C6—N2—N3—C7	-177.85 (12)	C7—C8—C10—O4	0.7 (2)
N2—N3—C7—C8	0.51 (16)		

Hydrogen-bond geometry (Å, °)

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
N4—H4A...O3	0.903 (18)	2.136 (18)	2.8233 (16)	132.3 (14)
N4—H4B...N3 ⁱ	0.876 (18)	2.239 (18)	3.0087 (17)	146.5 (15)
O4—H4...O3 ⁱⁱ	0.92 (2)	1.70 (2)	2.6189 (14)	178.4 (19)

Symmetry codes: (i) $x-1/2, -y+1/2, z+1/2$; (ii) $-x, -y, -z$.

Fig. 1




Fig. 2

