

This item was submitted to Loughborough's Institutional Repository (<https://dspace.lboro.ac.uk/>) by the author and is made available under the following Creative Commons Licence conditions.



CC creative commons  
COMMONS DEED

**Attribution-NonCommercial-NoDerivs 2.5**

**You are free:**

- to copy, distribute, display, and perform the work

**Under the following conditions:**

 **Attribution.** You must attribute the work in the manner specified by the author or licensor.

 **Noncommercial.** You may not use this work for commercial purposes.

 **No Derivative Works.** You may not alter, transform, or build upon this work.

- For any reuse or distribution, you must make clear to others the license terms of this work.
- Any of these conditions can be waived if you get permission from the copyright holder.

**Your fair use and other rights are in no way affected by the above.**

This is a human-readable summary of the [Legal Code \(the full license\)](#).

[Disclaimer](#) 

For the full text of this licence, please go to:  
<http://creativecommons.org/licenses/by-nc-nd/2.5/>

---

---

**ERRATA**

---

---

**Erratum: Colossal magneto-optical conductivity in doped manganites**  
**[Phys. Rev. B 60, 6215 (1999)]**

A. S. Alexandrov and A. M. Bratkovsky

There is a typo in the comment following Eq. (9) for  $n_c(h)$  close to the ferromagnetic phase transition on the paramagnetic side. It should read:

Here we assume that  $w \gg t_c$  (in the opposite limit,  $w \ll t_c$ , the preexponential factor will be 2 instead of  $t_c/w$ ). The factors result from assuming that the total number of states in the bipolaron (impurity) band equals the doping level  $x$ . If we were to normalize to unity instead of  $x$ , as in Ref. 1, we would obtain the prefactors  $(t_c/w)\sqrt{x/2}$  and  $\sqrt{2x}$ , respectively.

This correction has no effect on our results and discussion.

<sup>1</sup>A. S. Alexandrov and A. M. Bratkovsky, Phys. Rev. Lett. **82**, 141 (1999).