
This item was submitted to [Loughborough's Research Repository](#) by the author.
Items in Figshare are protected by copyright, with all rights reserved, unless otherwise indicated.

N-(3-Methoxyphenyl)-tert-butanesulfinamide

PLEASE CITE THE PUBLISHED VERSION

PUBLISHER

© International Union of Crystallography

VERSION

VoR (Version of Record)

LICENCE

CC BY-NC-ND 4.0

REPOSITORY RECORD

Datta, Mrityunjoy, Alan J. Buglass, and Mark R.J. Elsegood. 2019. "N-(3-methoxyphenyl)-tert-butanesulfinamide". figshare. <https://hdl.handle.net/2134/5712>.

This item was submitted to Loughborough's Institutional Repository (<https://dspace.lboro.ac.uk/>) by the author and is made available under the following Creative Commons Licence conditions.



CC creative commons
COMMONS DEED

Attribution-NonCommercial-NoDerivs 2.5

You are free:

- to copy, distribute, display, and perform the work

Under the following conditions:

 **Attribution.** You must attribute the work in the manner specified by the author or licensor.

 **Noncommercial.** You may not use this work for commercial purposes.

 **No Derivative Works.** You may not alter, transform, or build upon this work.

- For any reuse or distribution, you must make clear to others the license terms of this work.
- Any of these conditions can be waived if you get permission from the copyright holder.

Your fair use and other rights are in no way affected by the above.

This is a human-readable summary of the [Legal Code \(the full license\)](#).

[Disclaimer](#) 

For the full text of this licence, please go to:
<http://creativecommons.org/licenses/by-nc-nd/2.5/>

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368


Editors: W.T.A. Harrison, J. Simpson and
M. Weil

N-(3-Methoxyphenyl)-*tert*-butanesulfinamide

Mrityunjy Datta, Alan J. Buglass and Mark R. J. Elsegood

Acta Cryst. (2010). E66, o109

This open-access article is distributed under the terms of the Creative Commons Attribution Licence <http://creativecommons.org/licenses/by/2.0/uk/legalcode>, which permits unrestricted use, distribution, and reproduction in any medium, provided the original authors and source are cited.



Acta Crystallographica Section E: Structure Reports Online is the IUCr's highly popular open-access structural journal. It provides a simple and easily accessible publication mechanism for the growing number of inorganic, metal-organic and organic crystal structure determinations. The electronic submission, validation, refereeing and publication facilities of the journal ensure very rapid and high-quality publication, whilst key indicators and validation reports provide measures of structural reliability. In 2007, the journal published over 5000 structures. The average publication time is less than one month.

Crystallography Journals **Online** is available from journals.iucr.org

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

***N*-(3-Methoxyphenyl)-*tert*-butanesulfonamide**Mrityunjoy Datta,^a Alan J. Buglass^{a*} and Mark R. J. Elsegood^b^aDepartment of Chemistry, KAIST, Daejeon 305-701, Republic of Korea, and^bChemistry Department, Loughborough University, Loughborough LE11 3TU, England

Correspondence e-mail: ajbuglass@kaist.ac.kr


Received 27 November 2009; accepted 7 December 2009

Key indicators: single-crystal X-ray study; $T = 150$ K; mean $\sigma(C-C) = 0.003$ Å; disorder in main residue; R factor = 0.042; wR factor = 0.117; data-to-parameter ratio = 15.9.

In the title compound, $C_{11}H_{17}NO_2S$, the molecules interact in a head-to-tail fashion through pairs of $N-H \cdots O$ hydrogen bonds, giving discrete centrosymmetric dimers. The $N(H)S(O)Bu$ fragment is disordered over two sets of positions, with the major component comprising 90.0 (2)%.

Related literature

For *N*-arylalkanesulfonamides, see: Datta *et al.* (2008, 2009). For *N*-alkylalkanesulfonamides, see: Sato *et al.* (1975); Ferreira *et al.* (2005); Schuckmann *et al.* (1978). For the synthesis, see: Stretter *et al.* (1969).



Experimental

Crystal data

 $C_{11}H_{17}NO_2S$ $M_r = 227.32$ Monoclinic, $P2_1/n$ $a = 12.4068$ (13) Å $b = 7.3076$ (8) Å $c = 12.9230$ (13) Å $\beta = 93.2992$ (15)° $V = 1169.7$ (2) Å³ $Z = 4$ Mo $K\alpha$ radiation $\mu = 0.26$ mm⁻¹ $T = 150$ K $0.37 \times 0.22 \times 0.20$ mm

Data collection

Bruker APEXII CCD

diffractometer

Absorption correction: multi-scan

(SADABS; Sheldrick, 2007)

 $T_{min} = 0.911$, $T_{max} = 0.950$

10627 measured reflections

2633 independent reflections

2237 reflections with $I > 2\sigma(I)$ $R_{int} = 0.032$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.042$ $wR(F^2) = 0.117$ $S = 1.07$

2633 reflections

166 parameters

149 restraints

H-atom parameters constrained

 $\Delta\rho_{max} = 0.62$ e Å⁻³ $\Delta\rho_{min} = -0.40$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H \cdots A$	$D-H$	$H \cdots A$	$D \cdots A$	$D-H \cdots A$
$N1-H1 \cdots O1^i$	0.88	2.24	2.884 (2)	130
$N1X-H1X \cdots O1X^{ii}$	0.88	2.21	2.94 (2)	141

Symmetry codes: (i) $-x + 1, -y, -z + 1$; (ii) $-x + 1, -y + 1, -z + 1$.

Data collection: APEX2 (Bruker, 2006); cell refinement: SAINT (Bruker, 2006); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: SHELXTL (Sheldrick, 2008); software used to prepare material for publication: SHELXTL and local programs.

MD and AJB thank KAIST for financial assistance.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: NG2698).

References

- Bruker (2006). APEX2 and SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.
- Datta, M., Buglass, A. J. & Elsegood, M. R. J. (2009). *Acta Cryst.* **E65**, o2823.
- Datta, M., Buglass, A. J., Hong, C. S. & Lim, J. H. (2008). *Acta Cryst.* **E64**, o1393.
- Ferreira, F., Audoin, M. & Chemla, F. (2005). *Chem. Eur. J.* **11**, 5269–5278.
- Sato, S., Yoshioka, T. & Tamura, C. (1975). *Acta Cryst.* **B31**, 1385–1392.
- Schuckmann, W., Fuess, H., Möisinger, O. & Ried, W. (1978). *Acta Cryst.* **B34**, 1516–1520.
- Sheldrick, G. M. (2007). SADABS. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Stretter, H., Krause, M. & Last, W.-D. (1969). *Chem. Ber.* **102**, 3357–3363.

supplementary materials

Acta Cryst. (2010). E66, o109 [doi:10.1107/S1600536809052507]

N-(3-Methoxyphenyl)-*tert*-butanesulfinamide

M. Datta, A. J. Buglass and M. R. J. Elsegood

Comment

The molecular structure of the title compound (I) exhibits disorder of the N(H)S(O)^tBu fragment over two sets of positions. The major component comprises 90.0 (2)% (Fig. 1), in which the N—C_{aryl} bond length is 1.418 (2) Å, similar to that [1.4225 (14) Å] in *N*-(4-methoxyphenyl)-*tert*-butanesulfinamide (Datta *et al.*, 2009). The corresponding bond length [1.457 (16) Å] in the minor component is longer, though far less precisely determined. This perhaps suggests weaker delocalization of electrons over N and the aromatic ring, which would correlate with the greater non-coplanarity of the aromatic ring and sulfinyl moiety in this component. In either case, however, the N—C_{aryl} bond is shorter than the N—C_{alkyl} bonds [1.470–1.530 Å] observed in structures of *N*-alkylalkanesulfinamides (Sato *et al.*, 1975; Schuckmann *et al.*, 1978; Ferreira *et al.*, 2005). The crystal packing of (I) shows head-to-tail interaction through NH⋯OS hydrogen bonds, forming discrete centrosymmetric dimers, as illustrated for the major component in Fig. 2. The hydrogen bonding data for both components are listed in Table 1. There is no evidence of hydrogen bonding involving the methoxy group, nor of weak CH⋯OS hydrogen bonding, as observed in the packing of *N*-phenyladamantane-1-sulfinamide (Datta *et al.*, 2008).

Experimental

Compound (I) was prepared by the method of Stretter *et al.* (1969), using *tert*-butanesulfinyl chloride (281 mg, 2 mmol) and 3-methoxyaniline (492 mg, 4 mmol) in dry ether (20 ml). After 5 h (with TLC monitoring) the colourless solid amine salt was filtered off and the solvent was removed under reduced pressure. Column chromatography (silica gel, 1% methanol-dichloromethane) provided (I) as colourless crystals (430 mg, 95%), m.p. 367 K. Single crystals suitable for X-ray analysis were obtained by evaporation of a solution of (I) in dichloromethane:hexane (1:1) at room temperature. Spectroscopic analysis: FTIR (KBr) (cm⁻¹) 3024, 1603, 1496, 1473, 1368, 1278, 1227, 1214, 1156, 1069, 953, 834. ¹H NMR (400 MHz, CDCl₃ p.p.m. with respect to TMS) δ 7.13 (dd, J = 8.0, 8.5 Hz, 1H), 6.58–6.53 (m, 3H), 5.41 (bs, 1H), 3.75 (d, J = 0.6 Hz, 3H), 1.30 (s, 9H). ¹³C (100 MHz, CDCl₃ p.p.m. with respect to TMS) δ 160.6, 143.3, 130.2, 110.6, 108.4, 104.2, 56.4, 55.2, 22.4. EIMS *m/z* (%) 228 (MH⁺, 85), 227 (M⁺, 25), 213 (17), 171 (MH⁺ - *t*Bu, 100), 123 (MH⁺ - ^tBuSO, 6), 108 (MH⁺ - ^tBuSONH, 12), 95 (53). To our knowledge, these are the first reported analytical data for (I).

Refinement

H atoms were located in a difference Fourier map and refined geometrically using a riding model. Methyl groups were refined with rotational freedom. Lengths and displacement parameters were constrained as follows: C—H = 0.95–0.98 Å and *U*_{iso}(H) = 1.2 (1.5 for CH₃) times *U*_{eq}(C, N). The minor disorder component was refined isotropically. The disorder was modelled with the aid of geometrical and displacement parameter restraints.

Figures




Fig. 1. Molecular structure of the major component of (I) with atom labels and 50% probability displacement ellipsoids for non-H atoms.




Fig. 2. Centrosymmetric dimer of (I) in the crystal packing of the major component, showing intermolecular hydrogen bonding. Symmetry code $i = -x + 1, -y, -z + 1$.

N-(3-Methoxyphenyl)-*tert*-butanesulfinamide

Crystal data

$C_{11}H_{17}NO_2S$

$M_r = 227.32$

Monoclinic, $P2_1/n$

Hall symbol: $-P\ 2_1/n$

$a = 12.4068$ (13) Å

$b = 7.3076$ (8) Å

$c = 12.9230$ (13) Å

$\beta = 93.2992$ (15)°

$V = 1169.7$ (2) Å³

$Z = 4$

$F(000) = 488$

$D_x = 1.291$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 4390 reflections

$\theta = 2.2$ – 27.2 °

$\mu = 0.26$ mm⁻¹

$T = 150$ K

Block, colourless

$0.37 \times 0.22 \times 0.20$ mm

Data collection

Bruker APEXII CCD
diffractometer

Radiation source: fine-focus sealed tube
graphite

ω rotation with narrow frames scans

Absorption correction: multi-scan
(*SADABS*; Sheldrick, 2007)

$T_{\min} = 0.911$, $T_{\max} = 0.950$

10627 measured reflections

2633 independent reflections

2237 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.032$

$\theta_{\max} = 27.3$ °, $\theta_{\min} = 2.2$ °

$h = -15 \rightarrow 16$

$k = -9 \rightarrow 9$

$l = -16 \rightarrow 16$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.042$

Primary atom site location: structure-invariant direct methods

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

$$wR(F^2) = 0.117$$

$$S = 1.07$$

2633 reflections

166 parameters

149 restraints

H-atom parameters constrained

$$w = 1/[\sigma^2(F_o^2) + (0.056P)^2 + 0.7085P]$$

$$\text{where } P = (F_o^2 + 2F_c^2)/3$$

$$(\Delta/\sigma)_{\max} = 0.001$$

$$\Delta\rho_{\max} = 0.62 \text{ e } \text{\AA}^{-3}$$

$$\Delta\rho_{\min} = -0.40 \text{ e } \text{\AA}^{-3}$$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
O1	0.54940 (10)	-0.05258 (18)	0.36978 (10)	0.0229 (3)	0.900 (2)
S1	0.50816 (3)	0.13248 (6)	0.33748 (3)	0.01938 (15)	0.900 (2)
N1	0.42312 (12)	0.2035 (3)	0.42415 (12)	0.0236 (4)	0.900 (2)
H1	0.4482	0.2281	0.4877	0.028*	0.900 (2)
C1	0.31128 (13)	0.2253 (2)	0.39789 (13)	0.0253 (4)	
C2	0.24035 (14)	0.2103 (3)	0.47687 (14)	0.0277 (4)	
H2	0.2669	0.1811	0.5453	0.033*	
C3	0.13034 (14)	0.2383 (3)	0.45550 (14)	0.0289 (4)	
O2	0.06872 (11)	0.2258 (3)	0.54029 (11)	0.0462 (4)	
C11	-0.04594 (16)	0.2441 (4)	0.52212 (19)	0.0487 (6)	
H11A	-0.0626	0.3643	0.4916	0.073*	
H11B	-0.0808	0.2325	0.5879	0.073*	
H11C	-0.0727	0.1480	0.4744	0.073*	
C4	0.08935 (14)	0.2716 (3)	0.35566 (15)	0.0300 (4)	
H4	0.0140	0.2867	0.3407	0.036*	
C5	0.16122 (16)	0.2821 (3)	0.27816 (15)	0.0384 (5)	
H5	0.1340	0.3040	0.2090	0.046*	
C6	0.27117 (15)	0.2620 (3)	0.29758 (14)	0.0328 (4)	
H6	0.3188	0.2731	0.2429	0.039*	
C7	0.62017 (14)	0.2928 (3)	0.36663 (14)	0.0212 (4)	0.900 (2)
C8	0.57870 (17)	0.4793 (3)	0.32914 (18)	0.0314 (5)	0.900 (2)
H8A	0.5209	0.5198	0.3724	0.047*	0.900 (2)
H8B	0.5507	0.4698	0.2569	0.047*	0.900 (2)
H8C	0.6379	0.5682	0.3341	0.047*	0.900 (2)
C9	0.65533 (15)	0.2918 (3)	0.48138 (15)	0.0262 (4)	0.900 (2)

supplementary materials

H9A	0.6810	0.1694	0.5015	0.039*	0.900 (2)
H9B	0.5939	0.3248	0.5220	0.039*	0.900 (2)
H9C	0.7137	0.3807	0.4945	0.039*	0.900 (2)
C10	0.7108 (2)	0.2247 (6)	0.3005 (2)	0.0282 (7)	0.900 (2)
H10A	0.7359	0.1048	0.3258	0.042*	0.900 (2)
H10B	0.7710	0.3118	0.3054	0.042*	0.900 (2)
H10C	0.6835	0.2141	0.2281	0.042*	0.900 (2)
O1X	0.5509 (11)	0.5520 (16)	0.3695 (10)	0.032 (3)*	0.100 (2)
S1X	0.5084 (4)	0.3681 (6)	0.3374 (3)	0.0303 (15)*	0.100 (2)
N1X	0.4214 (12)	0.288 (2)	0.4216 (13)	0.026 (4)*	0.100 (2)
H1X	0.4450	0.2833	0.4871	0.032*	0.100 (2)
C7X	0.6173 (12)	0.204 (2)	0.3663 (12)	0.029 (4)*	0.100 (2)
C8X	0.5762 (16)	0.016 (2)	0.3308 (16)	0.034 (4)*	0.100 (2)
H8D	0.5135	-0.0179	0.3694	0.052*	0.100 (2)
H8E	0.6335	-0.0751	0.3436	0.052*	0.100 (2)
H8F	0.5553	0.0197	0.2565	0.052*	0.100 (2)
C9X	0.6514 (16)	0.199 (3)	0.4840 (12)	0.032 (5)*	0.100 (2)
H9D	0.6766	0.3204	0.5065	0.048*	0.100 (2)
H9E	0.7096	0.1097	0.4964	0.048*	0.100 (2)
H9F	0.5893	0.1635	0.5231	0.048*	0.100 (2)
C10X	0.7138 (19)	0.269 (4)	0.308 (2)	0.025 (8)*	0.100 (2)
H10D	0.7374	0.3893	0.3344	0.038*	0.100 (2)
H10E	0.6927	0.2793	0.2338	0.038*	0.100 (2)
H10F	0.7731	0.1813	0.3179	0.038*	0.100 (2)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0234 (6)	0.0218 (7)	0.0233 (6)	0.0001 (5)	0.0005 (5)	0.0005 (5)
S1	0.0152 (2)	0.0251 (2)	0.0179 (2)	0.00068 (16)	0.00179 (14)	0.00043 (17)
N1	0.0151 (8)	0.0366 (11)	0.0193 (8)	0.0024 (7)	0.0019 (6)	0.0015 (7)
C1	0.0168 (8)	0.0337 (10)	0.0254 (9)	0.0007 (6)	0.0014 (6)	0.0012 (7)
C2	0.0215 (9)	0.0385 (10)	0.0230 (8)	-0.0003 (7)	0.0009 (6)	0.0003 (7)
C3	0.0189 (8)	0.0400 (10)	0.0282 (9)	0.0001 (7)	0.0047 (7)	-0.0026 (7)
O2	0.0164 (7)	0.0900 (13)	0.0327 (8)	0.0034 (7)	0.0060 (5)	0.0010 (7)
C11	0.0156 (9)	0.0850 (19)	0.0464 (12)	0.0009 (10)	0.0083 (8)	-0.0025 (12)
C4	0.0180 (8)	0.0380 (10)	0.0336 (10)	0.0047 (7)	-0.0021 (7)	-0.0017 (8)
C5	0.0284 (10)	0.0603 (14)	0.0259 (9)	0.0128 (9)	-0.0022 (7)	0.0034 (9)
C6	0.0243 (9)	0.0504 (12)	0.0241 (9)	0.0076 (8)	0.0050 (7)	0.0068 (8)
C7	0.0165 (8)	0.0226 (10)	0.0251 (9)	-0.0010 (7)	0.0064 (6)	-0.0021 (8)
C8	0.0282 (10)	0.0240 (10)	0.0431 (12)	0.0031 (8)	0.0109 (9)	0.0028 (9)
C9	0.0187 (9)	0.0324 (12)	0.0277 (10)	-0.0042 (8)	0.0026 (7)	-0.0061 (8)
C10	0.0195 (12)	0.0340 (18)	0.0321 (14)	0.0012 (10)	0.0109 (8)	-0.0028 (13)

Geometric parameters (\AA , $^\circ$)

O1—S1	1.4967 (14)	C8—H8B	0.9800
S1—N1	1.6652 (16)	C8—H8C	0.9800
S1—C7	1.8401 (19)	C9—H9A	0.9800

N1—C1	1.418 (2)	C9—H9B	0.9800
N1—H1	0.8800	C9—H9C	0.9800
C1—C6	1.388 (3)	C10—H10A	0.9800
C1—C2	1.390 (2)	C10—H10B	0.9800
C1—N1X	1.457 (16)	C10—H10C	0.9800
C2—C3	1.392 (2)	O1X—S1X	1.493 (12)
C2—H2	0.9500	S1X—N1X	1.681 (13)
C3—O2	1.375 (2)	S1X—C7X	1.828 (13)
C3—C4	1.381 (3)	N1X—H1X	0.8800
O2—C11	1.435 (2)	C7X—C8X	1.528 (16)
C11—H11A	0.9800	C7X—C10X	1.528 (16)
C11—H11B	0.9800	C7X—C9X	1.555 (15)
C11—H11C	0.9800	C8X—H8D	0.9800
C4—C5	1.381 (3)	C8X—H8E	0.9800
C4—H4	0.9500	C8X—H8F	0.9800
C5—C6	1.381 (3)	C9X—H9D	0.9800
C5—H5	0.9500	C9X—H9E	0.9800
C6—H6	0.9500	C9X—H9F	0.9800
C7—C9	1.522 (3)	C10X—H10D	0.9800
C7—C8	1.526 (3)	C10X—H10E	0.9800
C7—C10	1.534 (3)	C10X—H10F	0.9800
C8—H8A	0.9800		
O1—S1—N1	108.28 (9)	H8B—C8—H8C	109.5
O1—S1—C7	106.04 (8)	C7—C9—H9A	109.5
N1—S1—C7	99.44 (9)	C7—C9—H9B	109.5
C1—N1—S1	121.54 (13)	H9A—C9—H9B	109.5
C1—N1—H1	119.2	C7—C9—H9C	109.5
S1—N1—H1	119.2	H9A—C9—H9C	109.5
C6—C1—C2	119.54 (16)	H9B—C9—H9C	109.5
C6—C1—N1	122.58 (16)	C7—C10—H10A	109.5
C2—C1—N1	117.88 (15)	C7—C10—H10B	109.5
C6—C1—N1X	114.7 (6)	H10A—C10—H10B	109.5
C2—C1—N1X	119.7 (6)	C7—C10—H10C	109.5
C1—C2—C3	119.86 (16)	H10A—C10—H10C	109.5
C1—C2—H2	120.1	H10B—C10—H10C	109.5
C3—C2—H2	120.1	O1X—S1X—N1X	111.3 (8)
O2—C3—C4	124.38 (16)	O1X—S1X—C7X	106.8 (8)
O2—C3—C2	114.58 (16)	N1X—S1X—C7X	97.7 (8)
C4—C3—C2	121.03 (17)	C1—N1X—S1X	127.0 (11)
C3—O2—C11	117.06 (16)	C1—N1X—H1X	116.5
O2—C11—H11A	109.5	S1X—N1X—H1X	116.5
O2—C11—H11B	109.5	C8X—C7X—C10X	112.9 (15)
H11A—C11—H11B	109.5	C8X—C7X—C9X	109.8 (14)
O2—C11—H11C	109.5	C10X—C7X—C9X	108.3 (16)
H11A—C11—H11C	109.5	C8X—C7X—S1X	107.4 (11)
H11B—C11—H11C	109.5	C10X—C7X—S1X	106.6 (14)
C5—C4—C3	117.97 (16)	C9X—C7X—S1X	111.8 (11)
C5—C4—H4	121.0	C7X—C8X—H8D	109.5
C3—C4—H4	121.0	C7X—C8X—H8E	109.5

supplementary materials

C6—C5—C4	122.33 (17)	H8D—C8X—H8E	109.5
C6—C5—H5	118.8	C7X—C8X—H8F	109.5
C4—C5—H5	118.8	H8D—C8X—H8F	109.5
C5—C6—C1	119.20 (17)	H8E—C8X—H8F	109.5
C5—C6—H6	120.4	C7X—C9X—H9D	109.5
C1—C6—H6	120.4	C7X—C9X—H9E	109.5
C9—C7—C8	112.73 (17)	H9D—C9X—H9E	109.5
C9—C7—C10	111.27 (18)	C7X—C9X—H9F	109.5
C8—C7—C10	110.95 (19)	H9D—C9X—H9F	109.5
C9—C7—S1	111.54 (13)	H9E—C9X—H9F	109.5
C8—C7—S1	105.49 (13)	C7X—C10X—H10D	109.5
C10—C7—S1	104.40 (18)	C7X—C10X—H10E	109.5
C7—C8—H8A	109.5	H10D—C10X—H10E	109.5
C7—C8—H8B	109.5	C7X—C10X—H10F	109.5
H8A—C8—H8B	109.5	H10D—C10X—H10F	109.5
C7—C8—H8C	109.5	H10E—C10X—H10F	109.5
H8A—C8—H8C	109.5		
O1—S1—N1—C1	-113.14 (17)	N1X—C1—C6—C5	-152.9 (8)
C7—S1—N1—C1	136.38 (17)	O1—S1—C7—C9	-59.63 (15)
S1—N1—C1—C6	-26.5 (3)	N1—S1—C7—C9	52.62 (15)
S1—N1—C1—C2	154.05 (16)	O1—S1—C7—C8	177.66 (12)
S1—N1—C1—N1X	-104.9 (15)	N1—S1—C7—C8	-70.09 (14)
C6—C1—C2—C3	-2.1 (3)	O1—S1—C7—C10	60.64 (17)
N1—C1—C2—C3	177.31 (18)	N1—S1—C7—C10	172.89 (16)
N1X—C1—C2—C3	149.0 (8)	C6—C1—N1X—S1X	-10.0 (16)
C1—C2—C3—O2	-177.99 (17)	C2—C1—N1X—S1X	-162.5 (9)
C1—C2—C3—C4	3.4 (3)	N1—C1—N1X—S1X	105 (2)
C4—C3—O2—C11	1.7 (3)	O1X—S1X—N1X—C1	125.7 (13)
C2—C3—O2—C11	-176.92 (19)	C7X—S1X—N1X—C1	-122.9 (14)
O2—C3—C4—C5	179.45 (19)	O1X—S1X—C7X—C8X	-178.2 (12)
C2—C3—C4—C5	-2.1 (3)	N1X—S1X—C7X—C8X	66.7 (13)
C3—C4—C5—C6	-0.5 (3)	O1X—S1X—C7X—C10X	-57.0 (15)
C4—C5—C6—C1	1.6 (3)	N1X—S1X—C7X—C10X	-172.0 (15)
C2—C1—C6—C5	-0.3 (3)	O1X—S1X—C7X—C9X	61.2 (14)
N1—C1—C6—C5	-179.7 (2)	N1X—S1X—C7X—C9X	-53.9 (14)

Hydrogen-bond geometry (Å, °)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
N1—H1 \cdots O1 ⁱ	0.88	2.24	2.884 (2)	130.
N1X—H1X \cdots O1X ⁱⁱ	0.88	2.21	2.94 (2)	141.

Symmetry codes: (i) $-x+1, -y, -z+1$; (ii) $-x+1, -y+1, -z+1$.

Fig. 1




Fig. 2

