
This item was submitted to [Loughborough's Research Repository](#) by the author.
Items in Figshare are protected by copyright, with all rights reserved, unless otherwise indicated.

Physical activity for primary dysmenorrhea: a systematic review and meta-analysis of randomized controlled trials

PLEASE CITE THE PUBLISHED VERSION

<https://doi.org/10.1016/j.ajog.2018.04.001>

PUBLISHER

© Elsevier

VERSION

AM (Accepted Manuscript)

PUBLISHER STATEMENT

This work is made available according to the conditions of the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0) licence. Full details of this licence are available at: <https://creativecommons.org/licenses/by-nc-nd/4.0/>

LICENCE

CC BY-NC-ND 4.0

REPOSITORY RECORD

Matthewman, Gemma, Alexandra Lee, Jaidev G. Kaur, and Amanda J. Daley. 2019. "Physical Activity for Primary Dysmenorrhea: A Systematic Review and Meta-analysis of Randomized Controlled Trials". figshare. <https://hdl.handle.net/2134/33273>.

Physical activity for primary dysmenorrhea: a systematic review and meta-analysis of randomized controlled trials

Authors

Dr Gemma MATTHEWMAN MSc Birmingham, West Midlands, UK; University of Birmingham

Dr Alexandra LEE MB ChB Birmingham, West Midlands, UK; University of Birmingham

Ms Jaidev G KAUR MPharm Birmingham, West Midlands, UK; University of Birmingham

Dr Amanda J DALEY PhD Birmingham, West Midlands, UK; University of Birmingham

Structured abstract word count: 333

Main text word count: 3182

Corresponding author:

Dr Gemma Matthewman

Institute of Applied Health Research

University of Birmingham

Edgbaston

Birmingham

B15 2TT

gemmamatthewman@doctors.org.uk

The authors report no conflict of interest

Condensation: A systematic review and meta-analysis of randomized controlled trials of the effectiveness of physical activity as a treatment for primary dysmenorrhea

Short version of title: Physical activity for primary dysmenorrhea

Implications and Contributions

- A. To determine whether physical activity can reduce pain in primary dysmenorrhea.
- B. Increased physical activity reduced pain intensity by almost 2cm on the VAS scale and pain duration by almost four hours in primary dysmenorrhea.
- C. This study provides improved and updated evidence that physical activity may be an effective treatment for primary dysmenorrhea.

Commented [GM1]: I've tried to make this more in keeping with the example they gave in the original email

Structured Abstract

Background: Primary dysmenorrhea is cramping abdominal pain associated with menses. It is prevalent, affects quality of life, and can cause absenteeism. Although evidence based medical treatment options exist, women may not tolerate these or may prefer to use non-medical treatments. Physical activity has been recommended by clinicians for primary dysmenorrhea since the 1930s, but there is still no high quality evidence on which to recommend exercise as treatment. use.

Objective: We sought to determine the effectiveness of physical activity for the treatment of primary dysmenorrhea

Data sources: Systematic literature searches of Medline, Embase, Cochrane, Web of Science, CINAHL, PsycINFO, SPORTDiscus, PEDro, AMED, WHO ICTRP, Clinicaltrials.gov and OpenGrey were performed, from database inception to 24th May 2017. Google searches and citation searching of previous reviews was also conducted.

Study eligibility criteria: Studies were selected using the following PICOS criteria: Participants: non-athlete females experiencing primary dysmenorrhea; Intervention: Physical activity delivered for at least two menstrual cycles; Comparator: Any comparator; Outcomes: Pain intensity or pain duration; Study type: Randomized controlled trials.

Study appraisal and synthesis methods: Study quality was assessed using the Cochrane Risk of Bias Tool. Random effects meta-analyses for pain intensity and pain duration were conducted, with pre-specified subgroup analysis by type of physical activity intervention. Strength of the evidence was assessed using GRADE.

Results: Searches identified 15 eligible randomized controlled trials; totalling 1681 participants. Data from 11 studies was included in the meta-analyses. Pooled results demonstrated effect estimates for physical activity versus comparators for pain intensity (-1.89cm on Visual Analogue Scale, 95% CI -2.96 to -1.09) and pain duration (-3.92 hours, 95% CI -4.86 to -2.97). Heterogeneity for both these results was high and only partly mitigated by subgroup analysis. Primary studies were of low or moderate methodological quality but results for pain intensity remained stable during sensitivity analysis by study quality. GRADE assessment found moderate quality evidence for pain intensity and low quality evidence for pain duration.

Conclusion: Clinicians can inform women that physical activity may be an effective treatment for primary dysmenorrhea but there is a need for high quality trials before this can be confirmed.

Key words: Exercise, Menstrual Pain, Physical Activity, Primary Dysmenorrhea

Main text

Introduction

Primary dysmenorrhea is pain occurring with menses in the absence of underlying pathology, commonly referred to as period pains or menstrual cramps by the lay press and public.¹⁻⁴ Women may consider primary dysmenorrhea to be a normal physiological state rather than a disorder.⁵ However, studies consistently find it to be the most common gynaecological condition of adolescence,⁶⁻⁸ also affecting 60 - 76% of adult menstruating women.^{9,10} Severe symptoms are reported by 13 - 33%¹¹⁻¹³ of women with primary dysmenorrhea and absenteeism by 24 - 43%.¹³⁻¹⁵ Approximately one third of women with primary dysmenorrhea have seen a health professional because of this condition.^{16,17}

Standard, evidence based treatment is with non-steroidal anti-inflammatory medications (NSAIDs)¹⁸ or oral hormonal contraceptives.¹⁹ Other hormonal contraceptives may be helpful but the evidence for these is less robust.⁸ Some women may not be able to use medications, or may prefer to avoid them. No complementary therapies have any high quality evidence of effectiveness.²⁰⁻²⁷

There are plausible mechanisms by which physical activity may reduce pain in primary dysmenorrhea. Pain during menstruation is thought to be mediated by uterine prostaglandins, which stimulate myometrial contractions.⁸ Pain sensitization,⁸ psychosocial^{28,29} and cultural factors³⁰ may also play a role. Physical activity reduces stress,^{31,32} has anti-nociceptive properties,³³⁻³⁶ reduces levels of $\text{PGF2}\alpha$ ^{37,38} (the prostaglandin subtype most closely linked with primary dysmenorrhea).⁸ Intense exercise has significant impacts on the menstrual cycle, with female athletes found to

Commented [GM2]: This was added because one of the reviewers wanted this paragraph expanding. I have mentioned moderate and intense exercise as most of the evidence is about intense exercise which is probably physiologically different

have fewer ovulatory cycles and lower oestrogen and progesterone levels.³⁹ However, the effects of moderate exercise during the menstrual cycle are less well understood.⁴⁰

Physical activity has been recommended by clinicians for primary dysmenorrhea since the 1930s,^{41,42} and this advice is reiterated on popular^{1,3} and medical^{4,43} websites, as well as in patient information provided by the American College of Obstetrics and Gynecology.⁴⁴ However, based on current evidence, the effectiveness of physical activity is uncertain,^{26,27,45} with even less known about which types of exercise might be beneficial or when these exercises should be performed. Four reviews of interventional studies of physical activity for primary dysmenorrhea have been published (two narrative reviews in 1998⁴⁵ and 2008²⁶ and two systematic reviews in 2010 and 2016).²⁷ Results from these reviews were inconclusive due to lack of primary studies.²⁷ The most recent systematic review published in 2016⁴⁶ deviates substantially from the Cochrane library guidelines and PRISMA reporting standards in a number of ways. The protocol was not registered, no inclusion criteria were reported for the types or length of intervention, no sample search strategy was provided, studies were excluded based on publication status and language, no information regarding excluded studies was reported, and no data regarding statistical heterogeneity was provided. Additionally, there appeared to be low return rates on the initial searches for potentially eligible studies and there are discrepancies in the methodological descriptions in different sections of the report. We performed scoping searches which identified a number of new trials since the searches of this previous review were performed. An updated review is therefore required.

Objective

Commented [AD3]: Isn't it four reviews if you include the 2016 review -

We sought to systematically review the evidence from randomized controlled trials (RCTs) of the use of physical activity as treatment for primary dysmenorrhea. We sought to perform subgroup analyses based on type of intervention, type of comparator, and whether participants were adolescents or adults.⁴⁷

Methods

This review was conducted in accordance with systematic review methodologies as per the Cochrane Handbook and has been reported in compliance with the PRISMA statement. It is based upon a prospectively registered protocol, available at:

www.crd.york.ac.uk/PROSPERO/ (registration number 42017062202).⁴⁷

The search strategy was developed building on search strategies from previous similar reviews.^{26-28,32,33,48,49} The following databases were used: Medline, Embase, Cochrane Database of Systematic Reviews, Cochrane Central Register of Controlled Trials, Science Citation Index, Social Sciences Citation Index, CINAHL, PsycINFO, SPORTDiscus, PEDro, AMED, Conference Proceedings Citation Index, Social Sciences Conference Proceedings Citation Index, WHO International Clinical Trials Registry Platform, Clinicaltrials.gov and OpenGrey. Google searches and citation searching of previous reviews were also conducted.

Indexing terms (where possible) and text words (title, abstract, key words and text search) were used for "physical activity" and "dysmenorrhea" terms. Language, date or publication type restrictions were not applied. "Humans" filters were used on some databases with large return rates (e.g. Medline) to enable easier handling of search results. Validated RCT filters were used where required,⁵⁰⁻⁵² the inbuilt search filter was used for CINAHL. The Medline search strategy (see Appendix A) was piloted for sensitivity and specificity using studies found during the initial scoping searches.

No changes were required following piloting. Searches performed on other databases used the same text terms as the piloted Medline search, with index terms adapted for the specific database.

Eligibility criteria

Published and unpublished studies, in any language, were included where the following PICOS criteria were met:

- Participants: Non-athlete females with regular menstruation, experiencing primary dysmenorrhea (diagnosis as defined by report), not using hormonal contraception
- Interventions: Physical activity interventions delivered over two or more menstrual cycles; as a single intervention or as a co-intervention, in any setting and via any mode of delivery
- Comparators: Any comparator that did not involve physical activity, including active comparators and usual care or no treatment
- Outcomes: Pain intensity (most painful day or average pain intensity on days that pain was experienced) measured by a validated tool, or pain duration measured in hours
- Study type: RCTs

Athletes were excluded as those exercising at very high levels have different menstrual cycle characteristics to moderate or low level exercisers.³⁹ Hormonal contraception also significantly alters menstrual cycle physiology. Those with irregular menstruation are likely to have an underlying gynecological disorder and therefore consideration to excluding these women should be made in the primary studies. An author defined diagnosis of primary dysmenorrhea was used as the diagnosis is

usually based on history and examination, with pelvic examination typically avoided in adolescents.⁵³

Title and abstract screening was performed independently by two reviewers and any discrepancies were resolved by consensus between the two reviewers. Full text screening for inclusion of eligible studies was completed by two independent reviewers; discrepancies were resolved by consensus between these reviewers. Study authors were contacted for missing information with a reminder sent after three weeks if there had been no reply. In total, 20 study authors were contacted for further information regarding 17 studies but only five replied.

Data extraction

The data extraction form was adapted from the Cochrane Good Practice Data Extraction form⁵⁴ and was piloted prior to use. Data from included studies was extracted for participants (setting, population, method of diagnosis, inclusion / exclusion criteria, sample size, age range), intervention (type of intervention, method, timing and frequency of delivery, duration), comparators (type of comparator, timing and duration), and outcomes (time point measured, measurement tool, mean, variance). Data extraction was completed by two independent reviewers using the full text copy and any supplementary information (protocols, correspondence from authors). The main publication was used as the reference and other sources were used to obtain any information that was not reported in the main study publication. Discrepancies were resolved by consensus between the two reviewers.

Assessment of risk of bias

The Cochrane Collaboration Risk of Bias Tool was used⁵⁵ with one adaption: “blinding of participants / personnel” was changed to “blinding to study purpose / group” as physical activity interventions do not allow complete blinding.^{26,56} Studies could therefore still be rated to be of high methodological quality despite being at high risk of bias. The main biases considered in the “other bias” section were recall bias, interviewer bias, contamination, the Hawthorne effect and the effect of co-interventions. Studies were assessed for quality at the study level by two independent reviewers using the Cochrane guidance.⁵⁰ Discrepancies were resolved by consensus. Quality assessment was used for descriptive purposes and sensitivity analysis only.

Data synthesis

Review Manager 5.3 (Revman) was used for statistical analyses. Meta-analyses of pain intensity and duration were performed as specified in the review protocol.⁴⁷ Where trials compared two physical activity interventions against one comparator, they were considered as two separate trials;⁵⁷⁻⁶⁰ the number of participants in the comparator group was evenly divided between the trials to avoid double-counting of comparators. The variance was adjusted accordingly where required. The final participant number (n) was not provided for three studies;⁶⁰⁻⁶² for these studies n was assumed to be the total randomized. Results for Ortiz 2015⁶³ were obtained from a graph; they did not specify the measure of variance so this was assumed to be standard deviation.

Results were combined using the weighted mean difference, as most studies reported pain intensity using a visual analogue scale (VAS) in centimetres and pain duration in hours. VAS is a 10cm, usually horizontal, line anchored by the phrases “no pain” and “worst pain imaginable” at each end. One study⁶⁴ used the McGill questionnaire,

which cannot be converted to VAS, so data from this trial could not be included in the meta-analysis. The remaining studies reported pain intensity using VAS in millimetres⁶³ and pain duration in days.^{62,65} These results were converted to centimetres and hours respectively before analysis. A correlation coefficient of 0.6 was used to estimate the standard deviation of the mean difference where this was not provided, based on the result obtained in an RCT of a physical activity intervention in a similar population.⁶⁶ Inverse variance methods were used for weighting in the meta-analyses. The random effects model was used as it was anticipated there may be a high degree of heterogeneity. I^2 was used to assess heterogeneity; an I^2 value greater than 50% was considered to indicate substantial heterogeneity.⁵⁰ Funnel plots were produced to look for publication bias.

Cluster RCTs could not be included in the meta-analyses as no intra-cluster correlation coefficient was reported in the eligible trials. Separate pooling of cluster RCTs was performed for pain duration but only one cluster randomized study reported pain intensity in a format that could be used. Subgroup analysis was not possible for comparator type as specified in the protocol due to insufficient primary studies. Subgroup analysis by age, which was also specified in the protocol, was not possible as most included studies did not provide enough detail on age ranges.

Strength of the evidence

The strength of the evidence was assessed by GRADE at the outcome level for pain intensity and pain duration using GRADE Pro / GDT. Two independent reviewers performed GRADE assessment with discrepancies resolved by consensus. A starting rating of high quality evidence was downgraded by one level for serious concerns (or

by two levels for very serious concerns) for risk of bias, inconsistency, indirectness, imprecision and publication bias.

Results

Searches were performed on 24th May 2017, resulting in 582 returns once duplicates were removed. The returns for individual databases are given in Appendix B. The PRISMA flow diagram, representing the flow of studies through the selection process, is shown in Figure 1. 69 articles were assessed at the full text stage, with 54 excluded at this stage. A list of studies excluded at this point can be found, with reasons for exclusion, in Appendix C.

Nine studies were only found in Persian or Mandarin. These papers were assessed with the assistance of native Persian and Mandarin speakers. Where the full text could not be located the study authors were contacted where possible. Two theses and one conference abstract could not be located by any method and were thus excluded at the full text stage.

Study characteristics

Fifteen RCTs, all published since 2011, met the review inclusion and exclusion criteria. This resulted in a total of 1681 participants across all included studies.^{57-65,67-76} Details of these studies are presented in Table 1. Included studies were small or medium sized single-centre trials from a range of countries but primarily Iran or India. Most studies recruited university students. Diagnosis of primary dysmenorrhea was usually based on clinical history,^{58,60-62,64,67,68,71-76} four studies performed a clinical examination for all participants,^{59,63,65,69} and three used ultrasound^{59,65,69} to exclude secondary causes. A range of physical activity interventions were used. These could

be categorised into: aerobic exercise,^{58,60,63,67,71,73,76} stretching exercises,^{59,61-64,67-69,74,75} yoga^{60,65,70,72} or Kegels exercises.^{57,58,63,76} Ortiz 2015 used a mixed intervention.⁶³ The majority of studies asked participants to perform exercises throughout the menstrual cycle, but not during menstruation.^{59,61,62,68,69,71,74} Reyhani 2013 asked participants to exercise by brisk walking for the first three days of menstruation.⁷³ Rakhshae 2011 asked participants to perform yoga in the luteal phase of the menstrual cycle.⁷²

Synthesis of results

Meta-analysis of pain intensity (Figure 3) produced a pooled effect estimate of -1.89cm (95% CI -2.96 to -1.09), representing a statistically significant reduction in pain intensity for those in the intervention (physical activity) group relative to comparators. Heterogeneity was high ($I^2 = 95\%$).

Subgroup analysis by intervention demonstrated effect sizes of -1.29cm (95% CI -2.38 to -0.21, I^2 83%) for aerobic exercise interventions; -1.67cm (95% CI -2.70 to -0.63, I^2 94%) for stretching exercise interventions; -1.81cm (95% CI -2.37 to -1.61, I^2 0%) for yoga interventions; -1.68cm (95% CI -2.43 to -0.93, I^2 0%) for Kegels exercise interventions and -4.70cm (95% CI -5.15 to -4.25) for the single mixed intervention trial. Studies that could not be included in the meta-analysis demonstrated the same direction of treatment effect.^{64,69,71,72,75}

Meta-analysis of pain duration (Figure 4) produced a pooled estimate of effect of -3.92 hours (95% CI -4.86 to -2.97), representing a reduction in pain duration for those in the intervention (physical activity) group relative to comparators.

Heterogeneity was high ($I^2 = 78\%$). Data from two cluster RCTs was combined with a similar pooled effect size of -3.34 hours (95% CI -4.15 to -2.53).

Subgroup analysis by intervention demonstrated effect sizes of -15.64 hours (95% CI -26.96 to -4.32, I^2 49%) for aerobic exercise interventions; -3.53 hours (95% CI -4.25 to -2.81, I^2 82%) for stretching exercise interventions; -6.74 hours (95% CI -13.4 to -0.03, I^2 32%) for yoga exercise interventions; and -21.00 hours (95% CI -38.70 to -3.30) for the single Kegels exercise intervention.

Four studies could not be included in the meta-analysis (WHY?) and six further studies did not report on both pain intensity and pain duration in a way that could be utilised for pooled effect estimates (see Appendix E for reasons for exclusion from the meta-analysis).

Sensitivity analysis was performed for type of comparator and timing of intervention (not specified in protocol), with no significant change in the combined estimate of treatment effect. Funnel plot asymmetry was seen for both outcomes. Pain intensity did not demonstrate the classical funnel shape, possibly due to the heterogeneity of primary studies. The funnel plot for pain duration suggested publication bias. This is potentially due to selective outcome reporting as five studies included in the pain intensity meta-analysis did not publish data on pain duration, and most studies were found to be at a high risk of selective outcome reporting. However, results remained statistically significant when the smaller studies contributing to this asymmetry were removed.

Analysis of absenteeism was planned but this was only reported in one study with no measure of variance given.⁶²

Risk of bias of included studies

Commented [AD4]: I don't think you give a reason for the 6 but not the 4?

Most included studies were at high risk of bias in multiple areas of study design, or did not report sufficiently in order for a conclusion to be made about the risk of bias (see Figure 2). The randomization process was not fully described for most studies^{58,60-62,64,68,69,71,72,74-76} and allocation concealment was only performed in two studies.^{63,70} No studies reported blinding participants to study purpose or group and only one study reported blinding outcome assessors.⁶³ Most studies did not report how or when they measured pain intensity.^{57-61,63,65,68-70} Registered protocols were found for three studies,^{58,62,64,74-76} of which two proposed outcomes that were not reported in the final study.^{74,75} Selective outcome reporting is also suggested by the range of outcomes reported across studies. Results were sometimes reported incompletely;^{63,64,68,75} for example, Aboushady 2016⁶⁸ did not report post-intervention pain intensity in the control group. Most studies reported no loss to follow up.^{57,58,62,64,65,68-70,73-76} Those studies that did report loss to follow up did not use intention to treat analysis.^{59,63,72} The remaining studies did not report how many participants completed the intervention and follow up.^{60,61,71}

Most biases would be expected to affect the results such that they increased the magnitude of the treatment effect. However, when low quality studies were removed (Score < 3 on risk of bias assessment in Figure 2), there was an increase in the pooled estimate of treatment effect for pain intensity (from -1.89cm (95% CI -2.96 to -1.09) to -2.87cm (95% CI -5.10 to -0.63)). Only one study of moderate quality assessed pain duration with a non-significant estimate of treatment effect of -2.64 hours (95% CI -11.58 to 6.30) suggesting that the evidence for the effect of physical activity on pain duration is less reliable.

Comment

Main findings

This systematic review and meta-analysis suggests that physical activity may be an effective intervention for primary dysmenorrhea. However, these results should be interpreted with caution, as heterogeneity was high and only partially mitigated by sub group analysis. Studies were of low or moderate quality, mainly due to performance bias and potential selective outcome reporting. Nevertheless, results for pain intensity remained stable when low quality studies were removed providing some reassurance of the treatment effect observed. All studies demonstrated an improvement in pain (intensity and / or duration) with intervention, including those that could not be included in the meta-analysis. The overall assessment of the strength of evidence using GRADE showed moderate quality evidence for pain intensity and low quality evidence for pain duration (see Figure 5).

As well as considering the statistical significance and methodological quality of the results it is important to place these within a clinical context. No minimal clinically important difference (MCID) is available in the literature for pain intensity measured by VAS in primary dysmenorrhea, but the MCID in endometriosis is 1cm.⁷⁷ This suggests that the pooled estimate, at almost 2cm, is clinically significant. There are no reported values for the MCID for pain duration in primary dysmenorrhea or similar conditions.

Strengths and limitations

This review was conducted in accordance with systematic review methodologies as described in the Cochrane Handbook⁵⁰ and has been reported in compliance with the PRISMA statement⁷⁸. A prospective protocol was registered on PROSPERO, ensuring methods were specified *a priori*, unlike previous reviews.⁴⁷ Substantially more RCTs

were found in this review than all previous reviews.. Searches used in this review were also more comprehensive than previous reviews; covering more databases, and identifying grey literature, such as theses and conference proceedings that were not identified in previous reviews. All eligible studies that were not published in English were translated so that they could be considered for inclusion. In compliance with current best practice guidelines for systematic reviews, eligibility screening, data extraction, quality assessment and strength of evidence assessment were all performed by two independent reviewers. The meta-analyses for this review contain the largest number of RCTs to date, and assess both pain intensity and pain duration (only the former has been previously assessed by meta-analysis). Our review is also the first to include subgroup analysis by type of physical activity. Interrogation of the data using sensitivity analysis and Funnel plots was performed, which was not the case in previous reviews. This review is also the first to report on strength of the evidence using GRADE. This review is therefore the most complete, up to date and methodologically rigorous review of the effectiveness of physical activity interventions for primary dysmenorrhea.

Despite this the findings remain limited by the number of primary studies, trial sample size and the quality of included studies. No high quality trials were identified, and reporting of trial methodology was not always clear. Publication bias was suggested for pain duration. The results of this review are subject to high levels of heterogeneity, introducing some uncertainty about the effectiveness of physical activity. Heterogeneity appeared to occur because studies evaluated a wide range of physical activity interventions. Attempts to resolve this by conducting subgroup analysis were somewhat limited because of insufficient primary studies. Insufficient data from primary studies also prevented reporting of one of the pre-specified outcomes

Commented [AD5]: Who by? This is incomplete?

(absenteeism) and two of the pre-specified subgroup analyses (adolescents and adults, comparator type).

Comparison with existing literature

Increased physical activity was identified as a small protective factor against experiencing dysmenorrhea in a 2006 systematic review of observational studies (odds ratio (OR) of 0.89, 95% CI 0.80 to 0.99).²⁸ A non-systematic review of controlled trials published in 1998 also found a beneficial effect, but noted there was a paucity of methodologically robust studies to confirm this.⁴⁵ Interestingly, the review authors considered three trials to be randomized despite not being reported as such, and not being considered as such in other reviews.^{26,27} A non-systematic review in 2009²⁶ and a Cochrane library systematic review in 2010 (including both primary and secondary dysmenorrhea)²⁷ identified just one small RCT which demonstrated a beneficial effect of treadmill running.⁷⁹ This single trial had some methodological limitations and therefore the previous reviews concluded that there was insufficient evidence to recommend the intervention. The most recent systematic review found a beneficial effect of physical activity but similarly reported that included trials contained methodological flaws limiting the strength of their conclusions.⁴⁶

Conclusions and Implications

This review provides moderate quality evidence that physical activity may reduce pain intensity and low quality evidence that it may reduce pain duration in primary dysmenorrhea. Whilst physical activity is currently recommended in clinical guidelines for primary dysmenorrhea, more high quality studies are needed before this can be confirmed. Future trials should adhere to international reporting guidelines, and seek

to minimise sources of bias. Trials that evaluate the optimum type and timing of physical activity interventions are also required.

Acknowledgements:

Sue Bayliss (University of Birmingham) for assistance with building the search strategy, Sayeed Haque (University of Birmingham) for assistance with statistical analysis, Natalie-Tyldesley-Marshall (University of Birmingham) for assistance with reviewing articles, Manadana Zanganeh (University of Birmingham) for assistance with translating and reviewing articles published in Persian, Zhaonan Wang (University of Birmingham) for assistance with translating and reviewing articles published in Mandarin.

References

1. Uzoma K. Can exercising stop menstrual cramps? 2015; <http://www.livestrong.com/article/384938-can-exercising-stop-menstrual-cramps/>. Accessed 10th April, 2017.
2. Wilkinson E. Exercise 'no aid' for period pain. 2009; <http://www.nhs.uk/news/2009/12December/Pages/Exercise-no-relief-for-period-pain.aspx>. Accessed 10th April, 2017.
3. 5-Minute Workout: Relief from Cramps. 2017; <http://www.fitnessmagazine.com/workout/express/5-minute/moves-to-relieve-cramps/>. Accessed 10th April, 2017.
4. NHS Choices. Period Pain. 2016; <http://www.nhs.uk/conditions/periods-painful/pages/introduction.aspx#Treating>. Accessed 10th April, 2017.
5. Wong LP. Attitudes towards dysmenorrhoea, impact and treatment seeking among adolescent girls: a rural school-based survey. *Australian Journal of Rural Health*. 2010;19:218 - 223.
6. Berkley KJ. Primary dysmenorrhoea: An urgent mandate. *Pain: Clinical Updates*. 2013;21(3).
7. Proctor M, Farquhar C. Diagnosis and management of dysmenorrhoea. *BMJ*. 2006;332(7550):1134 - 1138.
8. Iacovides S, Avidon I, Baker FC. What we know about primary dysmenorrhea today: a critical review. *Human Reproduction Update*. 2015;21(6):762 - 778.
9. Weissman AM, Hartz AJ, Hansen MD, Johnson SR. The natural history of primary dysmenorrhoea: a longitudinal study. *British Journal of Obstetrics and Gynaecology*. 2004;111(4):342 - 352.
10. Burnett MA, Antao V, Black A, et al. Prevalence of primary dysmenorrhea in Canada. *Journal of Obstetrics and Gynaecology Canada*. 2005;27(8):765 - 770.
11. Suvitie PA, Hallamaa MK, Matomaki JM, Makinen JI, Perheentupa AH. Prevalence of Pain Symptoms Suggestive of Endometriosis Among Finnish Adolescent Girls (TEENMAPS Study). *Journal of Pediatric and Adolescent Gynecology*. 2016;29(2):97 - 103.
12. Nooh AM, Abdul-Hady A, El-Attar N. Nature and prevalence of menstrual disorders among teenage female students at Zagazig University, Zagazig, Egypt. *Journal of Pediatric and Adolescent Gynecology*. 2016;29(2):137 - 142.
13. Esen I, Oğuz B, Serin HM. Menstrual characteristics of pubertal girls: A questionnaire-based study in Turkey. *Journal of Clinical Research in Pediatric Endocrinology*. 2016;8(2):192 - 196.
14. Banikarim C, Chacko MR, Kelder SH. Prevalence and impact of dysmenorrhea on Hispanic female adolescents. *Archives of Pediatric and Adolescent Medicine*. 2000;154:1226.
15. Agarwal A, Venkat A. Questionnaire study on menstrual disorders in adolescent girls in Singapore. *Journal of Pediatric and Adolescent Gynecology*. 2009;22:365.
16. Hewison A, Van den Akker OB. Dysmenorrhoea, menstrual attitude and GP consultation. *British Journal of Nursing*. 1996;5(8):480 - 484.
17. Hillena TIJ, Grbavac SL, Johnston PJ, Straton JAY, Keogh JMF. Primary dysmenorrhea in young Western Australian women: prevalence, impact, and knowledge of treatment. *Journal of Adolescent Health*. 1999;25(1):40 - 45.
18. Majoribanks J, Ayeleke RO, Farquar C, Proctor M. Nonsteroidal anti-inflammatory drugs for dysmenorrhoea. *Cochrane Database of Systematic Reviews*. 2015:CD001751.
19. Wong CL, Farquhar C, Roberts H, Proctor M. Oral contraceptive pill for primary dysmenorrhoea. *Cochrane Database of Systematic Reviews*. 2009(4):CD002120.
20. Smith CA, Armour M, Zhu X, Li X, Lu ZY, Song J. Acupuncture for dysmenorrhoea. *Cochrane Database of Systematic Reviews*. 2016:CD007854.
21. Pattanittum P, Kunyanone N, Brown J, et al. Dietary supplements for dysmenorrhoea. *Cochrane Database of Systematic Reviews*. 2016:CD002124.

22. Proctor ML, Murphy PA. Herbal and dietary therapies for primary and secondary dysmenorrhoea. *Cochrane Database of Systematic Reviews*. 2001;3:CD002124.
23. Proctor ML, Hing W, Johnson TC, Murphy PA. Spinal manipulation for primary and secondary dysmenorrhoea. *Cochrane Database of Systematic Reviews*. 2001;4:CD002119.
24. Zhu X, Proctor M, Bensoussan A, Wu E, Smith CA. Chinese herbal medicine for primary dysmenorrhoea. *Cochrane Database of Systematic Reviews*. 2008;2:CD005288.
25. Proctor ML, Smith CA, Farquhar CM, Stones RW. Transcutaneous electrical nerve stimulation and acupuncture for primary dysmenorrhoea. *Cochrane Database of Systematic Reviews*. 2002;1:CD002123.
26. Daley AJ. Exercise and primary dysmenorrhoea : a comprehensive and critical review of the literature. *Sports Medicine*. 2008;38(8):659 - 670.
27. Brown J, Brown S. Exercise for dysmenorrhoea. *Cochrane Database of Systematic Reviews*. 2010(2):CD004142.
28. Latthe P, Mignini L, Gray R, Hills R, Khan K. Factors predisposing women to chronic pelvic pain: systematic review. *BMJ*. 2006;332:749 - 755.
29. Ju H, Jones M, Mishra G. The prevalence and risk factors of dysmenorrhea. *Epidemiologic Review*. 2014;36:104 - 113.
30. Faramarzi M, Salmalia H. Association of Psychologic and Nonpsychologic Factors With Primary Dysmenorrhea. *Iranian Red Crescent Medical Journal*. 2014; 16(8).
31. Josefsson T, Lindwall M, Archer T. Physical exercise intervention in depressive disorders: meta-analysis and systematic review. *Scandinavian Journal of Medicine and Science in Sports*. 2014;24(2):259 - 272.
32. Byrne A, Byrne DG. The effect of exercise on depression, anxiety and other mood states: a review. *Journal of Psychosomatic Research*. 1993;37(6):565 - 574.
33. Geneen LJ, Moore RA, Clarke C, Martin D, Colvin LA, Smith BH. Physical activity and exercise for chronic pain in adults: an overview of Cochrane Reviews. *Cochrane Database of Systematic Reviews*. 2017:CD011279.
34. Stagg NJ, Mata HP, Ibrahim MM, et al. Regular exercise reverses sensory hypersensitivity in a rat neuropathic pain model: role of endogenous opioids. *Anaesthesiology*. 2011;114(4):940 - 948.
35. Bortz WM, Angwin P, Mefford IN. Catecholamines, dopamine and endorphin levels during extreme exercise. *New England Journal of Medicine*. 1981;305:466 - 467.
36. Colt EW, Wardlaw SL, Frantz AG. The effect of running on plasma beta endorphin. *Life Sciences*. 1981;28:1637 - 1640.
37. Anderson SD, Pojer R, Smith ID, Temple D. Exercise-related changes in plasma levels of 15-keto-13,14-dihydro-prostaglandin F2alpha and noradrenaline in asthmatic and normal subjects. *Scand J Respir Dis*. 1976;57(1):41 - 48.
38. Rong Z, Shan F. The efficacy and mechanism of yoga treatment for primary dysmenorrhea suffered by female college students. *Journal of Physical Education Guangzhou*. 2013;20(6):115 - 119.
39. Maïmoun L, Georgopoulos NA, Sultan C. Endocrine Disorders in Adolescent and Young Female Athletes: Impact on Growth, Menstrual Cycles, and Bone Mass Acquisition. *Journal of Clinical Endocrinology and Metabolism*. 2014;99(11):4037 - 4050.
40. Stoddard JL, Dent CW, Shames L, Bernstein L. Exercise training effects on premenstrual distress and ovarian steroid hormones. *European Journal of Applied Physiology*. 2006;99:27 - 37.
41. Clow AES. Treatment of dysmenorrhoea by exercise. *BMJ*. 1932;1:4 - 5.
42. Any Questions? Exercises for dysmenorrhoea. *BMJ*. 1960;1:71.
43. Smith RP, Kaunitz AM. Patient education: Painful menstrual periods (dysmenorrhea). *Beyond the Basics* 2017; <https://www.uptodate.com/contents/painful-menstrual-periods-dysmenorrhea-beyond-the-basics>. Accessed 10th April, 2017.

44. American College of Obstetrics and Gynaecology. Dysmenorrhea: Painful Periods. 2015; <https://www.acog.org/Patients/FAQs/Dysmenorrhea-Painful-Periods>. Accessed 22nd November, 2017.
45. Golomb LM, Solidum AA, Warren MP. Primary dysmenorrhea and physical activity. *Medicine and Science in Sports and Exercise*. 1998;30(6):906 - 909.
46. Abaraogu UO, Tabansi-Ochiogu CS, Igwe ES. Effectiveness of exercise therapy on pain and quality of life of patients with primary dysmenorrhea: a systematic review with meta-analysis. *Turkish Journal of Physical Medicine and Rehabilitation*. 2016;62(4):346 - 354.
47. Matthewman G, Daley AJ, Bates E. A systematic review of randomised controlled trials of the effectiveness of increased physical activity as a treatment option in primary dysmenorrhoea. *PROSPERO*. 2017:CRD42017062202.
48. Mishra SI, Scherer RW, Snyder C, Geigle PM, Berlanstein DR, Topaloglu O. Exercise interventions on health-related quality of life for people with cancer during active treatment. *Cochrane Database of Systematic Reviews*. 2012:CD008465.
49. Proctor D, Murphy PA, Pattison HM, Suckling JA, Farquhar C. Behavioural interventions for dysmenorrhoea. *Cochrane Database of Systematic Reviews*. 2007:CD002248.
50. Higgins JPT, Green S. *Cochrane Handbook for Systematic Reviews of Interventions*. In: Higgins JPT, Green S, eds.: The Cochrane Collaboration; 2011.
51. Wong SS-L, Wilczynski NL, Haynes RB. Developing optimal search strategies for detecting clinically sound treatment studies in EMBASE. *Journal of the Medical Library Association*. 2006;94(1):41 - 47.
52. Eady AM, Wilczynski NL, Haynes RB. PsycINFO search strategies identified methodologically sound therapy studies and review articles for use by clinicians and researchers. *Journal of Clinical Epidemiology*. 2008;61(1):34 - 40.
53. Dysmenorrhoea. *NICE Clinical Knowledge Summaries* 2014; <https://cks.nice.org.uk/dysmenorrhoea#!backgroundsub>. Accessed 7th June, 2017.
54. Effective Practice and Organisation of Care (EPOC) Group. Data extraction form. *EPOC Resources for Review Authors*. 2013.
55. Hartling L, Ospina M, Liang Y, et al. Risk of bias versus quality assessment of randomised controlled trials: cross sectional study. *BMJ*. 2009;339:1017.
56. Theou O, Stathokostas L, Roland KP, et al. The effectiveness of exercise interventions for the management of frailty: A systematic review. *Journal of Aging Research*. 2011;569194.
57. Kaur A, Gaurav S, Dhakshinamoorthy P. Comparison of effect of fast and slow kegels exercises in reducing pain in primary dysmenorrhoea: experimental design. *Physiotherapy and Occupational Therapy Journal*. 2013;6(135 - 41).
58. Nasri M, Barati AH, Ramezani AR. Comparison of aerobic training and pelvic floor muscle exercise on primary dysmenorrhoea in adolescent girls. *Journal of Clinical Nursing and Midwifery*. 2016;5(3):53 - 61.
59. Saleh HS, Mowafy HE, Abd El Hameid AA. Stretching or core strengthening exercises for managing primary dysmenorrhea. *Journal of Womens Health Care*. 2016;5(1):1 - 6.
60. Siahpour T, Nikbakht M, Rahimi E, Rabiee MA. The effect of 8 weeks aerobic exercise and yoga on primary dismenorrhea. *Armaghane-Danesh, Yasuj University of Medical Sciences Journal* 2013:483.
61. Patel NS, Tanna T, Bhatt S. Effect of active stretching exercises on primary dysmenorrhea in college going female students. *Indian Journal of Physiotherapy and Occupational Therapy*. 2015;9(3):72 - 76.
62. Motahari-Tabari N, Shirvani MA, Alipour A. Comparison of the effect of stretching exercises and mefenamic acid on the reduction of pain and menstruation characteristics in primary dysmenorrhoea: A randomized clinical trial. *Oman medical journal*. 2017;32(1):47 - 53.

63. Ortiz MI, Cortes-Marquez SK, Romero-Quezada LC, Murguía-Canovas G, Jaramillo-Diaz AP. Effect of a physiotherapy program in women with primary dysmenorrhea. *European Journal of Obstetrics Gynecology and Reproductive Biology*. 2015;194:24 - 29.
64. Behbahani BM, Ansari-pour L, Akbarzadeh M, Zare N, Hadianfard MJ. Comparison of the effects of acupressure and self-care behaviors training on the intensity of primary dysmenorrhea based on McGill pain questionnaire among Shiraz University students. *Journal of Research in Medical Sciences*. 2016;21:104.
65. Yang NY, Kim SD. Effects of a yoga program on menstrual cramps and menstrual distress in undergraduate students with primary dysmenorrhea: A single-blind, randomized controlled trial. *Journal of Alternative and Complementary Medicine*. 2016;22(9):732 - 738.
66. Ussher M, Lewis S, Aveyard P, et al. Physical activity for smoking cessation in pregnancy: Randomised controlled trial. *BMJ*. 2015:350.
67. Vaziri F, Hoseini A, Kamali F, Abdali K, Hadianfard M, Sayadi M. Comparing the effects of aerobic and stretching exercises on the intensity of primary dysmenorrhea in the students of universities of Bushehr. *Journal of Family and Reproductive Health*. 2015;9(1):23 - 28.
68. Aboushady RM-N, El-Saidy TMK. Effect of home based stretching exercises and menstrual care on primary dysmenorrhea and premenstrual symptoms among adolescent girls. *IOSR Journal of Nursing and Health Science*. 2016;5(2):10 - 17.
69. Shahr-Jerdy S, Hosseini RS, Eivazi M. Effects of stretching exercises on primary dysmenorrhea in adolescent girls. *Biomedical Human Kinetics*. 2012;4:127 - 132.
70. Yonglitthipagon P, Muansiangsai S, Wongkhumngern W, et al. Effect of yoga on the menstrual pain, physical fitness, and quality of life of young women with primary dysmenorrhea. *Journal of Bodywork and Movement Therapies*. 2017. <http://onlinelibrary.wiley.com/doi/10.1002/jbmt.1211>
71. Sutar A, Paldhikar S, Shikalgar N, Ghodey S. Effect of aerobic exercises on primary dysmenorrhoea in college students. *IOSR Journal of Nursing and Health Science*. 2016;5(5):20 - 24.
72. Rakhshae Z. Effect of three yoga poses (cobra, cat and fish poses) in women with primary dysmenorrhea: a randomized clinical trial. *Journal of Pediatric and Adolescent Gynecology*. 2011;24(4):192 - 196.
73. Reyhani T, Jafarnejad F, Behnam H, Ajam M, Baghaei M. The effect of brisk walking on primary dysmenorrhea in girl students. *Iranian Journal of Obstetrics, Gynecology and Infertility*. 2013;16(46):14 - 19.
74. IRCT201203118822N2. Comparative effectiveness of mefenamic acid capsules, ginger and stretching exercise on primary dysmenorrhea. In:2013.
75. IRCT2013071013940N1. Effect of acupressure on primary dysmenorrhea. In. Shiraz University of Medical S, trans2014.
76. IRCT2014042017362N1. The effects of two selected types of training on primary dysmenorrhea in adolescent girls. In:2014.
77. Gerlinger C, Schumacher U, Faustman T, Colligs A, Schmitz H, Seitz C. Defining a minimal clinically important difference for endometriosis-associated pelvic pain measured on a visual analog scale: analyses of two placebo-controlled, randomized trials. *Health and Quality of Life Outcomes*. 2010;8:138.
78. Moher D, Liberati A, Tetzlaff J, Altman DG, The PRISMA Group. Preferred reporting items for systematic reviews and meta-analyses: The PRISMA statement. *Annals of Internal Medicine*. 2009;151(4).
79. Israel RG, Sutton M, O'Brien KF. Effects of aerobic training on primary dysmenorrhea symptomatology in college females. *Journal of American college Health*. 1985;33:241 - 244.




Table 1 – Description of included studies

“Cycles” refers to menstrual cycles

Study	n	Participants	Intervention(s)	Comparator(s)	Outcome(s)	Results
Aboushady 2016 ⁶⁸	80	School / college students, Saudi Arabia (16–21yrs)	<ul style="list-style-type: none"> Instructional sessions (Menstrual care, stretches); exercises at home, 20-30mins, 2x/day, 3d/w for 8wks 	Menstrual care instructional session only	Pain duration Pain intensity via VAS	Statistically significant difference in pain duration Pain intensity only reported as pre/post-test
Behbahani 2016 ^{64,75}	120	Non-medical students, Iran (18–25yrs)	<ul style="list-style-type: none"> 4wks educational classes (physiology, nutrition, exercises), “isometric exercises” at home for 4wks 	Acupressure during pain Ibuprofen 400mg 3x/day	Pain intensity via McGill questionnaire	Statistically significant reduction in pain intensity in exercise / acupressure groups compared to ibuprofen
Kaur 2013a ⁵⁷ Kaur 2013b	24	Hostel at Post-Graduate Institute, India (19–25yrs)	Slow Kegels group: <ul style="list-style-type: none"> Hot pack, 90x Kegel exercises alt days, 5–10s hold; for 8wks Fast Kegels group as above, no hold	Hot pack over lower abdomen for 10mins	Pain intensity via VAS	No statistically significant difference in pain intensity between slow Kegels and control Statistically significant reduction in pain intensity in fast Kegels compared to control
Motahari-Tabari 2017 ^{62,74}	12	Medical students, Iran (Age range not reported)	<ul style="list-style-type: none"> 15mins abdominal / pelvic stretching exercises, taught initially, 3x/wk; for 2 cycles 	Mefenamic acid 250mg 3x/day	Pain intensity via VAS Pain duration	No statistically significant difference in pain intensity or pain duration
Nasri 2016a ^{58,76} Nasri 2016b	45	High school pupils, Iran (“Teenagers”)	Aerobic group: <ul style="list-style-type: none"> 45mins observed “aerobic exercise”, 3x/wk; for 8wks Kegel group: <ul style="list-style-type: none"> 15 mins Kegel exercises; 6s hold, 3x/day 	Usual care - “no exercise”, advised no salty / fatty foods, no medications	Pain intensity via VAS Pain duration	No statistically significant reduction in pain intensity / duration between exercise groups Statistically significant reduction in pain intensity / duration compared to control
Ortiz 2015 ⁶³	192	Uni students, Mexico (18–22yrs)	<ul style="list-style-type: none"> Stretches (inc Billig / Kegel), jogging, relaxation led / monitored by instructors; 50mins, 3x/wk; for 3 cycles 	Kept in courtyard; “walking, talking and standing”	Pain intensity via VAS	Statistically significant reduction in pain intensity
Patel 2015 ⁶¹	120	Students, India (17–25yrs)	<ul style="list-style-type: none"> 6 stretches; 2x/day, 3x/wk for 8wks 	Usual care	Pain intensity via VAS	Statistically significant reduction in pain intensity

Study	n	Participants	Intervention(s)	Comparator(s)	Outcome(s)	Results
Rakhshae 2011 ⁷²	1 2 0	Uni students, Iran (17–23yrs)	<ul style="list-style-type: none"> 3 yoga poses / breathing techniques taught by booklet, for 20mins/day, luteal phase (14d) of 2 cycles 	Usual care	Pain intensity via 0–3 scale Pain duration	Statistically significant reduction in pain intensity and pain duration
Reyhani 2013 ⁷³	9 0	Nursing/midwifery students, Iran (Age range not reported)	<ul style="list-style-type: none"> 30mins brisk walking (one training session), 1st 3d of menstruation; for 3 cycles 	Usual care	Pain intensity via VAS	Statistically significant reduction in pain intensity
Saleh 2016a ⁵⁹ Saleh 2016b	1 5 0	Women from outpatient clinic, Egypt (Age range not reported)	Stretching group: <ul style="list-style-type: none"> 4 stretches, 10mins, 3x/d, 3x/wk; for 8wks Core strengthening group: <ul style="list-style-type: none"> 4 core strengthening exercises, 20mins, 4x/wk 	Usual care	Pain intensity via VAS Pain duration	Statistically significant reduction in pain intensity / pain duration in both intervention groups when compared to control
Shahr-Jerdy 2012 ⁶⁹	1 7 9	High school pupils, Iran (15–17ys)	<ul style="list-style-type: none"> 6 stretches taught initially, 10mins, 2x/d, 3x/wk; for 8wk 	Usual care - exercises taught to controls after study	Pain intensity via VAS Pain duration	Statistically significant reduction in pain intensity and pain duration
Siahpour 2013a ⁶⁰ Siahpour 2013b	6 0	Uni students, Iran (20–25yrs)	Aerobic group: <ul style="list-style-type: none"> Aerobic dance for 60 mins, 3x/wk; for 8wks Yoga group: <ul style="list-style-type: none"> 60 mins yoga, 3x/wk; “trained” 	Usual care	Pain intensity via VAS Pain duration	Statistically significant reduction in pain intensity / pain duration between aerobic and yoga groups compared to control
Sutar 2016 ⁷¹	1 0 0	Medical students, India (18–22yrs)	<ul style="list-style-type: none"> Aerobic dance for 45mins, 3x/wk, for 8wks 	Usual care	Pain intensity via VAS	Statistically significant reduction in pain intensity
Yang 2016 ⁶⁵	1 4 0	Nursing students, S Korea (20–23yrs)	<ul style="list-style-type: none"> 60mins guided yoga, 1x/wk (poses, sun salutations, relaxation); for 12wks 	Usual care - told not to do yoga	Pain intensity via VAS Pain duration	Statistically significant reduction in pain intensity No statistically significant reduction in pain duration
Yonglitthipagon 2017 ⁷⁰	3 4	Uni students, Thailand (18–22yrs)	<ul style="list-style-type: none"> 30mins yoga taught by booklet, 2x/wk; for 12wks Diary / weekly phone calls to check adherence 	Usual care	Pain intensity via VAS	Statistically significant reduction in pain intensity

Figure legends

Figure 1 – PRISMA flow diagram.

*PRISMA flow diagram demonstrating flow of studies through identification process and eligibility screening *See Appendix C for further details*

Figure 2 – Risk of Bias Summary

Summary of Risk of Bias of included studies

Figure 3 – Pain intensity meta-analysis

Random effects meta-analysis of pain intensity via VAS in cm

Figure 4 – Pain duration meta-analysis

Random effects meta-analysis of pain duration in hours

Figure 5 - GRADE evidence profile

Evidence profile for pain intensity and pain duration

Appendix A – Search strategies

Medline / Medline in Process searched 05/24/17:

1 exp Dysmenorrhea/ 3600
 2 dysmenorrh*.ti,ab. 4950
 3 (menstrua* adj2 pain).ti,ab. 720
 4 (menstrua* adj2 cramp).ti,ab. 7
 5 (period* adj2 pain*).ti,ab. 1372
 6 1 or 2 or 3 or 4 or 5 7684
 7 exp Exercise/ or exp Exercise Therapy/ 184201
 8 exp Physical Exertion/ or exp Physical Fitness/ 79696
 9 exp running/ or exp swimming/ or exp walking/81381
 10 exp tai ji/ or exp yoga/ 2941
 11 exp dancing/ or exp gardening/ or exp sports/ 162811
 12 exercis*.ti,ab. 245761
 13 “physical activit*”.ti,ab. 82503
 14 sport*.ti,ab. 57350
 15 stretch*.ti,ab. 62854
 16 fitness.ti,ab. 55378
 17 jog*.ti,ab. 2026
 18 running.ti,ab. 49262
 19 swim*.ti,ab. 32718
 20 (cycl* adj2 train*).ti,ab. 2065
 21 walk*.ti,ab. 93124
 22 yoga.ti,ab. 3190
 23 “tai ji”.ti,ab. 25
 24 “tai chi”.ti,ab.1249
 25 pilates.ti,ab. 304
 26 “physical training”.ti,ab. 5245
 27 “resistance training”.ti,ab. 5278
 28 (athlete* adj2 train*).ti,ab. 4489
 29 “weight training”.ti,ab. 914
 30 isometric*.ti,ab. 30361
 31 danc*.ti,ab. 5670
 32 7 or 8 or 9 or 10 or 11 or 12 or 13 or 14 or 15 or 16 or 17 or 18 or 19
 or 20 or 21 or 22 or 23 or 24 or 25 or 26 or 27 or 28 or 29 or 30 or 31
 731463
 33 6 and 32 312
 34 randomized controlled trial.pt. 462560
 35 controlled clinical trial.pt. 94063
 36 randomized controlled trial.sh. 462560
 37 random allocation.sh.92576
 38 double blind method.sh. 147085

39 single-blind method.sh. 24526
 40 34 or 35 or 36 or 37 or 38 or 39 645852
 41 (animals not human).sh. 6109803
 42 40 not 41 571840
 43 clinical trial.pt. 521341
 44 exp clinical trial/ 803438
 45 (clin\$ adj25 trial\$.ti,ab. 368228
 46 ((singl\$ or doubl\$ or trebl\$ or tripl\$) adj25 (blind\$ or mask\$)).ti,ab. 161643
 47 placebos.sh. 34931
 48 placebo\$.ti,ab. 192528
 49 research design.sh. 96076
 50 43 or 44 or 45 or 46 or 47 or 48 or 49 491211208
 51 50 not 41 1110374
 52 51 not 42 562047
 53 comparative study.sh. 1809971
 54 exp evaluation studies/ 232296
 55 follow up studies.sh. 586124
 56 prospective studies.sh. 456986
 57 (control\$ or 31thlete3131ve\$ or volunteer\$.ti,ab. 3751440
 58 53 or 54 or 55 or 56 or 57 5717430
 59 58 not 41 4252943
 60 59 not (42 or 52) 3578680
 61 42 or 52 or 60 4712567
 62 33 and 61 136

EMBASE searched 05/24/17:

1 exp Dysmenorrhea/ 9975
 2 dysmenorrh*.ti,ab. 6530
 3 (menstrua* adj2 pain).ti,ab. 993
 4 (menstrua* adj2 cramp).ti,ab. 11
 5 (period* adj2 pain*).ti,ab. 2026
 6 1 or 2 or 3 or 4 or 5 13542
 7 exp Exercise/ or exp Physical Activity/ 539709
 8 exp Sport/ or exp Fitness/ 159023
 9 exp dynamic exercise/ or exp isotonic exercise/ or exp anaerobic exercise/ or
 exp static exercise/ or exp aerobic exercise/ or exp isokinetic exercise/ 16049
 10 exp stretching exercise/ or exp aquatic exercise/ or exp isometric exercise/
 5696
 11 exp stretching/ or exp muscle stretching/ or exp muscle training/ or exp
 resistance training/ or exp pelvic floor muscle training/ 29219
 12 exp running/ or exp swimming/ or exp walking/ 123921
 13 exp jogging/ or exp pilates/ 2023

14 exp tai ji/ or exp yoga/ 7329
 15 exp dancing/ or exp gardening/ or exp sports/ 133605
 16 exercis*.ti,ab. 316216
 17 "physical activit*".ti,ab. 108960
 18 sport*.ti,ab. 75241
 19 stretch*.ti,ab. 66632
 20 fitness.ti,ab. 61859
 21 jog*.ti,ab. 2419
 22 running.ti,ab. 59265
 23 swim*.ti,ab. 38714
 24 (cycl* adj2 train*).ti,ab. 2321
 25 walk*.ti,ab. 124213
 26 yoga.ti,ab. 4466
 27 "tai ji".ti,ab. 49
 28 "tai chi".ti,ab. 1729
 29 pilates.ti,ab. 449
 30 "physical training".ti,ab. 6938
 31 "resistance training".ti,ab. 6242
 32 (32thlete* adj2 train*).ti,ab. 5243
 33 "weight training".ti,ab. 997
 34 isometric*.ti,ab. 34963
 35 danc*.ti,ab. 7125
 36 exp health care quality/ 2478747
 37 random.tw. 1190652
 38 clinical trial.mp. 1442523
 39 7 or 8 or 9 or 10 or 11 or 12 or 13 or 14 or 15 or 16 or 17 or 18 or 19
 or 20 or 21 or 22 or 23 or 24 or 25 or 26 or 27 or 28 or 29 or 30 or 31 or 32
 or 33 or 34 or 35 1005054
 40 36 or 37 or 38 4188607
 41 6 and 39 and 40 243

PsycINFO searched 05/24/17:

1 exp Dysmenorrhea/ 190
 2 dysmenorrh*.ti,ab. 304
 3 (menstrua* adj2 pain).ti,ab. 122
 4 (menstrua* adj2 cramp).ti,ab. 1
 5 (period* adj2 pain*).ti,ab. 229
 6 1 or 2 or 3 or 4 or 5 617
 7 exp Exercise/ or exp Aerobic Exercise/ 22140
 8 exp Physical Fitness/ or exp Physical activity/ 35122
 9 exp running/ or exp swimming/ or exp walking/ 7599
 10 exp yoga/ 1407

11 exp dancing/ or exp dance therapy/ 938
 12 exp sports/ or exp athletic training/ 22874
 13 exercis*.ti,ab. 52521
 14 "physical activit*".ti,ab. 26010
 15 sport*.ti,ab. 27171
 16 stretch*.ti,ab. 4142
 17 fitness.ti,ab. 12800
 18 jog*.ti,ab. 417
 19 running.ti,ab. 12473
 20 swim*.ti,ab. 8017
 21 (cycl* adj2 train*).ti,ab. 217
 22 walk.ti,ab. 6195
 23 yoga.ti,ab. 2052
 24 "tai ji".ti,ab. 5
 25 "tai chi".ti,ab.443
 26 pilates.ti,ab. 55
 27 "physical training".ti,ab. 520
 28 "resistance training".ti,ab. 484
 29 (33thlete* adj2 train*).ti,ab. 797
 30 "weight training".ti,ab. 178
 31 isometric*.ti,ab. 1865
 32 danc*.ti,ab. 6309
 33 (control: or random*.tw. or exp treatment/ 1218890
 34 7 or 8 or 9 or 10 or 11 or 12 or 13 or 14 or 15 or 16 or 17 or 18 or 19
 or 20 or 21 or 22 or 23 or 24 or 25 or 26 or 27 or 28 or 29 or 30 or 31 or 32
 150179
 35 6 and 33 and 34 23

CINAHL searched 05/24/17:

1 MH exercise OR MH exercise therapy OR MH physical exertion OR MH
 physical fitness OR MH locomotion OR MH exercise movement techniques OR MH
 recreation 48,339S14
 2 AB dysmenorrh* OR AB "menstrual cramp*" OR AB "menstrual pain" OR
 AB "period pain" OR AB "painful periods" OR AB "painful menstrua*" 858
 3 TI dysmenorrh* OR TI "menstrual cramp*" OR TI "menstrual pain" OR TI
 "period pain" OR TI "painful periods" OR TI "painful menstrua*" 573
 4 KW dysmenorrh* OR KW "menstrual cramp*" OR KW "menstrual pain" OR
 KW "period pain" OR KW "painful periods" OR KW "painful menstrua*" 15,172
 5 2 OR 3 OR 4 1,147
 6 AB "physical activit*" OR AB 33thlete33* OR AB (stretch* or isometric)
 OR AB (fitness or sport*) OR AB "physical training" OR AB "weight training"

OR AB "resistance training" OR AB (jog* or running or swim* or walk* or danc*) OR AB (yoga or "tai ji" or "tai chi" or pilates) 131,871
 7 TI "physical activit*" OR TI 34thlete34* OR TI (stretch* or isometric) OR TI (fitness or sport*) OR TI "physical training" OR TI "weight training" OR TI "resistance training" OR TI (jog* or running or swim* or walk* or danc*) OR TI (yoga or "tai ji" or "tai chi" or pilates) 97,173
 8 KW "physical activity" OR KW exercise 35
 9 6 OR 7 OR 8 182,705
 10 1 OR 9 202,476
 11 MH dysmenorrhea 1,025
 12 5 OR 11 1,566
 13 10 AND 12 95
 14 Limiters – Clinical Queries: Therapy – High Sensitivity 46

SPORTDiscus searched 05/24/17:

1 AB dysmenorrh* OR AB "menstrual cramp*" OR AB "menstrual pain" OR AB "period pain" OR AB "painful periods" OR AB "painful menstrua*" 157
 2 TI dysmenorrh* OR TI "menstrual cramp*" OR TI "menstrual pain" OR TI "period pain" OR TI "painful periods" OR TI "painful menstrua*" 60
 3 KW dysmenorrhea OR KW dysmenorrhoea 35
 4 1 OR 2 OR 3 190
 5 AB "physical activit*" OR AB 34thlete34* OR AB (stretch* or isometric) OR AB (fitness or sport*) OR AB "physical training" OR AB "weight training" OR AB "resistance training" OR AB (jog* or running or swim* or walk* or danc*) OR AB (yoga or "tai ji" or "tai chi" or pilates) 420,655
 6 TI "physical activit*" OR TI 34thlete34* OR TI (stretch* or isometric) OR TI (fitness or sport*) OR TI "physical training" OR TI "weight training" OR TI "resistance training" OR TI (jog* or running or swim* or walk* or danc*) OR TI (yoga or "tai ji" or "tai chi" or pilates) 272,594
 7 KW "physical activity" OR KW exercise 20,595
 8 5 OR 6 OR 7 556,573
 9 4 AND 8 53

AMED searched 05/24/17:

1 AB dysmenorrh* OR AB "menstrual cramp*" OR AB "menstrual pain" OR AB "period pain" OR AB "painful periods" OR AB "painful menstrua*" 120
 2 TI dysmenorrh* OR TI "menstrual cramp*" OR TI "menstrual pain" OR TI "period pain" OR TI "painful periods" OR TI "painful menstrua*" 130
 3 KW dysmenorrhea OR KW dysmenorrhoea 116
 4 1 OR 2 OR 3 195

5 AB "physical activit*" OR AB 35thlete35* OR AB (stretch* or isometric)
 OR AB (fitness or sport*) OR AB "physical training" OR AB "weight training"
 OR AB "resistance training" OR AB (jog* or running or swim* or walk* or danc*
) OR AB (yoga or "tai ji" or "tai chi" or pilates) 24,404
 6 TI "physical activit*" OR TI 35thlete35* OR TI (stretch* or isometric) OR
 TI (fitness or sport*) OR TI "physical training" OR TI "weight training" OR TI
 "resistance training" OR TI (jog* or running or swim* or walk* or danc*) OR TI
 (yoga or "tai ji" or "tai chi" or pilates) 19,137
 7 KW "physical activity" OR KW exercise 15,600
 8 5 OR 6 OR 7 36,091
 9 4 AND 8 7

Cochrane searched 05/24/17:

1	"dysmenorrhea":ti,ab,kw (Word variations have been searched)	1188
2	MeSH descriptor: [Dysmenorrhea] explode all trees	465
3	"menstrual cramps":ti,ab,kw	27
4	"period pains":ti,ab,kw	4
5	"painful menstruation":ti,ab,kw	17
6	"menstrual pain":ti,ab,kw	174
7	1 or 2 or 3 or 4 or 5 or 6	1235
8	MeSH descriptor: [Exercise] explode all trees	18710
9	MeSH descriptor: [Exercise Therapy] explode all trees	10374
10	MeSH descriptor: [Physical Exertion] explode all trees	3389
11	MeSH descriptor: [Physical Fitness] explode all trees	2648
12	MeSH descriptor: [Running] explode all trees	1599
13	MeSH descriptor: [Swimming] explode all trees	407
14	MeSH descriptor: [Walking] explode all trees	3482
15	MeSH descriptor: [Sports] explode all trees	12907
16	MeSH descriptor: [Yoga] explode all trees	513
17	MeSH descriptor: [Tai Ji] explode all trees	343
18	MeSH descriptor: [Dancing] explode all trees	128
19	"35thlete35*":ti,ab,kw	57524
20	"physical activit*":ti,ab,kw	13568
21	sport*:ti,ab,kw	4824
22	stretch*:ti,ab,kw	2968
23	fitness:ti,ab,kw	6049
24	jog*:ti,ab,kw	318
25	running:ti,ab,kw	3623
26	swim*:ti,ab,kw	813
27	"cycl* train*":ti,ab,kw	129
28	walk*:ti,ab,kw	14632
29	yoga:ti,ab,kw	1449
30	"tai chi":ti,ab,kw	666

31 "tai ji":ti,ab,kw 355
 32 pilates:ti,ab,kw 187
 33 "physical training":ti,ab,kw 1083
 34 "resistance training":ti,ab,kw 4765
 35 "weight training":ti,ab,kw 310
 36 isometric:ti,ab,kw 3299
 37 danc*:ti,ab,kw 488
 38 8 or 9 or 10 or 11 or 12 or 13 or 14 or 15 or 16 or 17 or 18 or 19 or 20
 or 21 or 22 or 23 or 24 or 25 or 26 or 27 or 28 or 29 or 30 or 31 or 32 or 33
 or 34 or 35 or 36 or 37 81187
 39 7 and 38 40

Web of Science searched 05/24/17:

1 TS=(dysmenorrh*) 3,781
 2 TS=("menstrual cramp*") 72
 3 TS=(menstru* NEAR/1 pain*) 634
 4 TS=(period\$ NEAR/1 pain*) 811
 5 4 OR 3 OR 2 OR 1 4,861
 6 TS=("physical activit*") 120,487
 7 TS=(36thlete36*) 346,079
 8 TS=(sport*) 103,132
 9 TS=(stretch*) 145,828
 10 TS=(fitness) 104,942
 11 TS=(jog*) 3,483
 12 TS=(running) 408,116
 13 TS=(swim*) 52,369
 14 TS=(cycl* NEAR/2 train*) 3,333
 15 TS=(walk*) 171,784
 16 TS=(yoga) 3,304
 17 TS=("tai ji") 61
 18 TS=("tai chi") 1,945
 19 TS=(pilates) 341
 20 TS=("physical training") 5,087
 21 TS=("resistance training") 6,810
 22 TS=(36thlete* NEAR/2 train*) 5,548
 23 TS=("weight training") 1,330
 24 TS=(isometric*) 35,686
 25 TS=(danc*) 15,569
 26 25 OR 24 OR 23 OR 22 OR 21 OR 20 OR 19 OR 18 OR 17 OR 16 OR
 15 OR 14 OR 13 OR 12 OR 11 OR 10 OR 9 OR 8 OR 7 OR 6 1,340,672
 27 26 AND 5 231

PEDro searched 05/24/17:

1	dysmenorrh* AND exercise*	17
2	dysmenorrh* AND physical activit*	3
3	dysmenorrh* AND yoga	6
4	dysmenorrh* AND stretch*	28

Clinicaltrials.gov searched 05/24/17:

Condition: dysmenorrhoea OR “menstrual cramps” OR “menstrual pain” OR “period pain” OR “painful periods” OR “painful menstruation”

Intervention: “physical activity” OR exercise OR sport OR stretch OR fitness OR “physical training” OR “resistance training” OR “weight training” OR jogging OR running OR walking OR swimming OR yoga OR “tai chi” OR pilates (searches both tai ji and tai chi)

13

WHO ICTRP searched 05/24/17:

Condition: dysmenorrhea OR dysmenorrhoea OR menstrual cramps OR menstrual pain OR period pain OR painful periods OR painful menstruation

Intervention: physical activity OR exercise OR sport OR stretch OR fitness OR physical training OR resistance training OR weight training OR jogging OR running OR walking OR swimming OR yoga OR tai chi OR pilates

11

OpenGrey searched 05/24/17:

Dysmenorrhea	1
Dysmenorrhoea	6
Menstrual + exercise	11
Menstrual + sport	3
Menstrual + yoga	0
Menstrual + physical activity	0

Google searched 05/24/17:

Google was searched using the terms “dysmenorrhea” and “exercise”; all returns up to page 15 were reviewed at which point no new returns were being identified. This was repeated with the alternative spelling “dysmenorrhoea”.

The Menstruation Research website was also searched in greater detail (menstruationresearch.org)

Appendix B - Database search returns

Database	Interface	Dates	Returns
MEDLINE and MEDLINE In Process	Ovid	1946 – 05/24/17	136
EMBASE	Ovid	1974 – 05/24/17	243
PsycINFO	Ovid	1806 – 05/24/17	23
Cochrane Database of Systematic Reviews (CDSR)	Cochrane Library	N/A	40 total*
Cochrane Central Register of Controlled Trials (CENTRAL)	Cochrane Library	1966 – 05/24/17	
CINAHL	EBSCO	1937 – 05/24/17	46
SPORTDiscus	EBSCO	1980 – 05/24/17	53
AMED (Allied and Complimentary Medicine Database)	EBSCO	1995 – 05/24/17	7
PEDro	NeuRA	1929 – 05/14/17	41
Science Citation Index	Web of Science	1964- 05/24/17	231 total*
Conference Proceedings Citation Index	Web of Science	1990 – 05/24/17	
Social Sciences Citation Index	Web of Science	1900 – 05/24/17	
Conference Proceedings Citation Index – Social Sciences and Humanities	Web of Science	1990 – 05/24/17	
Clinicaltrials.gov	National Institute of Health	To 05/24/17	13
WHO ICTRP	WHO	To 05/24/17	11
OpenGrey	SIGLE	1993 – 05/24/17	21
Google	Google	N/A	42 new
Citation searching	N/A	N/A	1 new

SCI/CPCI/SSCI/SS-CPCI and CDSR/CENTRAL were searched simultaneously; there is therefore no individual return numbers for these databases.

Appendix C – Excluded studies with reasons for exclusion

A reference list of excluded studies can be obtained from the study authors

Study	Reason for exclusion
<i>Abbaspour 2006</i>	No inclusion / exclusion criteria reported, no reply from authors
<i>ACTRN12613001195741</i>	Registered trial and feasibility study, contacted authors who have completed trial and are in process of publishing, however unwilling to share any of data
<i>Anonymous 1945</i>	Letter
<i>Anonymous 1960</i>	Letter
<i>Anonymous 1993</i>	Review
<i>Arora 2014</i>	No inclusion / exclusion criteria reported, no reply from authors
<i>Azima 2015a</i>	Two arm trial which is also reported as 3 arm trial - see Azima 2015b below
<i>Azima 2015b</i>	No inclusion / exclusion criteria in either of papers reporting trial, protocol contains some inclusion / exclusion criteria but does not specify regarding secondary dysmenorrhoea, irregular menstruation or hormonal contraception
<i>Chaudhuri 2013</i>	Some participants were athletes, also some had irregular periods (from correspondence with author)
<i>Chien 2013</i>	Non-randomized controlled trial
<i>Dehghanzadeh 2014</i>	Before and after trial
<i>DeWitt 1981</i>	Asked women about hormonal contraception but did not specify whether these women were excluded
<i>El Refaye 2007</i>	Thesis, no online abstract, unable to obtain full text through inter-library loans, no contact details for authors
<i>Gamit 2014</i>	Physical activity intervention for four weeks only
<i>Golub 1960</i>	Non-randomized controlled trial
<i>Golub 1963</i>	Before and after trial
<i>Halder 2012</i>	Non-randomized controlled trial
<i>Haman1945</i>	Before and after trial
<i>Harris 1955</i>	Before and after trial
<i>Heidarianpour 2016</i>	Menstrual characteristics, dysmenorrhoea considered but did not exclude secondary dysmenorrhoea
<i>Huang 2007</i>	No inclusion / exclusion criteria reported, unable to find contact details for authors
<i>Hubbell 1949</i>	Non-randomized controlled trial
<i>IRCT2013071013940N1</i>	Registered trial, results published in Behbahani 2016
<i>IRCT2016103119024N2</i>	Registered trial, unable to find published paper, contacted authors and no reply; probably still ongoing as only registered in 2016
<i>ISRCTN75567759</i>	Registered trial and published protocol, mixed-methods study with no randomization
<i>Jahromi 2008</i>	Before and after trial

<i>Kang 2009</i>	Thesis, unable to obtain full text through inter-library loans, no contact details for authors
<i>Kanwal 2016</i>	Physical activity intervention for one month only
<i>Kashef 2014</i>	Physical activity intervention for four weeks only
<i>Khare 2016</i>	Physical activity intervention for three weeks only
<i>Kaur 2014</i>	Unclear whether secondary causes excluded. No contact details for authors. Mentions randomization in abstract but in text participants are "taken at random"
<i>Kumar 2013</i>	No inclusion / exclusion criteria reported, unable to find contact details for authors
<i>Locke 1999</i>	Review
<i>Locke 1999</i>	Review
<i>Lundquist 1947</i>	Non-randomized controlled trial
<i>Mahishale 2013</i>	Physical activity intervention for one menstrual cycle only
<i>Mahvash 2012</i>	Unclear whether those on hormonal contraception or with irregular periods were excluded as no inclusion / exclusion criteria were given. No contact details found
<i>Mathur 1986</i>	Non-randomized controlled trial
<i>Monika 2012</i>	Hormonal contraception specifically provided to the groups
<i>Motesharee 2013</i>	Hormonal contraception / irregular periods not excluded
<i>Nag 2013</i>	No inclusion / exclusion criteria reported, unable to find contact details for authors. Unclear whether athletes included (although have excluded those already practicing yoga), whether those on hormonal contraception or with irregular periods were excluded.
<i>Pazoki 2013</i>	Pre-menstrual symptoms; dysmenorrhoea considered but did not exclude secondary dysmenorrhoea
<i>Pazoki 2016</i>	Pre-menstrual symptoms; dysmenorrhoea considered but did not exclude secondary dysmenorrhoea
<i>Rani 2013</i>	Hormone profiles assessed as outcome
<i>Rozebahani 2015</i>	Did not exclude secondary dysmenorrhoea
<i>Rezvani 2013</i>	Unclear whether women using hormonal contraception were excluded, no reply from authors
<i>Rong 2013</i>	Unclear whether athletes or women using hormonal contraception were excluded, unable to find contact details for authors
<i>Shah 2016</i>	No inclusion / exclusion criteria, no reply from authors
<i>Steege 1993</i>	Pre-menstrual symptoms; dysmenorrhoea considered but did not exclude secondary dysmenorrhoea
<i>Thomas 2010</i>	Physical activity intervention for 3 weeks only
<i>Vaziri 2015</i>	Menstrual symptoms assessed as outcome, unable to extract data about pain
<i>Wilt 1976</i>	Not specified whether those with secondary cause excluded, unable to obtain information from authors as no contact details
<i>Yeknami 2015</i>	No inclusion / exclusion criteria, no reply from authors

Appendix D – Risk of Bias Assessment

	Risk of Bias	Justification
Random sequence generation	<i>Low / High / Unclear</i>	
Allocation concealment	<i>Low / High / Unclear</i>	
Blinding to study group / study purpose	<i>Low / High / Unclear</i>	Studies were considered at high risk of bias if they reported that no blinding was done, or blinding was not reported but the comparator group received no intervention.
Blinding of outcome assessment	<i>Low / High / Unclear</i>	
Incomplete outcome data	<i>Low / High / Unclear</i>	
Selective outcome reporting	<i>Low / High / Unclear</i>	
Other bias (observer bias, recall bias, contamination, co-interventions, Hawthorne effect)	<i>Low / High / Unclear</i>	<p>Studies were considered at high risk of bias if one of these biases was present. If none of these biases were adequately described studies were considered at unclear risk of bias</p> <p>Interviewer bias - high risk if outcomes assessed during interview</p> <p>Recall bias – high risk if outcomes assessed more than one day after the end of menstruation</p> <p>Contamination – high risk if participants were from schools / colleges / individual courses unless cluster randomized</p> <p>Co-interventions – high risk if there was a co-intervention or physical activity was performed in a group</p> <p>Hawthorne effect – high risk if physical activity was performed in a group or closely monitored setting</p>

Appendix E - Reasons For Exclusion From Meta-Analysis

	Pain intensity	Pain duration	Absenteeism
Aboushady 2016	Only reported for intervention group	Included	Not reported
Behbahani 2016	Pain intensity reported via McGill questionnaire	Not reported	Not reported
Kaur 2013	Included	Not reported	Not reported
Motahari-Tabari 2017	Included	Included	No measure of variance given
Nasri 2016	Included	Included	Not reported
Ortiz 2015	Included, derived from graph	Not reported	Not reported
Patel 2015	Included	Not reported	Not reported
Rakhshae 2011	VAS reported as 0 to 3 categorical scale	Included in cluster randomised meta-analysis	Not reported
Reyhani 2013	Included	Not reported	Not reported
Saleh 2016	Included	Included	Not reported
Shahr-Jerdy 2012	Cluster randomised; no other clusters to combine with	Included in cluster randomised meta-analysis	Not repeated
Siahpour 2013	Included	Included	Not reported
Sutar 2016	No measure of variance given	Not reported	Not reported
Yang 2016	Included	Included	Not reported
Yonglitthipagon 2017	Included	Not reported	Not reported