

Additional file 2

Table 1: Self-Reported Provision of Smoking Cessation Care compared to the Overall National Survey Sample, n(%)

| Smoking Cessation Care Component - Proportion reporting 'Often/Always' performing each component | GPs participating in the qualitative interviews (n=16) | GPs and Obstetricians participating in the national sample (n=378) |
|---|---|---|
| <i>Ask about smoking status</i> | 16 (100%) | 288 (77.2%) |
| <i>Give brief advise to quit if smoking</i> | 16 (100%) | 275 (74.7%) |
| <i>Assess nicotine dependence</i> | 14 (87.5%) | 89 (24.1%) |
| <i>Provide Cessation support to smokers (Assist)</i> | 15 (92.8%) | 124 (33.5%) |
| <i>Follow-up within 2 weeks (Arrange)</i> | 6 (37.5%) | 26 (7%) |
| <i>Prescribe/recommend NRT to assist quitting</i> | 9 (56.3%) | 40 (10.8%) |
| <i>Discuss the psychosocial context of smoking</i> | 13 (81.3%) | 82 (22.2%) |
| <i>Refer to Quit line/specialist service</i> | 8 (50%) | 95 (26.8%) |
| <i>Involve family members in counselling/tobacco management</i> | 4 (25%) | 15 (4.1%) |

Table 2: Barriers and Enablers to Provision of Smoking Cessation Care to Pregnant Smokers, compared to the Overall National Survey Sample, n(%)

| TDF domains –percentage answering ‘Agree/Strongly Agree’ n(%) | GPs participating in the qualitative interviews (n=16) | GPs and Obstetricians participating in the national sample (n=378) |
|--|---|---|
| <i>I am confident that I can counsel women about their smoking during pregnancy</i> | 15 (93.8%) | 299 (80.8%) |
| <i>I am confident that I can prescribe NRT for pregnant smokers</i> | 13 (81.3%) | 201 (54.5%) |
| <i>I am optimistic my intervention for smoking during pregnancy is likely to be effective</i> | 6 (37.5%) | 129 (35.1%) |
| <i>Raising the issue of smoking with a client during pregnancy will benefit our relationship</i> | 9 (56.3%) | 232 (65.2%) |
| <i>Addressing smoking during pregnancy is a high priority</i> | 16 (100%) | 351 (98.3%) |
| <i>I am comfortable raising the issue of smoking with a pregnant woman</i> | 15 (93.8%) | 338 (94.9%) |
| <i>In my workplace, it is routine to help pregnant women to quit smoking during pregnancy</i> | 15 (93.8%) | 271 (76.1%) |
| <i>I have sufficient time to help pregnant women to quit smoking</i> | 10 (62.5%) | 146 (41.1%) |
| <i>I have sufficient resources to help pregnant women to quit smoking</i> | 9 (56.3%) | 169 (47.5%) |