

1 What is the best method? Recovery of methicillin-resistant Staphylococcus
2 aureus and extended-spectrum beta-lactamase producing Escherichia coli
3 from inanimate hospital surfaces

4
5 Tânia Claro, PhD¹#; Sandra Galvin, PhD¹; Orla Cahill, PhD²; Deirdre
6 Fitzgerald-Hughes, PhD¹; Stephen Daniels, PhD²; Hilary Humphreys, MD^{1,3}.

7
8 ¹Department of Clinical Microbiology, Education and Research Centre, Royal
9 College of Surgeons in Ireland, Beaumont Hospital, Dublin 9, Ireland

10 ²School of Electronic Engineering and National Centre for Plasma Science
11 Technology, Dublin City University, Dublin 9, Ireland.

12 ³Department of Microbiology, Beaumont Hospital, Dublin 9, Ireland

13
14 #Corresponding author. Address: Department of Clinical Microbiology, Education
15 and Research Centre, Royal College of Surgeons in Ireland, Beaumont Hospital,
16 Dublin 9, Ireland; Telephone: +35318093748; Fax: +3538093709; E-mail:
17 tpedrosaclaro@rcsi.ie.

18
19 Preliminary data arising from this study were presented at the International
20 Healthcare Infection Society meeting in Liverpool, UK, November, 2012.

21
22 Running title: Detection of bacteria from hospital surfaces

23 Word count: 1199 words

25 ABSTRACT

26 Narrative abstract (50 words max)

27 Environmental sampling in hospitals, when required, needs to be reliable. We
28 evaluated different methods of sampling methicillin-resistant *Staphylococcus*
29 *aureus* and extended-spectrum beta-lactamase producing *Escherichia coli* on
30 five materials of the hospital setting. Petrifilms and contact-plates were superior
31 to swabs for all of the surfaces studied.

32

33 Keywords: MRSA, ESBL-*E. coli*, environmental contamination, hospital
34 surfaces, swabs, contact-plates, petrifilms.

35

36

37

38

39

40

41

42

43

44

45

46

47

Contamination of hospital surfaces by nosocomial microorganisms has long been recognized.¹ Transmission of these microorganisms occurs directly or indirectly, e.g. via contaminated healthcare workers' gloves.² Inadequate cleaning has also been associated with the transmission of healthcare-associated infections (HCAI).³ Furthermore, the use of novel decontamination methods such as gaseous plasma requires reliable methods to monitor their microbiological effectiveness. The Center for Disease Control provide recommendations for evaluating hospital environmental cleaning however, these are not standard and they do not consider specific outbreak scenarios or other research driven investigations.⁴ Sampling of the healthcare environment has been inconsistently reported using a variety of swabs (cotton, rayon or nylon flocked) or contact-plates.^{5,6} Although common in the food industry, petrifilms have rarely been used in assessing environmental contamination within the healthcare setting.⁷

We compared the recovery and limit of detection (LoD) of two important HCAI pathogens, i.e. methicillin-resistant *Staphylococcus aureus* (MRSA) and extended spectrum β -lactamase producing *Escherichia coli* (ESBL-E), from different materials commonly found in hospitals using swabs, contact-plates and petrifilms.

METHODS

Clinical strain 31(ST22-MRSA-IV) and ESBL-E clinical strain (CL2), both from our collection were used in this study.^{8,9} Columbia blood agar (CBA) and ESBL Brilliance agar (Oxoid Ltd, UK) were used for MRSA and ESBL-E, respectively.

71 Cultures were grown aerobically overnight at 37°C with rotation in tryptic-soy
72 broth (TSB) supplemented with 5% NaCl and Mueller-Hinton (MH) broth (Sigma
73 Aldrich, Ireland) for MRSA and ESBL-E, respectively. Cultures were centrifuged
74 for 10min at 15,500g and washed three times in phosphate buffered saline
75 (PBS). The culture was adjusted to the density of a 0.5McFarland standard
76 (approximately 1×10^8 MRSA and 3.2×10^7 ESBL-E CFU/mL) in PBS using a
77 DensiChek™ colorimeter (Vitek). Serial dilutions (10^{-1} to 10^{-6}) were prepared in
78 PBS and 50µl of each (including the original suspension) were applied to the
79 sterile test surfaces. These were sections (25cm²) of; linoleum flooring (Forbo
80 Flooring, Ireland), polyurethane mattress fabric (Meditec Medical, Ireland)
81 provided by the Maintenance Department, Beaumont Hospital, Dublin,
82 polypropylene (GoodFellow Cambridge Ltd., UK), powder-coated mild steel
83 (Watermark Engineering, Ireland) and stainless steel. Test sections were washed
84 for 30min (1% virkon solution for linoleum and mattress and 70% ethanol for
85 powder coated mild steel, polypropylene and stainless steel) before placing in
86 Petri-dishes under UV light for 30min before bacterial inoculation.

87 Inoculated sections were air dried in a laminar flow cabinet over 1h before
88 recovery. The recovery of bacteria from surfaces was assessed using rayon and
89 nylon flocked eSwabs (Copan, Italy), contact-plates - MRSA Chromagar (Cruinn
90 Diagnostics Ltd, Ireland) selective for MRSA, Brilliance UTI agar (Oxoid Ltd, UK)
91 selective for ESBL-E and 3M™Petrifilms; Staph Express Count for MRSA and
92 Enterobacteriaceae Count for ESBL-E (3M Petrifilm Trafalgar Scientific, UK).
93 Swabs were pre-moistened in PBS and the entire section then targeted. Swabs

were placed in 3mL of PBS in round-bottomed tubes and briefly vortex mixed. Serial dilutions were prepared to confirm the total viable count (TVC). Aliquots (10µl,100µl) of each suspension were spread onto CBA for MRSA and ESBL Brilliance agar for ESBL-E. Sterile contact-plates (65mm x 15mm) (VWR®) were poured with either MRSA Chromoagar (MRSA selective) or UTI brilliance agar (E. coli selective), dried and applied to the inoculated sections for 20 to 30s, ensuring firm contact with the surface. Petrifilms were prepared according to the supplier instructions. Briefly 1mL of sterile water was added to each petrifilm before storage at 4°C for 2h before use. Petrifilms were applied ensuring that the entire material section was covered. Sub-cultured plates from swabs, contact-plates and petrifilms were incubated at 37°C overnight. Colony enumeration was performed macroscopically the following day. The limit of detection (LoD) was defined as the lowest concentration of bacteria applied that was detected by a specific method.

Statistical data analysis was carried out using GraphPad Prism 5.00 software. The means of the $\log_{10}(\text{CFU/mL})$ recovered between methods or between materials was compared by one-way analysis of variance (ANOVA). When significant, i.e. $p < .05$, further analysis on the variance of the means between methods was carried out by Tukey's multiple comparison test. Comparison of the recoveries between microorganisms was analysed by t-test.

RESULTS

Figure 1 shows the recovery of MRSA from five materials, using four methods, at

different applied inocula (8 to 1 log₁₀CFU/mL). Petrifilms were best in recovering MRSA from all materials except stainless steel, when contact-plates were better. After petrifilms, contact-plates were best for recovering MRSA from linoleum, powder-coated mild steel and polypropylene. The second best method for stainless steel was the petrifilm and for mattress the flocked swabs. For all surfaces, rayon swabs had the lowest recovery of MRSA, compared to petrifilms (**P<.01). The LoD for MRSA was of 1x10² CFU/ml for all materials except linoleum (LoD of 1x10⁴ CFU/ml).

Figure 2 shows the recovery of ESBL-E from five materials, using four methods at different inocula concentrations (7.5 to 1.5 log₁₀CFU/mL). Petrifilms were best in recovering ESBL-E from all the materials followed by contact-plates. The lowest recovery of ESBL-E from all materials was with rayon and flocked swabs. Petrifilms and contact-plates were significantly better than either swab (**P<.001). The LoD for ESBL-E from petrifilms was 3.2x10¹ CFU/ml for powder-coated mild steel, polypropylene and stainless steel, and 3.2x10² CFU/ml and 3.2x10³ CFU/ml from mattress and linoleum, respectively.

The poorest recovery of MRSA and ESBL-E was with swabs. However, the recovery of MRSA using rayon and flocked swabs was significantly higher compared with ESBL-E from all surfaces (**P<.01 and ***P<.001, respectively). The recovery of MRSA from powder-coated mild steel, polypropylene and stainless steel was greater than from the mattress and linoleum but this difference was not statistically significant. Similarly, ESBL-E recovery was not significantly different among the different materials.

DISCUSSION

To our knowledge, this is the first study to evaluate petrifilms for assessing hospital surface contamination and compare it with other commonly used methods. Like contact-plates, petrifilms are a direct contact method advantageous over the aforementioned on its flexibility to adjust to non-flat surfaces (e.g. door handles). Interestingly, from those evaluated here, the petrifilm was the overall best method to recover both MRSA and ESBL-E from all surfaces tested followed by contact-plates. A previous study showed that Gram-positive contamination of the environment adjacent to 54 patients with Gram-positive infections versus 136 with Gram-negative infections was more heavily contaminated by Gram-positives (24.7%) than Gram-negatives (4.9%).¹⁰ Our findings suggest that contamination by Gram-negatives may be underestimated in studies where direct contact methods were not used. We show that the recovery of MRSA and ESBL-E differ significantly according to the method used and the type of surface being screened. Particularly, for MRSA, recovery was lower from linoleum and mattress compared to other surfaces, possibly due to the high porosity of these materials allowing bacteria to penetrate and being harder to culture.

This study limitation includes not evaluating the materials characteristics (e.g. porosity, roughness, hydrophobicity), bacteria were enumerated per volume rather than per area, and surfaces were free of protein unlike often in practice.

In conclusion, notwithstanding financial considerations, we suggest that direct contact methods, i.e. petrifilms and contact-plates and not swabs, are best

163 for the detection of MRSA and ESBL-E in the healthcare environment. They are
164 more rapidly processed than swabs and can be used as appropriate to the
165 surface type. Additional work is needed to confirm these findings in the actual
166 hospital environment.

167 ACKNOWLEDGEMENTS

168
169 Financial support. This study was funded by a Translational Research Award
170 from Science Foundation Ireland and the Health Research Board (TRA/2010/10).
171 Potential conflicts of interest. H.H. has recent research collaborations with
172 Steris Corporation, Inov8 Science, Pfizer & Cepheid and has also received
173 lecture & other fees from Novartis, AstraZenca & Astellas. All other authors
174 declare no potential conflict of interest.

Figure 1





Figure 2



185

186 REFERENCES

187

- 188 1. Hota B. Contamination, disinfection, and cross-colonization: are hospital
189 surfaces reservoirs for nosocomial infection? *Clin. Infect. Dis.*
190 2004;39(8):1182-1189.
- 191 2. Bhalla A, Pultz NJ, Gries DM, et al. Acquisition of nosocomial pathogens
192 on hands after contact with environmental surfaces near hospitalized
193 patients. *Infect. Control Hosp. Epidemiol.* 2004;25(2):164-167.
- 194 3. Dancer SJ. The role of environmental cleaning in the control of hospital-
195 acquired infection. *J. Hosp. Infect.* 2009;73(4):378-385.
- 196 4. Sehulster L, Chinn RY, Cdc, Hicpac. Guidelines for environmental
197 infection control in health-care facilities. Recommendations of CDC and
198 the Healthcare Infection Control Practices Advisory Committee (HICPAC).
199 *MMWR Recomm. Rep.* 2003;52(RR-10):1-42.
- 200 5. Dalmaso G, Bini M, Paroni R, Ferrari M. Qualification of high-recovery,
201 flocked swabs as compared to traditional rayon swabs for microbiological
202 environmental monitoring of surfaces. *PDA J. Pharm. Sci. Technol.*
203 2008;62(3):191-199.
- 204 6. Lutz JK, Crawford J, Hoet AE, Wilkins JR, 3rd, Lee J. Comparative
205 performance of contact plates, electrostatic wipes, swabs and a novel
206 sampling device for the detection of *Staphylococcus aureus* on
207 environmental surfaces. *J. Appl. Microbiol.* 2013;115(1):171-178.

- 208 7. Hooker EA, Allen S, Gray L, Kaufman C. A randomized trial to evaluate a
209 launderable bed protection system for hospital beds. *Antimicrob Resist*
210 *Infect Control*. 2012;1(1):27.
- 211 8. Burke L, Humphreys H, Fitzgerald-Hughes D. The revolving door between
212 hospital and community: extended-spectrum beta-lactamase-producing
213 *Escherichia coli* in Dublin. *J. Hosp. Infect.* 2012;81(3):192-198.
- 214 9. McNicholas WHJS, Humphreys H, Fitzgerald-Hughes D. Susceptibility of
215 *Staphylococcus aureus* Isolates Causing Bloodstream Infection to the
216 Antimicrobial Peptide LL-37. 2013;3(1):80-83.
- 217 10. Lemmen SW, Hafner H, Zolldann D, Stanzel S, Lutticken R. Distribution of
218 multi-resistant Gram-negative versus Gram-positive bacteria in the
219 hospital inanimate environment. *J. Hosp. Infect.* 2004;56(3):191-197.
220
221

222 FIGURE LEGENDS

223

224 Figure 1 – Numbers of MRSA inoculated and recovered using rayon and flocked
225 swabs, contact-plates and petrifilm from surfaces common to the hospital
226 environment. Each data point represents the mean of at least three individual
227 assays $n \geq 3$, error bars represent the standard error of the mean (SEM).

228

229 Figure 2 – Numbers of ESBL E. coli inoculated and recovered using rayon and
230 flocked swabs, contact-plates and petrifilm from surfaces common to the hospital
231 environment. Each data point represents the mean of at least three individual
232 assays $n \geq 3$, error bars represent the standard error of the mean (SEM).