



RCSI

UNIVERSITY
OF MEDICINE
AND HEALTH
SCIENCES

Royal College of Surgeons in Ireland

repository@rcsi.com

Systematic review and individual patient data meta-analysis of sex differences in depression and prognosis in persons with myocardial infarction: a MINDMAPS study

AUTHOR(S)

Frank Doyle, Hannah McGee, Ronán Conroy, Henk Jan Conradi, Anna Meijer, Richard Steeds, Hiroshi Sato, Donna E. Stewart, Kapil Parakh, Robert Carney, Kenneth Freedland, Matteo Anselmino, Roxanne Pelletier, Elisabeth H. Bos, Peter de Jonge

CITATION

Doyle, Frank; McGee, Hannah; Conroy, Ronán; Conradi, Henk Jan; Meijer, Anna; Steeds, Richard; et al. (2015): Systematic review and individual patient data meta-analysis of sex differences in depression and prognosis in persons with myocardial infarction: a MINDMAPS study. Royal College of Surgeons in Ireland. Journal contribution. <https://hdl.handle.net/10779/rcsi.10773272.v1>

HANDLE

[10779/rcsi.10773272.v1](https://hdl.handle.net/10779/rcsi.10773272.v1)

LICENCE

CC BY-NC-SA 4.0

This work is made available under the above open licence by RCSI and has been printed from <https://repository.rcsi.com>. For more information please contact repository@rcsi.com

URL

https://repository.rcsi.com/articles/journal_contribution/Systematic_review_and_individual_patient_data_meta-analysis_of_sex_differences_in_depression_and_prognosis_in_persons_with_myocardial_infarction_a_MINDMAPS_study/10773272/1

Systematic review and individual patient data meta-analysis of sex differences in depression and prognosis in persons with myocardial infarction: a MINDMAPS study

Running head: Sex differences in depression post-MI

Frank Doyle PhD¹, Hannah McGee PhD², R Conroy DSc³, Henk Jan Conradi PhD⁴, Anna Meijer MSc⁵, Richard Steeds MD⁶, Hiroshi Sato PhD⁷, Donna E. Stewart MD⁸, Kapil Parakh PhD⁹, Robert Carney PhD¹⁰, Kenneth Freedland PhD¹¹, Matteo Anselmino PhD¹², Roxanne Pelletier PhD¹³, Elisabeth H. Bos PhD¹⁴, Peter de Jonge PhD¹⁵.

Manuscript word count: 5987 (excluding authors and affiliations), including 4 Tables and 1 Figure

Affiliations

1. Frank Doyle, **Division of Population Health Sciences (Psychology)**, Royal College of Surgeons in Ireland: This author takes responsibility for all aspects of the reliability and freedom from bias of the data presented and their discussed interpretation
2. Hannah McGee, **Faculty of Medicine & Health Sciences**, Royal College of Surgeons in Ireland: This author takes responsibility for all aspects of the reliability and freedom from bias of the data presented and their discussed interpretation

3. Ronán Conroy, **Division of Population Health Sciences (Epidemiology & Public Health Medicine)**, Royal College of Surgeons in Ireland: This author takes responsibility for all aspects of the reliability and freedom from bias of the data presented and their discussed interpretation
4. Henk Jan Conradi, **Department of Clinical Psychology**, University of Amsterdam: This author takes responsibility for all aspects of the reliability and freedom from bias of the data presented and their discussed interpretation
5. Anna Meijer MSc, **Interdisciplinary Center Psychopathology and Emotion regulation**, University Medical Center Groningen: This author takes responsibility for all aspects of the reliability and freedom from bias of the data presented and their discussed interpretation
6. Richard Steeds, **Department of Cardiology**, Queen Elizabeth Hospital: This author takes responsibility for all aspects of the reliability and freedom from bias of the data presented and their discussed interpretation
7. Hiroshi Sato, **School of Human Welfare Studies**, Kwansai Gakuin University: This author takes responsibility for all aspects of the reliability and freedom from bias of the data presented and their discussed interpretation
8. Donna E. Stewart, **University Health Network**, University of Toronto: This author takes responsibility for all aspects of the reliability and freedom from bias of the data presented and their discussed interpretation

9. Kapil Parakh, **Department of Medicine**, Johns Hopkins University School of Medicine: This author takes responsibility for all aspects of the reliability and freedom from bias of the data presented and their discussed interpretation
10. Robert Carney, **Department of Psychiatry**, Washington University: This author takes responsibility for all aspects of the reliability and freedom from bias of the data presented and their discussed interpretation
11. Kenneth Freedland, **Department of Psychiatry**, Washington University: This author takes responsibility for all aspects of the reliability and freedom from bias of the data presented and their discussed interpretation
12. Matteo Anselmino, **Division of Cardiology, Department of Medical Sciences**, University of Turin: This author takes responsibility for all aspects of the reliability and freedom from bias of the data presented and their discussed interpretation
13. Roxanne Pelletier, McGill University **Health Centre**: This author takes responsibility for all aspects of the reliability and freedom from bias of the data presented and their discussed interpretation
14. Elisabeth H. Bos, **Interdisciplinary Center Psychopathology and Emotion regulation**, University Medical Center Groningen: This author takes responsibility for all aspects of the reliability and freedom from bias of the data presented and their discussed interpretation
15. Peter de Jonge, **Interdisciplinary Center Psychopathology and Emotion regulation**, University Medical Center Groningen: This author takes

responsibility for all aspects of the reliability and freedom from bias of the data presented and their discussed interpretation

Corresponding author:

Frank Doyle, PhD

Division of Population Health Sciences (Psychology),

Royal College of Surgeons in Ireland,

123 St Stephen's Green,

Dublin 2,

Ireland

Email: fdoyle4@csi.ie, Tel: + 353 1 4022718, Fax: + 353 1 4022764

Funding: A.M., H.J.C., and P. de J. have support from a Vidi grant from the Dutch Medical Research Council (grant no. 016.086.397)

Conflict of interest disclosures

Disclosures and grant support

Frank Doyle, PhD (No conflicts of interest or disclosures)

Hannah McGee PhD (No conflicts of interest or disclosures)

Ronan Conroy DSc (No conflicts of interest or disclosures)

Elisabeth H. Bos PhD (No conflicts of interest or disclosures)

Anna Meijer MSc (No conflicts of interest or disclosures)

Richard Steeds MD (No conflicts of interest or disclosures)

Hiroshi Sato PhD (supported by a grant-in-aid for University and Society Collaboration (15590743) from the Japanese Ministry of Education, Culture, Sports, Science, and Technology (Tokyo, Japan) and by a research grant from the Japan Arteriosclerosis Prevention Fund (Tokyo, Japan))

Donna E. Stewart MD (Potential conflicts are a grant from Canadian Institute of Health Research on Gender, Mental Health and Violence, and membership on the Duloxetine and Pregnancy Registry Scientific Advisory Board of Eli Lilly Inc. I also received travel support from Ranbaxy Pharma for a psychiatric meeting in Thailand)

Kapil Parakh PhD (No conflicts of interest or disclosures)

Robert Carney PhD (Dr. Carney or a member of his family own stock in Pfizer Inc and Forest Laboratories.)

Kenneth Freedland PhD (No conflicts of interest or disclosures)

Matteo Anselmino PhD (No conflicts of interest or disclosures)

Roxanne Pelletier PhD (no conflict of interest to disclose and I am funded by the Canadian Institutes of Health Research)

Henk Jan Conradi PhD (No conflicts of interest or disclosures)

Peter de Jonge PhD (No conflicts of interest or disclosures)

Keywords

Keywords– Depression; myocardial infarction; gender differences; prognosis; individual patient data meta-analysis

The MIND-MAPS Group:

E.H. Bos*, H.J. Conradi**, P. de Jonge*, A. Meijer*, J. Ormel*

* University of Groningen, University Medical Center Groningen, Interdisciplinary Center Psychopathology and Emotion regulation, 9713 GZ, Groningen, The Netherlands ** Department of Clinical Psychology, University of Amsterdam, 1018 XA, Amsterdam, The Netherlands

S. Bergerone, M. Anselmino

Division of Cardiology, Department of Medical Sciences, University of Turin, “Città della Salute e della Scienza” Hospital, Turin, Italy

L.F. Berkman*, R. Carney**, K.E. Freedland**

* Centre for Population and Development Studies, Harvard University, Boston, Massachusetts 02138, USA, ** Department of Psychiatry, Washington University, St. Louis, Missouri 63108, USA

J. Denollet

Centre of Research on Psychology in Somatic Diseases, Tilburg University, 5000 LE, Tilburg, The Netherlands

F. Doyle*, H.M. McGee*, Ronán Conroy**

* Division of Population Health Sciences (Psychology), Royal College of Surgeons in Ireland, Dublin 2, Ireland; ** Division of Population Health Sciences (Epidemiology & Public Health Medicine), Royal College of Surgeons in Ireland, Dublin 2, Ireland

S.L. Grace

Women’s Health Program, University Health Network; Faculty of Medicine, University of Toronto; School of Kinesiology and Health Science, York University, Toronto, Ontario M3J1P3, Canada

S.H. Hosseini

Psychiatry and Science Behavior Research Centre, Department of Psychiatry, Mazandaran University of Medical Sciences, Sari, 48171-19164, Iran

D.A. Lane

University of Birmingham Centre for Cardiovascular Sciences, City Hospital, Birmingham B18 7QH, UK

L. Pilote

Divisions of Epidemiology and Cardiology, McGill University Health Centre, Montréal H3A1A1, Canada

K. Parakh, R.C. Ziegelstein

Department of Medicine, Johns Hopkins University School of Medicine, Johns Hopkins Bayview Medical Centre, Baltimore, Maryland 21224, USA

C. Rafanelli

Department of Psychology, University of Bologna, 40127 Bologna, Italy

H. Sato

School of Human Welfare Studies, Kwansai Gakuin University, Nishinomiya, 662-8501 Japan

R.P. Steeds

Department of Cardiology, Queen Elizabeth Hospital, Birmingham B15 2TH, UK

B.D. Thombs

Lady Davis Institute for Medical Research, Jewish General Hospital and McGill University, Montréal H3T 1E4, Canada

C. Welin

College of Health Sciences, Department of Nursing, Göteborg University, SE-405 30 Göteborg, Sweden

Abstract

Objective—Using combined individual patient data (IPD) from prospective studies, we explored sex differences in depression and prognosis post-myocardial infarction (MI), and determined whether disease indices could account for found differences.

Methods—Meta-analysis of IPD from 10,175 MI patients who completed diagnostic interviews or depression questionnaires from 16 prospective studies of MI patients, identified by systematic review for the MINDMAPS study. Multilevel logistic and Cox regression models were used to determine sex differences in prevalence of depression and sex-specific effects of depression on **subsequent cardiovascular morbidity and all-cause mortality**.

Results—Combined interview and questionnaire data from observational studies showed that 36% (635/1760) of women and 29% (1575/5526) of men reported elevated levels of depression (age-adjusted OR=0.68, 95% CI 0.60 to 0.77, $p<0.001$). The risk for all-cause mortality associated with depression was higher in men (HR=1.38, 95% CI 1.30 to 1.47, $p<0.001$) than in women (HR=1.22, 95% CI 1.14 to 1.31, $p<0.001$) (**sex*depression interaction $p<0.001$**). Low left ventricular ejection fraction (LVEF) was associated with higher depression scores in men only (sex*LVEF interaction B=0.294, 95% CI 0.090 to 0.498, $p=0.005$), which attenuated the sex difference in the association between depression and prognosis.

Conclusions— The prevalence of depression post-MI was higher in women than men, but the association between depression and cardiac prognosis was worse for men. LVEF was associated with depression in men only, and accounted for the increased risk of all-cause

mortality in depressed men versus women, suggesting that depression in men post-MI may in part reflect cardiovascular disease severity.

Keywords: Depression; myocardial infarction; gender; prognosis; individual data meta-analysis

Acronyms

ACE = angiotensin converting enzyme

BMI – body mass index

CABG = coronary artery bypass graft

CI = confidence interval

CVE – cardiovascular events

HF = heart failure

HR = Hazard ratio

IPD = Individual patient data

LVEF = Left ventricular ejection fraction

MI = myocardial infarction

OR = odds ratio

PTCA = percutaneous transluminal coronary intervention

SD = standard deviation

Introduction

Depression in people with myocardial infarction (MI) has been the source of considerable study. While both clinically diagnosed depression and questionnaire-assessed depressive symptoms have been consistently associated with poorer cardiovascular prognosis in MI patients (1), the treatment of depression has not been demonstrated to reduce morbidity or mortality in randomized trials (2, 3). Recent estimates suggest a prevalence of diagnosed major depressive disorder in the US of 2.1% to 4.9% in adults aged 45 and over (4), with women having a higher prevalence (4.8%) than men (3.3%). Similarly, in adults aged 45 years and above, estimates of elevated depressive symptoms are also more prevalent among women (7-12%) than men (5-7%) (5). Similar findings are available for European countries (6, 7), although the differing methodologies adopted by different countries make overall estimates for major depression more difficult. According to a systematic review of the literature, the prevalence of depression is significantly higher in patients with MI, with 20% having major depression and up to 31% reporting elevated depressive symptoms, although sex differences were not reported (8).

The etiology of depression post-MI has been the subject of considerable debate. While depressive symptoms and trajectories of these symptoms post-MI appear to have similar risk factors to depression in the general population (e.g. stressful life events, personality etc.) (9-13), sex is not always a predictor of depression in these studies. This suggests that there may be a potentially different etiology for depressive symptoms in men and women with MI. Indeed, other researchers have demonstrated that elevated levels of depression, and its subsequent association with cardiovascular prognosis, may be a consequence of

cardiovascular disease severity (14, 15). For example, poor left ventricular ejection fraction (LVEF) has been demonstrated to not only correlate with depression, but also to predict the onset of major depression and depressive symptoms in patients with MI (14). In addition, this study found that poorer LVEF was a stronger predictor of onset of major depression for men than for women (14), again suggesting potentially different etiology for depression between men and women post-MI.

We aimed to establish whether the prevalence of diagnosed depression and elevated depressive symptomatology is similar in men and women with MI, and whether the diagnosed depression or depressive symptoms had a greater impact on prognosis for men than for women. We also investigated the role of LVEF in the etiology of depression and prognosis according to patients' sex. These questions have not been subject to systematic investigation and previous studies were not large enough to detect potential moderating effects. We therefore used data from the MINDMAPS individual patient meta-analytic database of 16 studies (16) to determine the following:

- 1 – whether cardiovascular disease severity indices account for differences in post-MI prevalence of depressive symptoms between men and women; and
- 2 – whether the association between depression and prognosis is stronger in men than in women, and whether disease severity indices (e.g. LVEF) account for these differences.

Methods

MINDMAPS

Details of MINDMAPS are available elsewhere (16). In brief, eligibility for inclusion in the MINDMAPS dataset was determined by a systematic review of published studies on post-MI depression **initiated on 5th January 2011** (within 3 months post-hospitalisation for MI, assessed by validated questionnaire or diagnostic interview) and two prognostic endpoints (all-cause mortality, or major cardiovascular events – CVE: recurrent MI, unstable angina, CABG) (1). Authors of all selected studies were contacted and original data was requested (covering demographics, depression, disease severity [*LVEF*<40%, *Killip class (I vs. II, III, IV), and history of MI*], comorbidities, diabetes, smoking, body mass index [BMI], medication use, and outcomes). Of 30 eligible studies, 16 provided data (16).

Statistical analysis

Stata 11 was used for the main analyses (Statacorp LP, TX, USA). Multilevel logistic regression, adjusting for age, was used to obtain odds ratios (ORs) for differences in depression status between men (1) and women (0), using a binary depression variable (based on interview or recommended scale thresholds). Hazard ratios (HRs) were calculated using multilevel Cox proportional hazards regression analysis, to predict time-to-event (all-cause mortality or CVE), using depression, sex, and the interaction between sex and depression as predictors, adjusting for age. A continuous depression variable was used in these Cox analyses (z-transformed total scale scores). As different studies used different methods and are likely to represent populations with different characteristics (such as date of study, differences in therapies and assessments used, etc.), observations within studies are unlikely to be fully independent. Therefore, a random intercept for

study was included, resulting in a multilevel model. When calculating the prevalence estimates we omitted data from the ENRICHD randomized trial and ancillary studies (2, 17) (which were combined in the original study (16)), as these were enrolled under strict inclusion criteria whereas observational studies would be more inclusive. To demonstrate the random effect variance in respect to the depression prevalence, we displayed the intraclass correlation coefficient, which shows what part of the total variance in prevalence is between studies. To investigate whether cardiovascular disease indices and other risk factors accounted for any sex differences in depression, a linear mixed model with random intercept was used, with depression z-scores as the dependent variable, and sex, disease indices and their interaction as predictors, adjusting for age. For each disease index a separate model was built. A contrast coding of -0.5/0.5 for dichotomous variables was used in the Cox regression analyses to ensure between-study variances between groups were equal (e.g. female vs. male) (18). Separate models were built for each disease and risk factor index. Also, two more multivariate models were built – including all disease and risk factors, but omitting LVEF, together as covariates, and then a final model including all disease and risk factors. To increase confidence interval robustness, and in view of the non-normal distribution of the depression z-scores, bootstrapping with 1000 replications was used in all analyses (19). The handling of the variables used for adjustment and the applied outcomes was identical to that in the main MINDMAPS study (16). All tests were two-tailed.

Results

Sample description

Of 30 eligible studies (16), 16 provided individual patient data (**see supplementary Table S1 for an overview of participating studies**). These studies were from 9 different countries. The description of the combined sample (N=10,175), stratified by sex, is displayed in Table 1:

Insert Table 1 here

Overall, the mean age was 61 years (range 20-97), 71% of participants were men, and mean follow-up time per study was 3.15 years (range 0.96 – 6.65).

Prevalence of depression

Data from four diagnostic interview studies (omitting the ENRICHD data, n=900) showed that 20% (47/240) of women and 12% (76/660) of men had major depression. The age-adjusted multilevel logistic regression showed this to be a significant difference (OR=0.42, 95% CI 0.27 to 0.65, p=0.001). There was little heterogeneity between these interview studies in depression prevalence: the intraclass correlation coefficient was 0.01 (Chi² = 1.4; **p=0.12**).

When using the data from the questionnaire studies, excluding data from the ENRICHD and ancillary studies (fourteen studies; n=7202), the sex differences in prevalence remained significant (age-adjusted OR=0.69, 95% CI 0.61 to 0.77, p<0.001), with a

higher percentage of women scoring above recommended thresholds for “clinically significant” depressive symptoms (38%, 657/1741) than men (30%, 1622/5461). There was some heterogeneity between these questionnaire studies in depression prevalence: the intraclass correlation coefficient was 0.10 ($\text{Chi}^2(1) = 436.3$; $\mathbf{p < 0.001}$). **This implies that 10% of the total variance in depression prevalence is due to differences between studies.**

Combining the interview and questionnaire studies, but omitting the ENRICHHD data (fifteen studies; $n=7286$), resulted in 36% (635/1760) of women and 29% (1575/5526) of men reporting elevated depressive symptoms (age-adjusted $\text{OR}=0.68$, 95% CI 0.60 to 0.77, $\mathbf{p < 0.001}$). The intraclass correlation coefficient was 0.14 ($\text{Chi}^2(1) = 547.7$; $\mathbf{p < 0.001}$), indicating significant, though not large, heterogeneity between studies in depression prevalence.

We also checked whether the sex difference in depression prevalence varied with age. We did so by including an interaction term $\text{sex} * \text{age}$ in the above analysis. The interaction was not significant ($\text{OR}=1.00$, 95% CI 0.99 to 1.01, $\mathbf{p} = 0.91$). So, the sex difference in depression prevalence did not vary with age. **The main effect of age on the prevalence of depression, adjusted for sex, was small and not significant ($\text{OR}=0.99$, 95% CI = 0.99 to 1.00, $\mathbf{p} = 0.069$).**

Disease severity

Various disease indices and risk factors were analysed to determine if these accounted for sex differences in depression scores. Interactions between sex and history of MI, Killip class, diabetes, smoking or BMI were not associated with depression z-scores (**Supplementary Table S2**). However, an interaction between sex and left ventricular ejection fraction was observed (see the upper panel of Table 2):

Insert Table 2 here

This interaction remained significant when adjusting for all other disease factors and risk indices (see the lower panel of Table 2). The results suggest that for men only, low LVEF was associated with higher levels of depression.

Moderation effects of sex in the association between depression and prognosis

Survival curves for all-cause mortality and depression status, and cardiovascular events and depression status are shown in Fig 1:

Insert Fig 1 about here

The survival curves show that depression was related to poorer prognosis, in both men and women, and that the effect is more important in men. This is further investigated in Table 3:

Insert Table 3 here

The Cox regression analyses for all-cause mortality showed that the interaction between sex and depression was statistically significant (HR for interaction=1.12; 95% CI: 1.05 to 1.19, **p<0.001**), suggesting that the association between depression and mortality was indeed stronger for men than for women. The HR associated with depression was 12% higher in men compared to women (men, HR=1.38, 95% CI 1.30 to 1.47, p<0.001; women, HR=1.22, 95% CI 1.14 to 1.31, p<0.001). Omitting data from the ENRICHHD studies had little effect on the interaction term (HR for interaction=1.10, 95% CI 1.02 to 1.19, p=0.014). The interaction effect for CVE pointed into the same direction, but did not reach significance.

We also checked whether the sex difference in the impact of depression on cardiovascular outcomes varied with age. We did so by including an interaction term sex*age*depression in the above analyses. The interaction was not significant (all-cause mortality HR =1.00. 95% CI 0.99 to 1.01, p = 0.92; CVE: HR =1.00. 95% CI 0.99 to 1.00, p = 0.59; **Supplementary Table S2**). So, the sex difference in the impact of

depression on cardiovascular outcomes did not vary with age. **The sex*depression interaction now did reach significance also in the CVE model (p = 0.038).**

Because sex differences between depression and cardiac prognosis may be dependent on post-menopausal status (20, 21), we did a sensitivity analysis repeating the analysis of Table 3 but now separately for individuals aged younger and older than 55 years. This did not yield different results. The sex*depression interaction in the prediction of all-cause mortality was significant in both age groups (age<=55, HR=1.20, 95% CI 1.01 to 1.43, p=0.038; age>55, HR=1.11, 95% CI 1.03 to 1.20, p=0.005). The HR for the sex*depression interaction in the prediction of CVE pointed in the same direction but was non-significant in both age groups (age<=55, HR=1.15, 95% CI 0.94 to 1.41, p=0.18; age>55, HR=1.05, 95% CI 0.90 to 1.23, p=0.51).

Finally, we determined if the moderating effect of sex in the association between depression and all-cause mortality could be explained by the cardiovascular disease indices or risk factors. The same multilevel Cox regression analyses were used, predicting all-cause mortality by age, depression, sex, and sex*depression. Then, the size of the change in the estimate for the interaction sex*depression was determined when each of the disease variables was included in these separate models, and in two multivariate models. Results are shown in Table 4:

Insert Table 4 here

In the separate models, LVEF accounted for the largest reduction in the size of the sex*depression interaction, and inclusion of this variable resulted in a non-significant interaction. History of MI also accounted for a substantial reduction in the interaction, but most other factors *increased* instead of reduced the size of the interaction. These differences could be the result of differences between studies or sample sizes, therefore, the analyses were repeated with only those participants who had data on all six cardiovascular disease variables and risk indices (History of MI, LVEF, and Killip class, diabetes, smoking, BMI). The last rows of Table 4 show the results of the multivariate models. While adjusting for all covariates except LVEF (5172 participants in 5 studies) resulted in an 8% increase in the sex*depression interaction, inclusion of all six covariates (2226 participants in 3 studies) resulted in a 32.5% reduction of the sex*depression interaction effect, and resulted in a non-significant interaction. Together, these results suggest that the moderating effect of sex on the association between depression and all-cause mortality can partially be explained by differences in LVEF and history of MI among depressed men and women, and that LVEF explained the largest part **of the sex*depression interaction**.

Furthermore, despite there being no statistically significant sex*depression interaction **in the cardiac disease and risk factor-adjusted models** for predicting CVE, the percentage reduction **of the interaction term** was again largest for LVEF, showing a consistency across outcomes (results not shown).

Discussion

In this largest study to date, we found a number of results that are important in interpreting the association between post-MI depression and prognosis. Firstly, low LVEF was associated with depressive symptoms, but only in men. Secondly, the association between depression and prognosis was stronger in men than in women, which could partly be explained by depressive symptoms being a reflection of low LVEF more often for men than for women. LVEF attenuated the interaction effect between depression and sex in the prediction of prognosis.

In general, the prevalence of depression by diagnostic interview and questionnaire studies are similar to that previously reported (8), with higher proportions of women having depression according to both diagnostic interview and questionnaire studies. This higher prevalence of depression in women, added to the fact that depression was associated with disease severity in men only, demonstrates how depression in this population mimics depression in the general population. Other studies have also demonstrated how psychosocial risk factors for depression are predictors not only of elevated post-MI depressive symptomatology, but also of trajectories of depression post-discharge (9-13, 22), suggesting that depressive symptoms cannot be attributed solely to coronary disease severity. However, the higher prevalence of depression reported using scale thresholds rather than interview techniques again provides evidence that psychometric scales overestimate the prevalence of depression in this population, with our findings

demonstrating 2.5- to 3-fold higher prevalence, as found previously (8). This higher prevalence by questionnaire probably represents confounding between somatic symptoms of depression and symptoms of coronary disease (14, 15, 23), which cannot easily be thoroughly investigated as may happen during a diagnostic interview. Conversely, it has been demonstrated that even low levels of depression predict cardiovascular prognosis (24), thus relying on diagnostic interviews may actually underestimate meaningful levels of depression. Furthermore, while it has previously been suggested that the higher prevalence of depression in women might lead to poorer cardiovascular prognosis (25), we have found no evidence for this. Indeed, depression was more strongly related to a worse cardiac prognosis in men than in women. Single studies may not have had the power to detect such effects.

Depression was associated with a poor LVEF in men but not in women. Poor LVEF has been associated with depression in patients with coronary disease (26, 27). Previously, van Melle et al. (14) have shown that in 1989 post-MI patients LVEF was inversely correlated with depression status (from diagnostic interviews) and depression scale scores at several time points in the year post-MI, even in those who did not demonstrate depression at baseline. Furthermore, men demonstrated a trend towards a stronger increase in rates of depression across more severe LVEF categories, which was confirmed in the present findings. The data from that sample were included in the present analyses. However, that was a single study, while the present analyses are from 16 studies. That depression was more strongly related to prognosis for men may be explained by the differential nature of MI in women and men. Women with MI tend to be

older by approximately 10 years, have more diffuse coronary disease, have non-ST-elevation MI thereby not being eligible for emergency primary percutaneous coronary intervention (21, 28) which is protective of subsequent lowering of LVEF (29-31), and therefore have worse hospital outcomes (28, 29). Men tend to have more focal lesions that result in ST-elevation MI. In the current data, women were less likely to receive percutaneous coronary intervention. As a result, women are more likely to end up with heart failure with preserved ejection fraction while men end up with heart failure with reduced ejection fraction (29-31). Thus, the differential attenuation by LVEF of the association of depression with outcomes may be partly explained by the fact that women are less likely to have the type of MI that results in a low LVEF. Further research is needed to clarify this possibility. Alternatively, the etiology of depression in post-MI patients may differ in women and men. Men may be more likely to have depression as a result of coronary disease severity, while women may more often have pre-existing depression (32). These sex differences are perhaps unlikely to be mediated by other biological (e.g. reduced cardiac autonomic tone, increased platelet aggregation, increased inflammatory processes) (33, 34) or behavioral (e.g. smoking, physical inactivity etc.) (34-36) mechanisms linking depression to MI, as no consistent links with men or women have been reported. Recent research also suggests that fetal exposure may have increased the risk of both depression and cardiovascular disease, and that this effect is stronger in women (37), although this requires further investigation. **The lack of variation in results in the sensitivity analysis was perhaps surprising given that younger women have consistently been shown to have poorer prognosis than younger men (21). This may be due to the fact that there were relatively few young women in each study included**

here, and that these studies took place over a large timeframe, perhaps making them less representative than larger registry studies, or that using age thresholds is too crude to represent actual menopausal status. It is unclear why there was not a significant moderating effect of sex in the association between depression and CVE. One reason may be that CVE is a less well-defined endpoint across the studies, and this may lead to measurement error, or that there were different numbers and types of studies and participants analysed for this outcome.

Limitations and strengths

The heterogeneity of study populations, depression, cardiovascular risk factor measurements, and endpoint assessments may be problematic, although the inclusion of the random intercept accounts for study differences. Measures of distress have been shown to demonstrate measurement invariance across sex and ethnic groups in a large cross-sectional study (38). All-cause mortality includes non-cardiac deaths, and may bias the results. The heterogeneity of cardiovascular risk factors reported in individual studies meant that full adjustment could not be made in the overall sample. Inclusion of variables such as New York Heart Association categorisation, diastolic function, blood pressure, and cholesterol were not included in the dataset, but might have changed the results. Similarly, psychiatric history was unavailable, as was alcohol abuse, and other non-cardiac co-morbidities or their treatments may also impact on the findings. Not all relevant studies were available for analysis (16), which may have biased the results, although most did not report sex differences or did not find any differences in prognosis.

Another major limitation of the present research, and indeed other studies in this area, is the lack of a plausible model that could explain the effects seen in this study. Other researchers have rightly called for more conceptual and theoretical clarity in the analysis of observational studies which investigate risk factors in disease etiology (39, 40). To establish any single variable as a true independent risk factor, confounder or mediator for a given outcome, knowledge of temporal precedence of variables is required (39, 40). This is extremely difficult in the case of cardiovascular disease and depression. In the present analyses we cannot, for example, assume that depression precedes LVEF, or vice versa, as this data simply was not available to us. We have therefore avoided the terms ‘mediate’ or ‘confound’, etc. Further research with repeated measures of important variables is required before true risk factors and mediators of sex differences in depression and prognosis post-MI are elucidated.

The major strength of the study is the use of individual patient data, which allows for time-to-event analysis, and adjustment for potentially confounding variables systematically across studies, which cannot be done in summary meta-analyses, resulting in more accurate estimates of effect sizes, increased statistical power, generalizability and reliability of results (41). The inclusion of the random intercept term allows important between-study differences to be accounted for statistically, such as changing acute treatments (percutaneous coronary intervention has superseded thrombolysis in the timeframe spanned by the present studies) (29, 30), or differential assessment of LVEF –

which may impact on its prognostic ability (42, 43). The ability to analyse both interview and questionnaire studies is a strength, and the results demonstrated consistent patterns across these methods.

Conclusions

Using IPD from a large sample, the present study revealed significant differences in the prevalence, possible etiology and consequences of depression post-MI in men versus women. The prevalence of depression post-MI was higher in women than men, but the association between depression and cardiac prognosis was worse for men. LVEF was associated with depression in men only, and accounted for the increased risk of all-cause mortality in depressed men, suggesting that depression in men post-MI may in part reflect cardiovascular disease severity. Future studies, including intervention studies, should account for these sex differences.

References

1. Meijer A, Conradi HJ, Bos EH, Thombs BD, van Melle JP, de Jonge P. Prognostic association of depression following myocardial infarction with mortality and cardiovascular events: a meta-analysis of 25 years of research. *Gen Hosp Psychiatry*. 2011;33:203-16.
2. The ENRICHD Investigators. Effects of treating depression and low perceived social support on clinical events after myocardial infarction: the Enhancing Recovery in Coronary Heart Disease Patients (ENRICHD) Randomized Trial. *JAMA*. 2003;289:3106-16.
3. van Melle JP, de Jonge P, Honig A, Schene AH, Kuyper AM, Crijns HJ, Schins A, Tulner D, van den Berg MP, Ormel J. Effects of antidepressant treatment following myocardial infarction. *Br J Psychiat*. 2007;190:460-6.
4. Centers for Disease Control and Prevention. Morbidity and Mortality Weekly Report: Erratum. 2010 [cited 2012 08/06/2012]; Available from: http://www.cdc.gov/Features/dsDepression/Revised_Table_Estimates_for_Depression_MMWR_Erratum_Feb%202011.pdf.
5. Centers for Disease Control and Prevention. QuickStats: Prevalence of Current Depression* Among Persons Aged ≥ 12 Years, by Age Group and Sex—United States, National Health and Nutrition Examination Survey, 2007-2010. National Center for Health Statistics, 2012 Jan 6. Report No.: Contract No.: 60.
6. Wittchen HU, Jacobi F, Rehm J, Gustavsson A, Svensson M, Jonsson B, Olesen J, Allgulander C, Alonso J, Faravelli C, Fratiglioni L, Jennum P, Lieb R, Maercker A, van Os J, Preisig M, Salvador-Carulla L, Simon R, Steinhausen HC.

The size and burden of mental disorders and other disorders of the brain in Europe 2010. *Eur Neuropsychopharm.* 2011;21:655-79.

7. Van de Velde S, Bracke P, Levecque K. Gender differences in depression in 23 European countries. Cross-national variation in the gender gap in depression. *Soc Sci Med.* 2010;71:305-13.

8. Thombs BD, Bass EB, Ford DE, Stewart KJ, Tsilidis KK, Patel U, Fauerbach JA, Bush DE, Ziegelstein RC. Prevalence of depression in survivors of acute myocardial infarction. *J Gen Intern Med.* 2006;21:30-8.

9. Rieckmann N, Burg MM, Gerin W, Chaplin WF, Clemow L, Davidson KW. Depression vulnerabilities in patients with different levels of depressive symptoms after acute coronary syndromes. *Psychotherap Psychosom.* 2006;75:353-61.

10. Doyle F, McGee H, Conroy R, Delaney M. What predicts depression in cardiac patients: Sociodemographic factors, disease severity or theoretical vulnerabilities? *Psychol Health.* 2011;26:619-34.

11. Doyle F, McGee HM, Delaney M, Motterlini N, Conroy RM. Depressive vulnerabilities predict depression status and trajectories of depression over one year in persons with acute coronary syndrome. *Gen Hosp Psychiat.* 2011;33:224-31.

12. Kaptein KI, de Jonge P, van den Brink RH, Korf J. Course of depressive symptoms after myocardial infarction and cardiac prognosis: a latent class analysis. *Psychosom Med.* 2006;68:662-8.

13. Martens EJ, Smith OR, Winter J, Denollet J, Pedersen SS. Cardiac history, prior depression and personality predict course of depressive symptoms after myocardial infarction. *Psychol Med.* 2008;38:257-64.
14. van Melle JP, de Jonge P, Ormel J, Crijns HJ, van Veldhuisen DJ, Honig A, Schene AH, van den Berg MP. Relationship between left ventricular dysfunction and depression following myocardial infarction: data from the MIND-IT. *Eur Heart J.* 2005;26:2650-6.
15. Nicholson A, Kuper H, Hemingway H. Depression as an aetiologic and prognostic factor in coronary heart disease: a meta-analysis of 6362 events among 146 538 participants in 54 observational studies. *Eur Heart J.* 2006;27:2763-74.
16. Meijer A, Conradi HJ, Bos EH, Anselmino M, Carney R, Denollet J, Doyle F, Freedland KE, Grace SL, Hosseini SH, Lane DA, Pilote L, Parakh K, Rafanelli C, Sato H, Steeds RP, Welin C, de Jonge P. Adjusted prognostic association of depression following myocardial infarction with mortality and cardiovascular events: an individual patient data meta-analysis. *Br J Psychiat.* 2013;203:90-102.
17. Carney RM, Blumenthal JA, Stein PK, Watkins L, Catellier D, Berkman LF, Czajkowski SM, O'Connor C, Stone PH, Freedland KE. Depression, heart rate variability, and acute myocardial infarction. *Circulation.* 2001;104:2024-8.
18. Smith CT, Williamson PR, Marson AG. An overview of methods and empirical comparison of aggregate data and individual patient data results for investigating heterogeneity in meta-analysis of time-to-event outcomes. *J Eval Clin Pract.* 2005;11:468-78.

19. Sauerbrei W, Schumacher M. A bootstrap resampling procedure for model building: application to the Cox regression model. *Stat Med.* 1992;11:2093-109.
20. **Vaccarino V, Shah AJ, Rooks C, Ibeanu I, Nye JA, Pimple P, Salerno A, D'Marco L, Karohl C, Bremner JD, Raggi P. Sex differences in mental stress-induced myocardial ischemia in young survivors of an acute myocardial infarction. *Psychosom Med.* 2014;76:171-80.**
21. **Vaccarino V, Badimon L, Corti R, de Wit C, Dorobantu M, Manfrini O, Koller A, Pries A, Cenko E, Bugiardini R. Presentation, management, and outcomes of ischaemic heart disease in women. *Nat Rev Cardiol.* 2013;10:508-18.**
22. Sorensen C, Brandes A, Hendricks O, Thrane J, Friis-Hasche E, Haghfelt T, Bech P. Psychosocial predictors of depression in patients with acute coronary syndrome. *Acta Psychiatr Scand.* 2005;111:116-24.
23. Doyle F, Conroy R, McGee H. Challenges in reducing depression-related mortality in cardiac populations: cognition, emotion, fatigue or personality? *Health Psychol Rev.* 2007;1:137-72.
24. Bush DE, Ziegelstein RC, Tayback M, Richter D, Stevens S, Zahalsky H, Fauerbach JA. Even minimal symptoms of depression increase mortality risk after acute myocardial infarction. *Am J Cardiol.* 2001;88:337-41.
25. Naqvi TZ, Naqvi SS, Merz CN. Gender differences in the link between depression and cardiovascular disease. *Psychosom Med.* 2005;67 Suppl 1:S15-8.

26. Lehto S, Koukkunen H, Hintikka J, Viinamaki H, Laakso M, Pyorala K. Depression after coronary heart disease events. *Scandinav Cardiovasc J*. 2000;34:580-3.
27. Freedland KE, Rich MW, Skala JA, Carney RM, Davila-Roman VG, Jaffe AS. Prevalence of depression in hospitalized patients with congestive heart failure. *Psychosom Med*. 2003;65:119-28.
28. Milcent C, Dormont B, Durand-Zaleski I, Steg PG. Gender differences in hospital mortality and use of percutaneous coronary intervention in acute myocardial infarction: microsimulation analysis of the 1999 nationwide French hospitals database. *Circulation*. 2007;115:833-9.
29. Poon S, Goodman SG, Yan RT, Bugiardini R, Bierman AS, Eagle KA, Johnston N, Huynh T, Grondin FR, Schenck-Gustafsson K, Yan AT. Bridging the gender gap: Insights from a contemporary analysis of sex-related differences in the treatment and outcomes of patients with acute coronary syndromes. *Am Heart J*. 2012;163:66-73.
30. Gulati M, Shaw LJ, Bairey Merz CN. Myocardial ischemia in women: lessons from the NHLBI WISE study. *Clin Cardiol*. 2012;35:141-8.
31. Shaw LJ, Bugiardini R, Merz CN. Women and ischemic heart disease: evolving knowledge. *J Am Coll Cardiol*. 2009;54:1561-75.
32. Kendler KS, Gardner CO, Fiske A, Gatz M. Major depression and coronary artery disease in the Swedish twin registry: phenotypic, genetic, and environmental sources of comorbidity. *Arch Gen Psychiat*. 2009;66:857-63.

33. Carney RM, Freedland KE, Miller GE, Jaffe AS. Depression as a risk factor for cardiac mortality and morbidity: a review of potential mechanisms. *J Psychosom Res.* 2002;53:897-902.
34. Ormel J, de Jonge P. Unipolar depression and the progression of coronary artery disease: toward an integrative model. *Psychother Psychosom.* 2011;80:264-74.
35. Ziegelstein RC, Fauerbach JA, Stevens SS, Romanelli J, Richter DP, Bush DE. Patients with depression are less likely to follow recommendations to reduce cardiac risk during recovery from a myocardial infarction. *Arch Intern Med.* 2000;160:1818-23.
36. Doyle F, Rohde D, Kowalczyk A, Morgan K, Cousins G, McGee H. Systematic review and meta-analysis of the impact of depression on subsequent smoking cessation in patients with coronary heart disease: 1990–2013. *Psychosom Med.* 2014;76:44-57.
37. Goldstein JM, Cherkerzian S, Buka SL, Fitzmaurice G, Hornig M, Gillman M, O'Toole S, Sloan RP. Sex-specific impact of maternal-fetal risk factors on depression and cardiovascular risk 40 years later. *J Develop Origin Health Disease.* 2011;2:353-64.
38. Clark CJ, Henderson KM, de Leon CF, Guo H, Lunos S, Evans DA, Everson-Rose SA. Latent constructs in psychosocial factors associated with cardiovascular disease: an examination by race and sex. *Front Psychiat.* 2012;3:5.

39. Christenfeld NJ, Sloan RP, Carroll D, Greenland S. Risk factors, confounding, and the illusion of statistical control. *Psychosom Med.* 2004;66:868-75.
40. Kraemer HC, Stice E, Kazdin A, Offord D, Kupfer D. How do risk factors work together? Mediators, moderators, and independent, overlapping, and proxy risk factors. *Am J Psychiat.* 2001;158:848-56.
41. Stewart LA, Tierney JF. To IPD or not to IPD? Advantages and disadvantages of systematic reviews using individual patient data. *Eval Health Prof.* 2002;25:76-97.
42. Nagueh SF, Appleton CP, Gillebert TC, Marino PN, Oh JK, Smiseth OA, Waggoner AD, Flachskampf FA, Pellikka PA, Evangelisa A. Recommendations for the evaluation of left ventricular diastolic function by echocardiography. *Eur J Echocard.* 2009;10:165-93.
43. Schepis T, Gaemperli O, Koepfli P, Valenta I, Strobel K, Brunner A, Leschka S, Desbiolles L, Husmann L, Alkadhi H, Kaufmann PA. Comparison of 64-slice CT with gated SPECT for evaluation of left ventricular function. *J Nuclear Med.* 2006;47:1288-94.

Table 1: Baseline characteristics by sex

	Men (n=7200)	Women (n=2975)	p-value	N measured
Demographic characteristics				
Age (mean, (SD))	59.7 (11.5)	64.1 (12.2)	<0.001***	10,171
Employment status (% employed)	53.6	27.0	<0.001***	6,528
Partner status (% with partner)	77.5	51.4	<0.001***	6,412
Cardiac disease severity				
History of MI (% yes)	19.2	18.0	<0.001***	9,646
LVEF (% of patients <40%)	22.6	22.4	0.56	3,505
Killip class (% poor)	16.1	22.9	<0.001***	7,532
PTCA (%)	47.2	42.1	<0.001***	7,679
History of PTCA (%)	11.4	11.0	0.11	4,830
CABG (%)	9.8	9.3	<0.001***	8,139
History of CABG (%)	10.8	7.9	<0.001***	4,849
Thrombolysis (%)	33.8	34.0	0.061	8,065
Congestive HF (%)	17.2	24.3	0.013*	6,104
Other risk factors				
Diabetes (%)	18.0	29.8	<0.001***	10,060
Smoking (%)	49.5	32.8	<0.001***	9,942
BMI (mean, (SD))	26.8 (4.5)	27.8 (6.0)	0.002**	7,188
Hyperlipidemia / hypercholesterolemia (%)	45.3	50.2	<0.001***	8,405

Hypertension (%)	31.6	38.6	<0.001***	8,301
History of hypertension (%)	43.1	62.2	<0.001***	5,348
Medication use				
Hypolipidemics (%)	43.5	44.1	0.29	4,004
Beta-blockers (%)	72.3	71.2	0.003**	8,833
Aspirin (%)	87.2	87.2	0.49	7,561
Calcium-channel blockers / antagonists (%)	14.5	21.3	<0.001***	7,056
ACE-inhibitors (%)	48.7	49.8	0.26	8,550
Antidepressant use (%)	5.8	8.1	0.013*	5,507
Study type				
Diagnostic interview (n)	660	240		900
Questionnaire (n)	7108	2942		10,050

ACE: angiotensin converting-enzyme; BMI: body mass index; CABG: coronary artery bypass graft; HF: heart failure;

LVEF: left ventricular ejection fraction; MI: myocardial infarction; PTCA: percutaneous transluminal coronary angioplasty; SD: standard deviation

***p < 0.001; **p < 0.01; *p < 0.05

Table 2: Linear mixed model with random intercept for study, predicting depression z-score (n=3115, 5 studies)

	B	Bootstrapped 95% CI	p
Adjusted for age (n=3115, 5 studies)			
Age	-0.001	-0.004 – 0.002	0.57
LVEF	-0.015	-0.191 – 0.161	0.87
Sex	-0.371	-0.476 – -0.265	<0.001***
Sex*LVEF	0.294	0.090 – 0.498	0.005**
Adjusted for all confounders (n=2226, 3 studies)			
Age	-0.002	-0.008 – 0.003	0.36
History of MI	0.148	0.064 – 0.233	0.001**
Killip class	0.132	0.029 – 0.235	0.012*
Diabetes	0.145	0.059 – 0.231	0.001**
Smoking	0.057	0.024 – 0.090	0.001**
BMI	-0.011	-0.028 – 0.006	0.21
LVEF	-0.019	-0.089 – 0.052	0.61
Sex	-0.348	-0.426 – -0.269	<0.001***
Sex*LVEF	0.274	0.061 – 0.488	0.012*

***p < 0.001; **p < 0.01; *p < 0.05. Sex (0=female, 1=male)

Table 3: Multilevel Cox proportional hazards regression assessing sex differences in the association between depression and cardiovascular prognosis

	All-cause mortality (n=7628, 10 studies)			Cardiovascular events (CVE) (n=6556, 7 studies)		
	HR	Bootstrapped 95% CI	P	HR	Bootstrapped 95% CI	P
Age	1.07	1.06 – 1.08	< 0.001***	1.01	1.01 – 1.02	< 0.001***
Depression z-score	1.30	1.24 – 1.37	< 0.001***	1.17	1.13 – 1.22	< 0.001***
Sex	1.07	0.90 – 1.28	0.44	0.93	0.78 – 1.11	0.42
Sex * depression z-score	1.12	1.05 – 1.19	< 0.001***	1.07	0.98 – 1.17	0.118

***p < 0.001; **p < 0.01; *p < 0.05. Sex (-0.5=female, 0.5=male)

Table 4: Percentage change in sex*depression interaction effects in the prediction of all-cause mortality by additional disease indices or risk factors

Factor	HR for interaction sex*depression	Adjusted HR for interaction sex*depression	% change	N/n studies
History of MI	1.097***	1.088**	-8.9%	7543/10
LVEF	1.126*	1.086	-30.5%	3115/5
Killip class	1.137***	1.162***	+16.9%	5924/6
Diabetes	1.107***	1.130***	+20.2%	7587/10
Smoking	1.118**	1.119**	+0.8%	7485/10
BMI	1.128**	1.125**	-2.2%	6133/7
Adjusting for all covariates except LVEF	1.140***	1.152**	+8.0%	5172/5
Adjusting for all covariates	1.107**	1.071	-32.5%	2226/3

Adjusted for age, depression z-score, and sex, with study as a random factor

Note: The percentage change was estimated by using the log HR for the sex*depression interaction effect.

***p < 0.001; **p < 0.01; *p < 0.05

Figure Legends

Figure 1: Survival curves for all-cause mortality (ACM) and cardiovascular events (CVE), by depression category and sex (adjusted for age). depr = depression. Depression categories are based on a split-half of the continuous depression z-scores on self-report questionnaires. (Note: In all Cox regression analyses, the continuous depression z-score is used.)

Systematic review and individual patient data meta-analysis of sex differences in depression and prognosis in persons with myocardial infarction: a MINDMAPS study

Supplementary Table S1: Overview of participating studies

PI, study name, country	Start of study	N ³ / % Women (n women)	Inclusion criteria	Exclusion criteria	Mean age (SD)	Depression measurement instrument ¹ and cut-off score	N / % depressed	Mean depression score (SD)	Mean follow-up time (days)	No. of events ACM / CVE
R. Carney, J. A. Blumenthal ENRICHD and ancillary HRV study ² USA (1-3)	1996 / 1997	2848 / 42% (1196)	standardized criteria for MI; low social support or depression ancillary study: no depression or social isolation, otherwise eligible for ENRICHD	other life-threat. med. illn.; cogn. impairm.; too ill; other major psych. disord.; unable/refused to participate	61 (12.3)	BDI-1A ≥ 10	1951 / 69%	14.13 (8.87)	ACM: 834 CVE: 609	ACM: 350 CVE: 1151
J. Ormel, P. de Jonge, MIND-IT The Netherlands (4)	1999	1814 / 23% (417)	standardized criteria for MI age ≥ 18	other life-threat. med. illn.; MI during hosp. for other reason; psych. depr. treatm; part. in conflicting clin. trial	61 (11.6)	BDI 1-A ≥ 10	BDI: 474 / 26%	6.77 (6.18)	ACM: 2167 CVE: 1517	ACM: 278 CVE: 784
J. Ormel, P. de Jonge DepreMI The Netherlands (5)	1997	528 / 19% (100)	standardized criteria for MI	other life-threat. med. illn.; cogn. impairm.; too ill; unable to communicate; MI	61 (11.7)	modified Dutch version of CIDI 2.1 BDI 1A ≥ 10	CIDI: 73 / 15% BDI:	6.80 (6.12)	ACM: 2663 CVE: 1851	ACM: 116 CVE: 237

				during hosp. for other reason				117 / 23%		
J. Denollet The Netherlands (6)	2003	501 / 22% (110)	Standardized criteria for MI; Age > 30	other life-threat. med. illn.; cogn. Impairm.	60 (11.6)	BDI-1A \geq 10	132 / 26%	7.03 (6.18)	ACM: 1374 CVE: 1284	ACM: 38 CVE: 82
F. Doyle, H. M. McGee Ireland(7)	2003	433 / 25% (108)	standardized criteria for ACS (only MI patients were included in the IPD meta-analysis)	nr	63 (12.3)	HADS-D > 7 BDI-FS > 3	HADS- D or BDI- FS: 75 / 17%	HADS-D: 4,14 (3,26) BDI-FS: 1.88 (2.69)	ACM: 356 CVE: na	ACM: 17 CVE: na
F. Doyle, H.M. McGee Ireland(8)	2006	285 / 20% (57)	standardized criteria for ACS (only MI patients were included in the IPD meta-analysis)	patient too distressed	61 (10.9)	HADS-D > 7 BDI-FS (6- item) > 3	either HADS- D or BDI- FS: 78 / 27% HADS- D: 33 / 12%; BDI-FS 69 / 24%	HADS-D: 3,52 (3,07) BDI-FS: 2.15 (2.84)	ACM: 427 CVE: 427	ACM: 19 CVE: 67
S. Bergerone Italy(9)	1999	98 / 22% (22)	standardized criteria for MI	in-hosp. mort.; other major psych. disord.; alcoholism; antidepr. therapy	62 (12.6)	struct. int. based on DSM-IV criteria BDI 1A \geq 10	MDD: 14 / 14% BDI: 35 / 34%	BDI: 8.34 (8.48)	ACM: 1485 CVE: 1485	ACM: 6 CVE: 29

S.H. Hosseini	2004	540 / 31% (167)	standardized criteria for MI	other life-threat. med. illn.; MI result of CABG or angiogr.; too ill	58 (12.0)	BDI 1A ≥ 10	355 / 66%	14.20 (9.80)	na (24 months)	ACM: na CVE: 55
Iran(10)										
D.A. Lane	1997	288 / 25% (72)	standardized criteria for MI	other life-threat. med. illn.; cogn. impairm.; MI result of CABG or angiogr.; unable to communicate; too ill	63 (11.5)	BDI 1A ≥ 10	89 / 31%	7.72 (6.26)	ACM: 976 CVE:na	ACM: 38 CVE: na
UK(11, 12)										
L. Pilote	1996	553 / 19% (105)	acute MI; surv. up to 24 hours after hosp. adm.; adm. through emergency dept.	phys. unable to respond to quest.; unable to communicate	60 (12.2)	BDI 1 ≥ 10	193 / 35%	9.07 (7.93)	ACM: 350 CVE: 205	ACM: 32 CVE: 222
Canada(13)										
K. Parakh, R.C. Ziegelstein	1995	284 / 43% (122)	standardized criteria for MI	other life-threat. med. illn.; cogn. impairm.; too ill	65 (12.1)	SCID BDI 1A ≥ 10	MDD: 29 / 10% BDI: 56 / 20%	BDI: 5.76 (6.15)	ACM: 2428 CVE: na	ACM: 153 CVE: na
USA (14)										
C. Rafanelli	1995	61 / 15% (9)	standardized criteria for MI; first MI	nr	59 (9.9)	modified SCID (minor and major depression	7 / 11%	na	na (24 months)	ACM: 4 CVE: 22
Italy(15)										
H. Sato OACIS Japan(16)	1998	1042 / 20% (208)	standardized criteria for MI	in-hosp. mort.; other major psych. disord.; unable to communicate	63 (11.2)	ZSDS ≥ 40	438 / 42%	38.03 (9.07)	na (12 months)	ACM: na CVE: 283
R.P. Steeds	1999	131 / 67%	MI; age < 75	nr	60	BDI-II ≥ 14	52 /	12.42 (9.09)	ACM: 457	ACM: 11

UK(17)	(88)				(9.4)		40%		CVE: na	CVE: na
S.L. Grace	1997	468 / 28%	confirmed MI; age ≥ 18	too ill; unable to communicate	61 (12.2)	BDI 1A ≥ 10	136 / 28%	7.93 (7.22)	na (12 months)	ACM: 29 CVE: 101
Canada(18)		(131)								
C. Welin	1985	270 / 16%	standardized criteria for MI; first MI; age <	nr	56 (7.0)	ZSDS ≥ 40	96/ 36%	36.76 (8.58)	na (120 months)	ACM: 65 CVE: 73
Sweden(19)		(43)	65							

¹ Depression measurement instrument used in the current individual patient data meta-analysis.

² Part of the non-depressed control group came from an ancillary study and part of the patients in the non-depressed control group had low social support. Depressed patients were oversampled for the purpose of the study.

³ Number of patients included in IPD-meta-analysis.

Abbreviations: ACM: All-cause mortality; ACS: acute coronary syndrome; BDI: Beck Depression Inventory; BDI-FS: Beck Depression Inventory Fast Scale; CIDI: Composite International Diagnostic Interview; CVE: cardiovascular events; DepreMI: Depression after Myocardial Infarction; ENRICHD: Enhancing Recovery in Coronary Heart Disease; HADS-D: Hospital Anxiety and Depression Scale-Depression subscale; HRV: heart rate variability; MDD: major depression disorder; MI: myocardial infarction; MIND-IT: Myocardial Infarction and Depression Intervention Trial; na: not available; nr: not reported; OACIS: Osaka Acute Coronary Insufficiency Study; PI: primary investigator; SCID: Structured Clinical Interview for DSM; SD: standard deviation; ZSDS: Zung Self-rating Depression Scale

Note: Most data in Supplementary Table 1 has been reproduced with permission from the original MINDMAPS study published in the British Journal of Psychiatry (Meijer A, Conradi HJ, Bos EH, Anselmino M, Carney R, Denollet J, Doyle F, Freedland KE, Grace SL, Hosseini SH, Lane DA, Pilote L, Parakh K, Rafanelli C, Sato H, Steeds RP, Welin C, de Jonge P. Adjusted prognostic association of post-myocardial infarction depression with mortality and cardiovascular events: an

individual patient data meta-analysis. Br J Psychiat. 2013;203:90-102. <http://bjp.rcpsych.org/content/203/2/90.short>). The authors gratefully acknowledge this permission.

Supplementary Table S2: Interactions between sex and disease indices and other risk factors in the prediction of depression z-scores

	B	Bootstrapped 95% CI	p	N/n studies
Sex*History of MI	0.080	-0.051 – 0.212	0.23	7543/10
Sex*LVEF	0.294	0.090 – 0.498	0.005**	3115/5
Sex*Killip class	0.066	-0.210 – 0.078	0.37	5924/6
Sex*Diabetes	-0.045	-0.166 – 0.075	0.46	7587/10
Sex*Smoking	0.002	-0.101 – 0.105	0.97	7485/10
Sex*BDI	-0.008	-0.018 – 0.002	0.13	6133/7

Linear mixed models with random intercept, with z-depression scores as dependent variable, and sex, disease index/risk factor, and their interaction as predictors, adjusting for age. **p<0.01. Sex (0=female, 1=male)

Supplementary Table S3: Multilevel Cox proportional hazards regression assessing sex and age differences in the association between depression and cardiovascular prognosis

	All-cause mortality (n=7628, 10 studies)			Cardiovascular events (CVE) (n=6556, 7 studies)		
	HR	Bootstrapped 95% CI	P	HR	Bootstrapped 95% CI	P
Age	1.07	1.07 – 1.08	< 0.001***	1.01	1.01 – 1.02	< 0.001***
Depression z-score	1.32	1.21 – 1.43	< 0.001***	1.19	1.13 – 1.26	< 0.001***
Sex	0.89	0.77 – 1.04	0.44	0.53	0.23 – 1.22	0.14
Sex * depression z-score	1.14	1.05 – 1.24	< 0.001***	1.09	1.00 – 1.17	0.038*
Age * depression z-score	1.00	1.00 – 1.01	0.58	1.00	1.00 – 1.00	0.50
Sex * age	1.02	1.01 – 1.03	< 0.001***	1.01	1.00 – 1.02	0.17
Sex * age * depression z-score	1.00	0.99 – 1.01	0.92	1.00	0.99 – 1.00	0.59

***p < 0.001; **p < 0.01; *p < 0.05.

Note. Age and sex are grand-mean centered in this analysis, to facilitate interpretation of the main effects

References for supplementary Table S1

1. Carney RM, Freedland KE, Steinmeyer B, Blumenthal JA, de Jonge P, Davidson KW, Czajkowski SM, Jaffe AS. History of depression and survival after acute myocardial infarction. *Psychosom Med*. 2009;71:253-9.
2. Carney RM, Blumenthal JA, Stein PK, Watkins L, Catellier D, Berkman LF, Czajkowski SM, O'Connor C, Stone PH, Freedland KE. Depression, heart rate variability, and acute myocardial infarction. *Circulation*. 2001;104:2024-8.
3. The ENRICHD Investigators. Enhancing recovery in coronary heart disease patients (ENRICHD): study design and methods. *Am Heart J*. 2000;139:1-9.
4. van den Brink RH, van Melle JP, Honig A, Schene AH, Crijns HJ, Lambert FP, Ormel J. Treatment of depression after myocardial infarction and the effects on cardiac prognosis and quality of life: rationale and outline of the Myocardial INfarction and Depression-Intervention Trial (MIND-IT). *Am Heart J*. 2002;144:219-25.
5. de Jonge P, van den Brink RH, Spijkerman TA, Ormel J. Only incident depressive episodes after myocardial infarction are associated with new cardiovascular events. *J Am Coll Cardiol*. 2006;48:2204-8.
6. Denollet J, Martens EJ, Smith OR, Burg MM. Efficient assessment of depressive symptoms and their prognostic value in myocardial infarction patients. *J Affect Disord*. 2010;120:105-11.
7. Doyle F, McGee HM, De La Harpe D, Shelley E, Conroy R. The Hospital Anxiety and Depression Scale depression subscale, but not the Beck Depression

- Inventory-Fast Scale, identifies patients with acute coronary syndrome at elevated risk of 1-year mortality. *J Psychosom Res.* 2006;60:461-7.
8. Doyle F, Conroy R, McGee H, Delaney M. Depressive symptoms in persons with acute coronary syndrome: specific symptom scales and prognosis. *J Psychosom Res.* 2010;68:121-30.
 9. Drago S, Bergerone S, Anselmino M, Varalda PG, Cascio B, Palumbo L, Angelini G, Trevi PG. Depression in patients with acute myocardial infarction: influence on autonomic nervous system and prognostic role. Results of a five-year follow-up study. *Int J Cardiol.* 2007;115:46-51.
 10. Hosseini SH, Yousefnejad K, Tabiban S, Nesarhoseyni V, Bagheri B, Kiasari AM, Ghaemian A, Ghadirnejad SN, Lolati HA, Amiri FN, Ashraf H, Mokhberi V. Effects of depression and anxiety symptoms on cardiac mortality following myocardial infarction: a 2-year follow-up. *Int J Psychiat Clin Pract.* 2011;15:91-6.
 11. Lane D, Carroll D, Ring C, Beevers DG, Lip GY. Do depression and anxiety predict recurrent coronary events 12 months after myocardial infarction? *QJM.* 2000;93:739-44.
 12. Lane D, Carroll D, Ring C, Beevers DG, Lip GY. Mortality and quality of life 12 months after myocardial infarction: effects of depression and anxiety. *Psychosom Med.* 2001;63:221-30.
 13. Lauzon C, Beck CA, Huynh T, Dion D, Racine N, Carignan S, Diodati JG, Charbonneau F, Dupuis R, Pilote L. Depression and prognosis following hospital admission because of acute myocardial infarction. *CMAJ.* 2003;168:547-52.

14. Parakh K, Thombs BD, Fauerbach JA, Bush DE, Ziegelstein RC. Effect of depression on late (8 years) mortality after myocardial infarction. *Am J Cardiol.* 2008;101:602-6.
15. Rafanelli C, Milaneschi Y, Roncuzzi R, Pancaldi LG. Dysthymia before myocardial infarction as a cardiac risk factor at 2.5-year follow-up. *Psychosomatics.* 2010;51:8-13.
16. Shiotani I, Sato H, Kinjo K, Nakatani D, Mizuno H, Ohnishi Y, Hishida E, Kijima Y, Hori M, Sato H. Depressive symptoms predict 12-month prognosis in elderly patients with acute myocardial infarction. *J Cardiovasc Risk.* 2002;9:153-60.
17. Steeds RP, Bickerton D, Smith MJ, Muthusamy R. Assessment of depression following acute myocardial infarction using the Beck depression inventory. *Heart.* 2004;90:217-8.
18. Grace SL, Abbey SE, Pinto R, Shnek ZM, Irvine J, Stewart DE. Longitudinal course of depressive symptomatology after a cardiac event: effects of gender and cardiac rehabilitation. *Psychosom Med.* 2005;67:52-8.
19. Welin C, Lappas G, Wilhelmsen L. Independent importance of psychosocial factors for prognosis after myocardial infarction. *J Int Med.* 2000;247:629-39.