

Supplementary Table 2: Fatty acids present in feed components. Feed was sampled twice and an average was calculated (mg FAME/g dw).

Fatty acid	Content (mg FAME/g dw)
14:0	7.87±0.12
15:0	0.46±0.01
16:0	15.70±0.27
16:1n-9	0.21±0.00
16:1n-7	7.52±0.12
17:0	0.45±0.01
17:1n-7	0.00±0.00
18:0	3.19±0.05
18:1n-9t	0.09±0.01
18:1n-9c	22.76±0.31
18:2n-6t	0.09±0.01
18:2n-6c (LIN)	10.09±0.21
20:0	0.46±0.01
18:3n-6	0.14±0.00
20:1n-9	10.61±0.10
18:3n-3 (ALA)	3.66±0.07
21:0	0.02±0.05
18:4n-3	3.13±0.04
20:2n-6	0.41±0.01
22:0	0.90±0.01
20:3n-6	0.13±0.00
22:1n-9	1.05±0.01
20:3n-3	0.35±0.01
20:4n-6	0.46±0.01
23:0	0.10±0.01
20:4n-3	0.91±0.01
22:2n-6	None detected
24:0	0.00±0.00
20:5n-3 (EPA)	9.02±0.01
24:1n-9	0.78±0.00
22:3n-3	0.06±0.00
22:4n-6	0.06±0.01
22:5n-3	1.19±0.01
22:6n-3 (DHA)	9.41±0.02

Values are mean±S.E