

1 Contents

2	1 Cheilostome genera: range-through geologic durations	2
3	2 Text-mining references	11

1 Cheilostome genera: range-through geologic durations

Table S2: First and last appearances for cheilostome bryozoan genera. These are derived from the text-mined fossil occurrence data, and extant cheilostomes in the World Register of Marine Species. For each genus, the first appearance is the early boundary for the first time interval, and the last appearance is the late boundary of the last interval. Note that these first and last appearances have not been validated in any other way. Figs. 3 and S6 are based on the information given in this table. Ma = million years ago. $n = 337$.

Genus	First appearance (Ma)	Last appearance (Ma)
<i>Acoscinopectora</i>	83.60	72.10
<i>Actisecos</i>	17.46	0.00
<i>Adeonella</i>	8.70	0.00
<i>Aechmella</i>	100.50	66.00
<i>Aeolopora</i>	86.30	72.10
<i>Aetea</i>	41.30	0.00
<i>Akatopora</i>	86.30	0.00
<i>Alderina</i>	83.60	0.00
<i>Allantopora</i>	83.60	0.00
<i>Amphiblestrella</i>	72.10	68.03
<i>Amphiblestrum</i>	86.30	0.00
<i>Anaptopora</i>	100.50	93.90
<i>Anornithopora</i>	83.60	61.60
<i>Anoteropora</i>	20.44	0.00
<i>Anotopora</i>	100.50	93.90
<i>Antropora</i>	83.60	0.00
<i>Aplousina</i>	72.10	0.00
<i>Arachnopusia</i>	16.30	0.00
<i>Arctonula</i>	100.50	0.00
<i>Arthropoma</i>	23.03	0.00
<i>Aspidostoma</i>	83.60	0.00
<i>Bactrellaria</i>	83.60	72.10
<i>Bactridium</i>	16.30	12.80
<i>Balantiostoma</i>	72.10	66.00
<i>Basslerinella</i>	72.10	66.00
<i>Basyaylella</i>	11.62	7.25
<i>Bathosella</i>	72.10	0.00
<i>Bathystomella</i>	83.60	72.10
<i>Batopora</i>	56.00	0.00
<i>Batrachopora</i>	72.10	61.60
<i>Beisselina</i>	83.60	61.60
<i>Beisselinopsis</i>	72.10	61.60
<i>Bellulopora</i>	100.50	0.00
<i>Biflustra</i>	72.10	0.00
<i>Biselenaria</i>	41.30	37.80
<i>Bitectipora</i>	11.62	0.00

Table S2: First and last appearances for cheilostome bryozoan genera. These are derived from the text-mined fossil occurrence data, and extant cheilostomes in the World Register of Marine Species. For each genus, the first appearance is the early boundary for the first time interval, and the last appearance is the late boundary of the last interval. Note that these first and last appearances have not been validated in any other way. Figs. 3 and S6 are based on the information given in this table. Ma = million years ago. $n = 337$. (*continued*)

Genus	First appearance (Ma)	Last appearance (Ma)
<i>Boreas</i>	72.10	66.00
<i>Boreasina</i>	89.80	66.00
<i>Brydonella</i>	72.10	61.60
<i>Bryopesanser</i>	17.46	0.00
<i>Buffonellaria</i>	33.90	0.00
<i>Bullaconopeum</i>	75.93	66.00
<i>Caberea</i>	17.46	0.00
<i>Caberooides</i>	47.80	15.97
<i>Calloporina</i>	13.82	0.00
<i>Calpidopora</i>	100.50	93.90
<i>Calvina</i>	66.00	61.60
<i>Calypthotheca</i>	17.46	0.00
<i>Canda</i>	17.46	0.00
<i>Castanopora</i>	86.30	66.00
<i>Catencella</i>	47.80	0.00
<i>Cauloramphus</i>	11.62	0.00
<i>Cellaria</i>	166.10	0.00
<i>Cellarinidra</i>	100.50	93.90
<i>Celleporaria</i>	66.00	0.00
<i>Celleporella</i>	72.10	0.00
<i>Celleporina</i>	16.30	0.00
<i>Chaperia</i>	12.70	0.00
<i>Charixa</i>	132.90	0.00
<i>Cheethamia</i>	83.60	66.00
<i>Cheilopora</i>	38.00	0.00
<i>Cheiloporina</i>	38.00	0.00
<i>Chiastosella</i>	34.30	0.00
<i>Chiplonkarina</i>	113.00	66.00
<i>Cigclisula</i>	20.44	0.00
<i>Cleidochasmidra</i>	0.78	0.00
<i>Coleopora</i>	11.62	0.00
<i>Conopeum</i>	125.00	0.00
<i>Copidozoum</i>	20.44	0.00
<i>Corymboporella</i>	100.50	93.90
<i>Cosciniopsis</i>	47.80	0.00
<i>Coscinopleura</i>	66.00	61.60
<i>Costatimorpha</i>	37.80	33.90
<i>Cranosina</i>	13.82	0.00
<i>Crassimarginatella</i>	93.90	0.00

Table S2: First and last appearances for cheilostome bryozoan genera. These are derived from the text-mined fossil occurrence data, and extant cheilostomes in the World Register of Marine Species. For each genus, the first appearance is the early boundary for the first time interval, and the last appearance is the late boundary of the last interval. Note that these first and last appearances have not been validated in any other way. Figs. 3 and S6 are based on the information given in this table. Ma = million years ago. $n = 337$. (*continued*)

Genus	First appearance (Ma)	Last appearance (Ma)
<i>Craticulacella</i>	83.60	75.93
<i>Cribrilaria</i>	56.00	0.00
<i>Cribrilina</i>	93.90	0.00
<i>Criserpia</i>	168.30	166.10
<i>Crisidmonea</i>	83.60	0.00
<i>Cryptostomella</i>	83.60	66.00
<i>Ctenopora</i>	100.50	93.90
<i>Cupuladria</i>	16.30	0.00
<i>Cyclicopora</i>	38.00	0.00
<i>Cymulopora</i>	75.93	0.00
<i>Dacryoporella</i>	89.80	86.30
<i>Decurtaria</i>	66.00	61.60
<i>Diacanthopora</i>	66.00	61.60
<i>Dicasignetella</i>	41.30	37.80
<i>Dightonia</i>	21.70	19.00
<i>Dimorphomicropora</i>	72.10	66.00
<i>Dionella</i>	72.10	66.00
<i>Discoflustraria</i>	47.80	41.30
<i>Discopora</i>	72.10	66.00
<i>Discoporella</i>	23.03	0.00
<i>Discoradius</i>	47.80	33.90
<i>Discovibracella</i>	66.00	61.60
<i>Distelopora</i>	100.50	75.93
<i>Ditaxiporina</i>	47.80	28.10
<i>Dittosaria</i>	56.00	28.10
<i>Dysnoetocella</i>	98.30	96.10
<i>Ehrhardina</i>	100.50	93.90
<i>Ellisina</i>	72.10	0.00
<i>Emballotheca</i>	8.70	0.00
<i>Encicellaria</i>	72.10	15.97
<i>Escharella</i>	72.10	0.00
<i>Escharifora</i>	75.93	66.00
<i>Escharina</i>	100.50	0.00
<i>Escharipora</i>	89.80	86.30
<i>Eucratea</i>	56.00	0.00
<i>Euritina</i>	100.50	89.80
<i>Exechonella</i>	47.80	0.00
<i>Erochella</i>	83.60	0.00

Table S2: First and last appearances for cheilostome bryozoan genera. These are derived from the text-mined fossil occurrence data, and extant cheilostomes in the World Register of Marine Species. For each genus, the first appearance is the early boundary for the first time interval, and the last appearance is the late boundary of the last interval. Note that these first and last appearances have not been validated in any other way. Figs. 3 and S6 are based on the information given in this table. Ma = million years ago. $n = 337$. (*continued*)

Genus	First appearance (Ma)	Last appearance (Ma)
<i>Figularia</i>	66.00	0.00
<i>Filaguria</i>	13.82	0.00
<i>Floridina</i>	72.10	0.00
<i>Floridinella</i>	72.10	0.00
<i>Flustrellaria</i>	100.50	0.00
<i>Foratella</i>	72.10	66.00
<i>Foveolaria</i>	93.90	0.00
<i>Francopora</i>	89.80	86.30
<i>Frurionella</i>	72.10	61.60
<i>Fusicellaria</i>	93.90	89.80
<i>Gaudryanella</i>	47.80	41.30
<i>Gemellipora</i>	16.30	0.00
<i>Gemelliporella</i>	16.30	0.00
<i>Gephyrotes</i>	15.97	0.00
<i>Gigantopora</i>	13.82	0.00
<i>Gilbertopora</i>	113.00	75.93
<i>Gontarella</i>	13.82	0.00
<i>Graptoporella</i>	86.30	83.60
<i>Hagenowinella</i>	72.10	66.00
<i>Haplocephalopora</i>	66.00	61.60
<i>Haplostoechios</i>	104.67	93.90
<i>Hapsidopora</i>	100.50	93.90
<i>Hemiphylactella</i>	20.44	15.97
<i>Hemismittoidea</i>	37.80	0.00
<i>Herentia</i>	0.26	0.00
<i>Herpetopora</i>	125.00	61.60
<i>Heteractis</i>	47.80	28.10
<i>Heteroconopeum</i>	100.50	61.60
<i>Hiantopora</i>	66.00	0.00
<i>Hippaliosina</i>	0.78	0.00
<i>Hippodiplosia</i>	20.44	2.59
<i>Hippomenella</i>	38.00	0.00
<i>Hippopleurifera</i>	16.30	0.00
<i>Hippopodina</i>	17.46	0.00
<i>Hippoporella</i>	16.30	0.00
<i>Hippoporidra</i>	37.80	0.00
<i>Hippoporina</i>	28.10	0.00
<i>Hippothoa</i>	100.50	0.00

Table S2: First and last appearances for cheilostome bryozoan genera. These are derived from the text-mined fossil occurrence data, and extant cheilostomes in the World Register of Marine Species. For each genus, the first appearance is the early boundary for the first time interval, and the last appearance is the late boundary of the last interval. Note that these first and last appearances have not been validated in any other way. Figs. 3 and S6 are based on the information given in this table. Ma = million years ago. $n = 337$. (*continued*)

Genus	First appearance (Ma)	Last appearance (Ma)
<i>Hoeverella</i>	83.60	66.00
<i>Homalostega</i>	83.60	61.60
<i>Hoplitaechmella</i>	83.60	75.93
<i>Hoplocheilina</i>	100.50	61.60
<i>Houzeauina</i>	37.80	20.44
<i>Hybopora</i>	100.50	93.90
<i>Ichnopora</i>	93.90	61.60
<i>Inversaria</i>	75.93	61.60
<i>Iodictyum</i>	21.70	0.00
<i>Kionidella</i>	47.80	41.30
<i>Kleidionella</i>	72.10	61.60
<i>Kristerina</i>	83.60	72.10
<i>Kunradocella</i>	72.10	66.00
<i>Kylonisa</i>	47.80	43.47
<i>Labioporella</i>	20.44	0.00
<i>Lacrimula</i>	37.80	0.00
<i>Lagenipora</i>	37.80	0.00
<i>Lagynopora</i>	72.10	68.03
<i>Laterotecatia</i>	68.03	66.00
<i>Leieschara</i>	11.62	0.00
<i>Leptocheilopora</i>	86.30	66.00
<i>Lifuella</i>	17.46	11.62
<i>Lunulites</i>	93.90	33.90
<i>Macropora</i>	59.20	0.00
<i>Mamillopora</i>	37.80	0.00
<i>Margaretta</i>	37.80	0.00
<i>Marginaria</i>	104.67	20.44
<i>Melychocella</i>	59.20	47.80
<i>Membranipora</i>	137.50	0.00
<i>Membraniporidra</i>	66.00	0.00
<i>Metracolposa</i>	35.27	33.90
<i>Metrarabdotos</i>	20.44	0.00
<i>Metroperiella</i>	56.00	0.00
<i>Micropora</i>	132.90	0.00
<i>Microporella</i>	17.46	0.00
<i>Mollia</i>	100.50	0.00
<i>Monoceratopora</i>	93.90	61.60
<i>Monoporella</i>	83.60	0.00
<i>Multescharellina</i>	72.10	0.00

Table S2: First and last appearances for cheilostome bryozoan genera. These are derived from the text-mined fossil occurrence data, and extant cheilostomes in the World Register of Marine Species. For each genus, the first appearance is the early boundary for the first time interval, and the last appearance is the late boundary of the last interval. Note that these first and last appearances have not been validated in any other way. Figs. 3 and S6 are based on the information given in this table. Ma = million years ago. $n = 337$. (*continued*)

Genus	First appearance (Ma)	Last appearance (Ma)
<i>Multescharipora</i>	83.60	72.10
<i>Myriapora</i>	11.62	0.00
<i>Mystriopora</i>	113.00	100.50
<i>Nannopora</i>	89.80	86.30
<i>Nellia</i>	72.10	0.00
<i>Ogiva</i>	86.30	83.60
<i>Ogivalina</i>	36.63	35.27
<i>Onychocellaria</i>	72.10	66.00
<i>Opisthornithopora</i>	72.10	66.00
<i>Orbiculipora</i>	17.46	15.97
<i>Ornatella</i>	100.50	93.90
<i>Ostrovskia</i>	11.62	7.25
<i>Otionellina</i>	0.34	0.00
<i>Otopora</i>	100.50	93.90
<i>Oviexechonella</i>	13.82	11.62
<i>Pachyderma</i>	72.10	61.60
<i>Pachydermopora</i>	83.60	72.10
<i>Pachythecella</i>	68.03	61.60
<i>Parantropora</i>	13.82	0.00
<i>Parasmittina</i>	47.80	0.00
<i>Parellisina</i>	13.82	0.00
<i>Parkermavella</i>	1.63	0.00
<i>Pavobeisselina</i>	66.00	61.60
<i>Pavolunulites</i>	89.80	66.00
<i>Peedeesella</i>	72.10	66.00
<i>Pelmatopora</i>	93.90	61.60
<i>Pentapora</i>	0.78	0.00
<i>Petalostegus</i>	100.50	0.00
<i>Petralia</i>	23.03	0.00
<i>Phonicosia</i>	34.30	0.00
<i>Phylactella</i>	33.90	0.00
<i>Planicellaria</i>	66.00	15.97
<i>Platyglena</i>	72.10	66.00
<i>Plesiocleidochasma</i>	17.46	0.00
<i>Pleurocodonellina</i>	20.44	0.00
<i>Pleuromucrum</i>	20.44	0.00
<i>Pliophloea</i>	83.60	0.00
<i>Porella</i>	11.62	0.00

Table S2: First and last appearances for cheilostome bryozoan genera. These are derived from the text-mined fossil occurrence data, and extant cheilostomes in the World Register of Marine Species. For each genus, the first appearance is the early boundary for the first time interval, and the last appearance is the late boundary of the last interval. Note that these first and last appearances have not been validated in any other way. Figs. 3 and S6 are based on the information given in this table. Ma = million years ago. $n = 337$. (*continued*)

Genus	First appearance (Ma)	Last appearance (Ma)
<i>Poricella</i>	13.82	0.00
<i>Poricellaria</i>	72.10	0.00
<i>Porina</i>	100.50	0.00
<i>Predanophora</i>	17.46	11.62
<i>Prenantia</i>	5.33	0.00
<i>Pseudoallantopora</i>	72.10	66.00
<i>Pseudothyracella</i>	72.10	0.00
<i>Psilosecos</i>	63.07	61.60
<i>Puellina</i>	17.46	0.00
<i>Puncturiella</i>	72.10	66.00
<i>Pyripora</i>	113.00	0.00
<i>Pyriporella</i>	66.00	61.60
<i>Pyriporopsis</i>	163.50	132.90
<i>Pyrisinella</i>	72.10	66.00
<i>Quadricellaria</i>	93.90	0.00
<i>Quasitrilaminopora</i>	66.00	61.60
<i>Ramicosticella</i>	66.00	61.60
<i>Ramphonotus</i>	83.60	0.00
<i>Rectonychocella</i>	38.00	0.00
<i>Reniporella</i>	47.80	41.30
<i>Reptadeonella</i>	17.46	0.00
<i>Reptescharella</i>	89.80	86.30
<i>Reptoflustrella</i>	100.50	93.90
<i>Reptolunulites</i>	100.50	66.00
<i>Reptoporina</i>	20.44	15.97
<i>Retelepralia</i>	20.44	0.00
<i>Reussirella</i>	13.82	0.00
<i>Rhabdopora</i>	93.90	89.80
<i>Rhammatopora</i>	113.00	100.50
<i>Rhamphostomella</i>	56.00	0.00
<i>Rhiniopora</i>	83.60	66.00
<i>Rhynchozoon</i>	72.10	0.00
<i>Rosseliana</i>	34.30	0.00
<i>Saevitella</i>	17.46	0.00
<i>Sandalopora</i>	93.90	61.60
<i>Schizomavella</i>	20.44	0.00
<i>Schizoporella</i>	83.60	0.00
<i>Schizorthosecos</i>	35.27	33.90

Table S2: First and last appearances for cheilostome bryozoan genera. These are derived from the text-mined fossil occurrence data, and extant cheilostomes in the World Register of Marine Species. For each genus, the first appearance is the early boundary for the first time interval, and the last appearance is the late boundary of the last interval. Note that these first and last appearances have not been validated in any other way. Figs. 3 and S6 are based on the information given in this table. Ma = million years ago. $n = 337$. (*continued*)

Genus	First appearance (Ma)	Last appearance (Ma)
<i>Schizosmittina</i>	37.80	0.00
<i>Schizotheca</i>	11.62	0.00
<i>Scorpiodinipora</i>	23.03	0.00
<i>Scrupocellaria</i>	38.00	0.00
<i>Selenaria</i>	1.63	0.00
<i>Semieschara</i>	100.50	41.30
<i>Semiescharipora</i>	86.30	15.90
<i>Semihaskellia</i>	38.00	0.00
<i>Sendinopora</i>	13.82	11.62
<i>Sertulipora</i>	0.26	0.00
<i>Setosellina</i>	56.00	0.00
<i>Setosinella</i>	72.10	13.82
<i>Smittina</i>	17.46	0.00
<i>Smittipora</i>	38.00	0.00
<i>Smittoidea</i>	34.30	0.00
<i>Solenonychocella</i>	72.10	66.00
<i>Solenophragma</i>	83.60	66.00
<i>Spinicharixa</i>	125.00	100.50
<i>Spinisinella</i>	100.50	13.82
<i>Stamenocella</i>	93.90	89.80
<i>Staurosteginopora</i>	86.30	66.00
<i>Steginopora</i>	72.10	66.00
<i>Steginoporella</i>	47.80	0.00
<i>Stenopsella</i>	38.00	0.00
<i>Stenosipora</i>	47.80	41.30
<i>Stephanollona</i>	23.03	0.00
<i>Stephanotheca</i>	0.26	0.00
<i>Stichocados</i>	72.10	66.00
<i>Stichomicropora</i>	100.50	61.60
<i>Stichopora</i>	72.10	66.00
<i>Stichoporina</i>	47.80	41.30
<i>Stictostega</i>	83.60	72.10
<i>Stylopoma</i>	13.82	0.00
<i>Systemostoma</i>	72.10	68.03
<i>Taeniocellaria</i>	72.10	66.00
<i>Taenioporina</i>	72.10	66.00
<i>Tamanicella</i>	11.62	7.25
<i>Tecatia</i>	72.10	0.00
<i>Tetraplaria</i>	38.00	0.00

Table S2: First and last appearances for cheilostome bryozoan genera. These are derived from the text-mined fossil occurrence data, and extant cheilostomes in the World Register of Marine Species. For each genus, the first appearance is the early boundary for the first time interval, and the last appearance is the late boundary of the last interval. Note that these first and last appearances have not been validated in any other way. Figs. 3 and S6 are based on the information given in this table. Ma = million years ago. $n = 337$. (*continued*)

Genus	First appearance (Ma)	Last appearance (Ma)
<i>Thalamoporella</i>	47.80	0.00
<i>Therenia</i>	47.80	0.00
<i>Thoracopora</i>	100.50	93.90
<i>Thornelya</i>	13.82	0.00
<i>Tornipora</i>	72.10	61.60
<i>Trematooecia</i>	17.46	0.00
<i>Tremogasterina</i>	11.62	5.33
<i>Tricellaria</i>	0.26	0.00
<i>Tricephalopora</i>	89.80	61.60
<i>Trichinopolia</i>	83.60	66.00
<i>Triphyllozoon</i>	17.46	0.00
<i>Tripurula</i>	16.30	0.00
<i>Trypostega</i>	17.46	0.00
<i>Tubiporella</i>	17.46	0.00
<i>Turbicellepora</i>	33.90	0.00
<i>Turnerellina</i>	72.10	66.00
<i>Tyloporella</i>	100.50	93.90
<i>Ubagsia</i>	72.10	66.00
<i>Umbonula</i>	47.80	0.00
<i>Unidistelopora</i>	100.50	72.10
<i>Urceolipora</i>	21.70	0.00
<i>Vasignyella</i>	7.25	0.00
<i>Vibracella</i>	83.60	33.90
<i>Vibracellina</i>	56.00	0.00
<i>Vincularia</i>	86.30	5.33
<i>Vix</i>	20.44	0.00
<i>Voorthuyseniella</i>	23.03	20.44
<i>Wawalia</i>	139.80	113.00
<i>Wilbertopora</i>	113.00	66.00
<i>Woodipora</i>	72.10	66.00
<i>Xaveropora</i>	11.62	7.25
<i>Zimmerella</i>	83.60	72.10

2 Text-mining references

- Abdelsalam, K. M. 2014. Benthic bryozoan fauna from the northern Egyptian coast. *Egyptian Journal of Aquatic Research* 40:269–282.
- Adamczyk, A. D., and J. F. Pachut. 2013. Cladistic analysis of the Paleozoic bryozoan families Monticuliporidae and Mesotrypidae. *Journal of Paleontology* 87:635–649.
- Aguirre, J., J. C. Braga, J. M. Martín, and C. Betzler. 2012. Palaeoenvironmental and stratigraphic significance of Pliocene rhodolith beds and coralline algal bioconstructions from the Carboneras Basin (SE Spain). *Geodiversitas* 34:115–136.
- Almeida, A. C. S., F. B. C. Souza, C. M. S. Menegola, J. Sanner, and L. M. Vieira. 2014. Taxonomic review of the family Colatoeciidae Winston, 2005 (Bryozoa, Cheilostomata), with description of seven new species. *Zootaxa* 3868:1–61.
- Almeida, A. C. S., F. B. C. Souza, J. Sanner, and L. M. Vieira. 2015. Taxonomy of recent Adeonidae (Bryozoa, Cheilostomata) from Brazil, with the description of four new species. *Zootaxa* 4013:348–368.
- Altman, S., and R. B. Whitlatch. 2007. Effects of small-scale disturbance on invasion success in marine communities. *Journal of Experimental Marine Biology and Ecology* 342:15–29.
- Amui, A.-M. 2005. Adeonellas from the Gulf of Aden including one new species. *Zootaxa* 1012:45–52.
- Amui-Vedel, A.-M., P. J. Hayward, and J. S. Porter. 2007. Zooid size and growth rate of the bryozoan *Cryptosula pallasiana* Moll in relation to temperature, in culture and in its natural environment. *Journal of Experimental Marine Biology and Ecology* 353:1–12.
- Anstey, R. L., and J. F. Pachut. 2004. Cladistic and phenetic recognition of species in the Ordovician bryozoan genus *Peronopora*. *Journal of Paleontology* 78:651–674.
- Ariunchimeg, Y. 2012. *Admirandopora* nom. nov., a New Replacement Generic Name for a Carboniferous Bryozoan. *Paleontological Journal* 46:330–330.
- Arnold, P. W., and P. L. Cook. 1997. Some Recent species of the genus *Anaskopora* Wass, 1975 (Bryozoa: Cribriomorpha) from Queensland. *Memoirs of the Queensland Museum* 42:1–11.
- Álvarez, J. A. 1995. New data on the family Lichenoporidae Smitt (Bryozoa: Cyclostomida) from the Mediterranean region. *Journal of Natural History* 29:1067–1079.
- Bader, B., and P. Schäfer. 2004. Skeletal morphogenesis and growth check lines in the Antarctic bryozoan *Melicerita obliqua*. *Journal of Natural History* 38:2901–2922.
- Badve, R. M., and M. A. Sonar. 1995. Bryozoa Cheilostomata from Holocene, West Coast of Maharashtra, India. *Geobios* 28:317–335.
- Banta, W. C. 1971. The body wall of cheilostome Bryozoa. IV. The frontal wall of *Schizoporella unicornis* (Johnston). *Journal of Morphology* 135:165–184.
- Banta, W. C., N. Gray, and D. P. Gordon. 1997. A cryptocystal operculum and a new method of lophophore protrusion in the cheilostome bryozoan *Macropora levinseni*. *Invertebrate Biology* 116:161–170.
- Banta, W. C., F. K. McKinney, and R. L. Zimmer. 1974. Bryozoan monticules: excurrent water outlets? *Science* 185:783–784.
- Banta, W. C., F. M. Perez, and S. Santagata. 1995. A setigerous collar in *Membranipora chesapeakeensis* n. sp. (Bryozoa): implications for the evolution of cheilostomes from ctenostomes. *Invertebrate Biology* 114:83–88.
- Barnes, D. K. A. 2002. Invasions by marine life on plastic debris. *Nature* 416:808–809.
- Barnes, D. K. A. 2006. Temporal–spatial stability of competition in marine boulder fields. *Marine Ecology Progress Series* 314:15–23.
- Baud, A., H. A. Nakrem, B. Beauchamp, T. W. Beatty, A. F. Embry, and C. M. Henderson. 2008. Lower Triassic bryozoan beds from Ellesmere Island, High Arctic, Canada. *Polar Research* 27:428–440.
- Benton, M. J., and P. N. Pearson. 2001. Speciation in the fossil record. *Trends in Ecology & Evolution* 16:405–411.
- Bernecker, M., and O. Weidlich. 2005. Azooxanthellate corals in the Late Maastrichtian–Early Paleocene of

- 51 the Danish basin: bryozoan and coral mounds in a boreal shelf setting. Pages 3–25 in A. Freiwald and J. M.
52 Roberts, editors. Cold-water corals and ecosystems. Springer-Verlag, Berlin.
- 53 Berning, B. 2006. The cheilostome bryozoan fauna from the Late Miocene of Niebla (Guadalquivir Basin, SW
54 Spain): environmental and biogeographic implications. *Mitteilungen aus dem Geologisch-Paläontologischen*
55 *Institut der Universität Hamburg* 90:7–156.
- 56 Berning, B. 2007a. Evidence for sublethal predation and regeneration among living and fossil ascophoran bryozoans.
57 Pages 1–7 in S. J. Hageman, M. M. Key, and J. E. Winston, editors. *Bryozoan studies 2007*.
- 58 Berning, B. 2007b. The Mediterranean bryozoan *Myriapora truncata* (Pallas, 1766): a potential indicator of
59 (palaeo-) environmental conditions. *Lethaia* 40:221–232.
- 60 Berning, B. 2012. Taxonomic notes on some Cheilostomata (Bryozoa) from Madeira. *Zootaxa* 3236:36–54.
- 61 Berning, B. 2013. New and little-known Cheilostomata (Bryozoa, Gymnolaemata) from the NE Atlantic. *European*
62 *Journal of Taxonomy* 44:1–25.
- 63 Berning, B., and P. Kuklinski. 2008. North-east Atlantic and Mediterranean species of the genus *Buffonellaria*
64 (Bryozoa, Cheilostomata): implications for biodiversity and biogeography. *Zoological Journal of the Linnean*
65 *Society* 152:537–566.
- 66 Berning, B., M. Reuter, W. E. Piller, M. Harzhauser, and A. Kroh. 2009. Larger foraminifera as a substratum for
67 encrusting bryozoans (Late Oligocene, Tethyan Seaway, Iran). *Facies* 55:227–241.
- 68 Berning, B., K. J. Tilbrook, and A. N. Ostrovsky. 2014. What, if anything, is a lyrula? *Studi Trentini di Scienze*
69 *Naturali* 94:21–28.
- 70 Berning, B., K. J. Tilbrook, and A. Rosso. 2008. Revision of the north-eastern Atlantic and Mediterranean species
71 of the genera *Herentia* and *Therenia* (Bryozoa: Cheilostomata). *Journal of Natural History* 42:1509–1547.
- 72 Bishop, J. D. D. 1986. The identity of *Cribrilaria innominata* (Couch, 1844) (Bryozoa, Cheilostomata). *Bulletin*
73 *of the British Museum (Natural History), Zoology* 50:93–102.
- 74 Bishop, J. D. D. 1987. Hermit crabs associated with the bryozoan *Hippoporidra* in British waters. *Bulletin of the*
75 *British Museum (Natural History), Zoology* 52:29–30.
- 76 Bishop, J. D. D. 1994. The genera *Cribrilina* and *Collarina* (Bryozoa, Cheilostomatida) in the British Isles and
77 North Sea Basin, Pliocene to present day. *Zoologia Scripta* 23:225–249.
- 78 Bishop, J. D. D., and B. C. Househam. 1987. *Puellina* (Bryozoa; Cheilostomata; Cribriliniidae) from British and
79 adjacent waters. *Bulletin of the British Museum (Natural History), Zoology* 53:1–63.
- 80 Bjerager, M., and F. Surlyk. 2007. Benthic palaeoecology of Danian deep-shelf bryozoan mounds in the Danish
81 Basin. *Palaeogeography, Palaeoclimatology, Palaeoecology* 250:184–215.
- 82 Boardman, R. S. 1971. Mode of growth and functional morphology of autozooids in some Recent and Paleozoic
83 tubular Bryozoa. *Smithsonian Contributions to Paleobiology* 8:1–51.
- 84 Boardman, R. S. 1984. Origin of the post-Triassic Stenolaemata (Bryozoa): a taxonomic oversight. *Journal of*
85 *Paleontology* 58:19–39.
- 86 Boardman, R. S. 1998. Reflections on the morphology, anatomy, evolution, and classification of the class
87 Stenolaemata (Bryozoa). *Smithsonian Contributions to Paleobiology* 86:1–59.
- 88 Boardman, R. S. 1999. Indications of polypides in feeding zooids and polymorphs in lower Paleozoic Trepostomata
89 (Bryozoa). *Journal of Paleontology* 73:803–815.
- 90 Boardman, R. S. 2001. The growth and function of skeletal diaphragms in the colony life of lower Paleozoic
91 Trepostomata (Bryozoa). *Journal of Paleontology* 75:225–240.
- 92 Boardman, R. S., and C. J. Buttler. 2005. Zooids and extrazoooidal skeleton in the order Trepostomata (Bryozoa).
93 *Journal of Paleontology* 79:1088–1104.
- 94 Boardman, R. S., and A. H. Cheetham. 1969. Skeletal growth, intracolony variation, and evolution in Bryozoa: a
95 review. *Journal of Paleontology* 43:205–233.
- 96 Boardman, R. S., and F. K. McKinney. 1976. Skeletal architecture and preserved organs of four-sided zooids in
97 convergent genera of Paleozoic Trepostomata (Bryozoa). *Journal of Paleontology* 50:25–78.

- 98 Boardman, R. S., F. K. McKinney, and P. D. Taylor. 1992. Morphology, anatomy, and systematics of the
99 Cinctiporidae, new family (Bryozoa: Cheilostomata). *Smithsonian Contributions to Paleobiology* 70:1–81.
- 100 Bock, P. E., and P. L. Cook. 2001a. A review of Australian *Siphonicytara* Busk (Bryozoa: Cheilostomatida).
101 *Records of the Western Australian Museum* 20:307–322.
- 102 Bock, P. E., and P. L. Cook. 2001b. Revision of the multiphased genus *Corbulipora* MacGillivray (Bryozoa:
103 *Cribrimorpha*). *Memoirs of Museum Victoria* 58:191–213.
- 104 Bock, P. E., and P. L. Cook. 2002. First fossil finds of some Australian Bryozoa (Cheilostomata). *Alcheringa*
105 25:407–424.
- 106 Bock, P. E., and P. L. Cook. 2004a. Dimorphic brooding zooids in the genus *Adeona* Lamouroux from Australia
107 (Bryozoa: Cheilostomata). *Memoirs of Museum Victoria* 61:129–133.
- 108 Bock, P. E., and P. L. Cook. 2004b. A review of Australian Conescharellinidae (Bryozoa: Cheilostomata).
109 *Memoirs of Museum Victoria* 61:135–182.
- 110 Bone, Y., and N. P. James. 1993. Bryozoans as carbonate sediment producers on the cool-water Lacedpede Shelf,
111 southern Australia. *Sedimentary Geology* 86:247–271.
- 112 Bork, K. D., and T. G. Perry. 1968. Bryozoa (Ectoprocta) of Champlanian age (Middle Ordovician) from
113 northwestern Illinois and adjacent parts of Iowa and Wisconsin. Part III. *Homotrypa*, *Orbignyella*, *Prasopora*,
114 *Monticulipora*, and *Cyphotrypa*. *Journal of Paleontology* 42:1042–1065.
- 115 Braga, G. 2003. Salvador Reguant, Bryozoologist. *Geologica Acta* 1:169–171.
- 116 Braga, G., and F. Finotti. 2010. Aggiornamenti alla Bibliografia sul Phylum Bryozoa (fossili e viventi) dal 2006
117 al 2009 e integrazione dei lavori mancanti nella banca dati. *Annali del Museo Civico, Rovereto* 25:1–36.
- 118 Branch, M. L., and P. Hayward. 2005. New species of cheilostomatous Bryozoa from subantarctic Marion and
119 Prince Edward Islands. *Journal of Natural History* 39:2671–2704.
- 120 Brood, K. 1976a. Cyclostomatous Bryozoa from the coastal waters of East Africa. *Zoologica Scripta* 5:277–300.
- 121 Brood, K. 1976b. Note on the bryozoan *Pustulopora (Diaperoecia) intricaria* Busk, 1875. *Zoologia Scripta*
122 5:231–233.
- 123 Brood, K. 1976c. Cyclostomatous Bryozoa from the Paleocene and Maestrichtian of Majunga Basin, Madagascar.
124 *Géobios* 9:393–423.
- 125 Brown, D. A. 1956. Some Pliocene Polyzoa from South Australia. *Annals and Magazine of Natural History, series*
126 12 9:593–611.
- 127 Brown, G. D. 1965. Trepostomatous Bryozoa from the Logana and Jessamine limestones (Middle Ordovician) of
128 the Kentucky Bluegrass Region. *Journal of Paleontology* 39:974–1006.
- 129 Burton, E. S. B. 1929. The horizons of Bryozoa (Polyzoa) in the Upper Eocene beds of Hampshire. *Quarterly*
130 *Journal of the Geological Society of London* 85:223–241.
- 131 Busk, G. 1858. Zoophytology. On some Polyzoa collected by J.Y. Johnson, Esq., at Madeira. *Quarterly Journal*
132 *of Microscopical Science* 6:124–130.
- 133 Busk, G. 1859. A Monograph of the Fossil Polyzoa of the Crag. Palaeontographical Society, London.
- 134 Cadée, G. C., P. J. Chimonides, and P. L. Cook. 1989. *Pseudolunularia* gen.n. (Cheilostomata), a lunulitiform
135 bryozoan from the Indo-West Pacific. *Zoologica Scripta* 18:43–48.
- 136 Canu, F., and R. S. Bassler. 1919. Fossil Bryozoa from the West Indies. *Publications of the Carnegie Institution*
137 291:75–102.
- 138 Canu, F., and R. S. Bassler. 1923. North American later Tertiary and Quaternary Bryozoa. *United States*
139 *National Museum Bulletin* 125:1–302.
- 140 Canu, F., and R. S. Bassler. 1930. The bryozoan fauna of the Galapagos Islands. *Proceedings of the United*
141 *States National Museum* 76:1–78.
- 142 Canu, F., and G. Lecointre. 1927. Les Bryozoaires cheilostomes des Faluns de Touraine et d'Anjou. *Mémoires de*
143 *la Société Géologique de France, nouvelle série* 3:19–50.

- 144 Carter, M. C. 2008. The functional morphology of avicularia in cheilostome bryozoans. PhD thesis, Victoria
145 University, Wellington.
- 146 Carter, M. C., D. P. Gordon, and J. P. A. Gardner. 2010. Polymorphism and vestigiality: comparative anatomy
147 and morphology of bryozoan avicularia. *Zoomorphology* 129:195–211.
- 148 Cheetham, A. H. 1962. Eocene Bryozoa from the McBean Formation in Georgia. *Micropaleontology* 8:323–336.
- 149 Cheetham, A. H. 1966. Cheilostomatous Polyzoa from the Upper Bracklesham Beds (Eocene) of Sussex. *Bulletin*
150 *of the British Museum (Natural History)*, *Geology* 13:1–115.
- 151 Cheetham, A. H. 1971. Functional morphology and biofacies distribution of cheilostome Bryozoa in the Danian
152 Stage (Paleocene) of southern Scandinavia. *Smithsonian Contributions to Paleobiology* 6:1–87.
- 153 Cheetham, A. H. 1975. Preliminary report on Early Eocene cheilostome bryozoans from Site 308 - Leg 32,
154 Deep-Sea Drilling Project. *Initial Reports of the Deep Sea Drilling Project* 32:835–851.
- 155 Cheetham, A. H., L. C. Hayek, and E. Thomsen. 1981. Growth models in fossil arborescent cheilostome bryozoans.
156 *Paleobiology* 7:68–86.
- 157 Cheetham, A. H., J. B. C. Jackson, and J. Sanner. 2001. Evolutionary significance of sexual and asexual modes
158 of propagation in Neogene species of the bryozoan *Metrarabdotos* in tropical America. *Journal of Paleontology*
159 75:564–577.
- 160 Cheetham, A. H., and D. M. Lorenz. 1976. A vector approach to size and shape comparisons among zooids in
161 cheilostome bryozoans. *Smithsonian Contributions to Paleobiology* 29:1–55.
- 162 Cheetham, A. H., and P. A. Sandberg. 1964. Quaternary Bryozoa from Louisiana mudlumps. *Journal of*
163 *Paleontology* 38:1013–1046.
- 164 Cheetham, A. H., J. Sanner, and J. B. C. Jackson. 2007. *Metrarabdotos* and related genera (Bryozoa: Cheilosto-
165 mata) in the late Paleogene and Neogene of tropical America. *Journal of Paleontology* 81:1–91.
- 166 Cheetham, A. H., J. Sanner, P. D. Taylor, and A. N. Ostrovsky. 2006. Morphological differentiation of avicularia
167 and the proliferation of species in mid-Cretaceous *Wilbertopora* Cheetham, 1954 (Bryozoa: Cheilostomata).
168 *Journal of Paleontology* 80:49–71.
- 169 Cheetham, A. H., and E. Thomsen. 1981. Functional morphology of arborescent animals: strength and design of
170 cheilostome bryozoan skeletons. *Paleobiology* 7:355–383.
- 171 Chimonides, P. J. 1987. Notes on some species of the genus *Amathia* (Bryozoa, Ctenostomata). *Bulletin of the*
172 *British Museum (Natural History)*, *Zoology* 52:307–358.
- 173 Chimonides, P. J., and P. L. Cook. 1994. Notes on the genus *Cranosina* (Bryozoa, Cheilostomida). *Zoologia*
174 *Scripta* 23:43–49.
- 175 Clark, N., M. Williams, B. Okamura, J. Smellie, A. Nelson, T. Knowles, P. D. Taylor, M. J. Leng, J. Zalasiewicz,
176 and A. M. Haywood. 2010. Early Pliocene Weddell Sea seasonality determined from bryozoans. *Stratigraphy*
177 7:199–206.
- 178 Cleary, D., and P. N. Wyse Jackson. 2007. *Stenophragmidium* Bassler, 1952 (Trepotomida: Bryozoa) from the
179 Mississippian of Ireland and Britain. *Irish Journal of Earth Sciences* 25:1–25.
- 180 Clin, B., C. Stosse-Guevel, M.-F. Marquignon, L. Verneuil, and M. Letourneux. 2008. Professional photosensitive
181 eczema of fishermen by contact with bryozoans: disabling occupational dermatosis. *International Maritime*
182 *Health* 59:1–4.
- 183 Cocito, S., and F. Ferdeghini. 2000. Morphological variations in *Pentapora fascialis* (Cheilostomatida, Ascophorina).
184 Pages 176–181 in A. Herrera Cubilla and J. B. C. Jackson, editors. *Proceedings of the 11th international*
185 *bryozoology association conference, 1998*. Smithsonian Tropical Research Institute, Balboa, Republic of Panama.
- 186 Cook, P. L. 1964. Polyzoa from West Africa. I. Notes on the Steganoporellidae, Thalamoporellidae and
187 Onychocellidae (Anasca, Coilostega). *Annales de l'Institut Oceanographique (Calypso IV)* 41:43–79.
- 188 Cook, P. L. 1965a. Polyzoa from West Africa. The Cupuladriidae (Cheilostomata, Anasca). *Bulletin of the British*
189 *Museum (Natural History)*, *Zoology* 13:189–227.
- 190 Cook, P. L. 1965b. Notes on the Cupuladriidae (Polyzoa, Anasca). *Bulletin of the British Museum (Natural*

- 191 History), Zoology 13:151–187.
- 192 Cook, P. L. 1967. Polyzoa (Bryozoa) from West Africa. The Pseudostega, the Cribrimorpha and some Ascophora.
193 Bulletin of the British Museum (Natural History), Zoology 15:321–351.
- 194 Cook, P. L. 1982. Notes on some African Adeonellidae (Bryozoa, Cheilostomata). Journal of Natural History
195 16:833–846.
- 196 Cook, P. L., and P. E. Bock. 1994. The astogeny and morphology of *Rhabdozoum wilsoni* Hincks (Anasca,
197 Buguloidea). Pages 47–50 in P. J. Hayward, J. S. Ryland, and P. D. Taylor, editors. Biology and palaeobiology
198 of bryozoans. Olsen & Olsen, Fredensborg.
- 199 Cook, P. L., and P. E. Bock. 2002. Notes on astogeny of some Petraliellidae (Bryozoa) from Australia. Journal of
200 Natural History 36:1601–1619.
- 201 Cook, P. L., and P. J. Chimonides. 1981a. Morphology and systematics of some rooted cheilostome Bryozoa.
202 Journal of Natural History 15:97–134.
- 203 Cook, P. L., and P. J. Chimonides. 1981b. Morphology and systematics of some interior-walled cheilostome
204 Bryozoa. Bulletin of the British Museum (Natural History), Zoology 41:53–89.
- 205 Cook, P. L., and P. J. Chimonides. 1984a. Recent and fossil Lunulitidae (Bryozoa: Cheilostomata). 1. The genus
206 *Otionella* from New Zealand. Journal of Natural History 18:227–254.
- 207 Cook, P. L., and P. J. Chimonides. 1984b. Recent and fossil Lunulitidae (Bryozoa: Cheilostomata). 2. Species of
208 *Helixotionella* gen. nov. from Australia. Journal of Natural History 18:255–270.
- 209 Cook, P. L., and P. J. Chimonides. 1985a. Recent and fossil Lunulitidae (Bryozoa: Cheilostomata). 3. 'Opesiulate'
210 and other species of *Selenaria* sensu lato. Journal of Natural History 19:285–322.
- 211 Cook, P. L., and P. J. Chimonides. 1985b. Recent and fossil Lunulitidae (Bryozoa: Cheilostomata). 5. *Selenaria*
212 *alata* Tenison Woods, and related species. Journal of Natural History 19:337–358.
- 213 Cook, P. L., and P. J. Chimonides. 1986. Recent and fossil Lunulitidae (Bryozoa, Cheilostomata). 6. *Lunulites*
214 sensu lato and the genus *Lunularia* from Australasia. Journal of Natural History 20:681–705.
- 215 Cook, P. L., and P. J. Chimonides. 1987. Recent and fossil Lunulitidae (Bryozoa, Cheilostomata). 7. *Selenaria*
216 *maculata* (Busk) and allied species from Australasia. Journal of Natural History 21:933–966.
- 217 Cook, P. L., and P. J. Hayward. 1983. Notes on the family Lekythoporidae (Bryozoa, Cheilostomata). Bulletin of
218 the British Museum (Natural History), Zoology 45:55–76.
- 219 Cook, P. L., and R. Lagaaij. 1976. Some Tertiary and Recent conescharelliniform Bryozoa. Bulletin of the British
220 Museum (Natural History), Zoology 29:319–376.
- 221 Crowley, S. F., and P. D. Taylor. 2000. Stable isotope composition of modern bryozoan skeletal carbonate from
222 the Otago Shelf, New Zealand. New Zealand Journal of Marine and Freshwater Research 34:331–351.
- 223 Cuffey, R. J., R. M. Feldmann, and K. E. Pohlable. 1981. New Bryozoa from the Fox Hills Sandstone (Upper
224 Cretaceous, Maestrichtian) of North Dakota. Journal of Paleontology 55:401–409.
- 225 Deudero, S., A. Blanco, A. Box, G. Mateu-Vicens, M. Cabanellas-Reboredo, and A. Sureda. 2010. Interaction
226 between the invasive macroalga *Lophocladia lallemandii* and the bryozoan *Reteporella grimaldii* at seagrass
227 meadows: density and physiological responses. Biological Invasions 12:41–52.
- 228 Di Geronimo, I., C. Messina, A. Rosso, R. Sanfilippo, F. Sciuto, and A. Vertino. 2005. Enhanced biodiversity
229 in the deep: Early Pleistocene coral communities from southern Italy. Pages 61–86 in A. Freiwald and J. M.
230 Roberts, editors. Cold-water corals and ecosystems. Springer-Verlag, Berlin.
- 231 Di Martino, E., and A. Rosso. 2015. Revision of the bryozoan genus *Gephyrotes* Norman, 1903 (Cheilostomata,
232 Cribriliniidae) with the description of two new taxa. Zootaxa 3941:261–283.
- 233 Di Martino, E., and P. D. Taylor. 2012a. Pyrisinellidae, a new family of anascan cheilostome bryozoans. Zootaxa
234 3534:1–20.
- 235 Di Martino, E., and P. D. Taylor. 2012. Morphology and palaeobiogeography of *Retelepralia*, a distinctive
236 cheilostome bryozoan new to the fossil record. Neues Jahrbuch für Geologie und Paläontologie-Abhandlungen
237 263:67–74.

- 238 Di Martino, E., and P. D. Taylor. 2012b. Systematics and life history of *Antonietta* *exigua*, a new genus and
239 species of cribrimorph bryozoan from the Miocene of East Kalimantan (Indonesia). *Bollettino della Società*
240 *Paleontologica Italiana* 51:99–108.
- 241 Di Martino, E., and P. D. Taylor. 2014a. A brief review of seagrass-associated bryozoans, Recent and fossil. *Studi*
242 *Trentini di Scienze Naturali* 94:79–94.
- 243 Di Martino, E., and P. D. Taylor. 2014b. Miocene Bryozoa from East Kalimantan, Indonesia. Part I: Cyclostomata
244 and ‘Anascan’ Cheilostomata. *Scripta Geologica* 146:17–126.
- 245 Di Martino, E., and P. D. Taylor. 2015. Miocene Bryozoa from East Kalimantan, Indonesia. Part II: ‘Ascophoran’
246 Cheilostomata. *Scripta Geologica* 148:1–142.
- 247 Dick, M. H., A. V. Grischenko, and S. F. Mawatari. 2005. Intertidal Bryozoa (Cheilostomata) of Ketchikan,
248 Alaska. *Journal of Natural History* 39:3687–3784.
- 249 Dick, M. H., A. Herrera-Cubilla, and J. B. C. Jackson. 2003. Molecular phylogeny and phylogeography of
250 free-living Bryozoa (Cupuladriidae) from both sides of the Isthmus of Panama. *Molecular Phylogenetics and*
251 *Evolution* 27:355–371.
- 252 Dick, M. H., M. Hirose, R. Takashima, T. Ishimura, H. Nishi, and S. F. Mawatari. 2008. Application of MART
253 analysis to infer paleoseasonality in a Pleistocene shallow marine benthic environment. Pages 93–99 *in* H. Okada,
254 S. F. Mawatari, N. Suzuki, and P. Gautam, editors. *Origin and evolution of natural diversity*. Cambridge
255 University Press.
- 256 Dick, M. H., T. Komatsu, R. Takashima, and A. N. Ostrovsky. 2014. A mid-Cretaceous (Albian–Cenomanian)
257 shell-rubble bryozoan fauna from the Goshoura Group, Kyushu, Japan. *Journal of Systematic Palaeontology*
258 12:401–425.
- 259 Dick, M. H., S. Lidgard, D. P. Gordon, and S. F. Mawatari. 2009. The origin of ascophoran bryozoans was
260 historically contingent but likely. *Proceedings of the Royal Society of London, B* 276:3141–3148.
- 261 Dick, M. H., C. Sakamoto, and T. Komatsu. 2018. Cheilostome Bryozoa from the Upper Cretaceous Himenoura
262 Group, Kyushu, Japan. *Paleontological Research* 22:239–264.
- 263 Dominici, S. 2001. Taphonomy and paleoecology of shallow marine macrofossil assemblages in a collisional setting
264 (Late Pliocene–Early Pleistocene, Western Emilia, Italy). *Palaios* 16:336–353.
- 265 Dudley, J. E. 1973. Observations on the reproduction, early larval development, and colony astogeny of *Conopeum*
266 *tenuissimum* (Canu). *Chesapeake Science* 14:270–278.
- 267 Dumont, J. P. C. 1981. A report on the cheilostome Bryozoa of the Sudanese Red Sea. *Journal of Natural History*
268 15:623–637.
- 269 Duncan, H. 1939. Trepostomatous Bryozoa from the Traverse Group of Michigan. *Contributions from the Museum*
270 *of Paleontology, University of Michigan* 5:171–270.
- 271 Dzik, J. 1975. The origin and early phylogeny of the cheilostomatous Bryozoa. *Acta Palaeontologica Polonica*
272 20:395–423.
- 273 Dzik, J. 1992. Early astogeny and relationships of the Ordovician rhabdomesine bryozoans. *Acta Palaeontologica*
274 *Polonica* 37:37–54.
- 275 D’Hondt, J.-L. 2006. Nouvelle explications des planches de "Polypes" de la Description de l’Égypte. II - Bryozoaires.
276 *Collection Nouvelle Description de l’Égypte* 2:1–86.
- 277 D’Onghia, G., F. Capezzuto, F. Cardone, R. Carlucci, A. Carluccio, G. Chimienti, G. Corriero, C. Longo, P.
278 Maiorano, F. Mastrototaro, P. Panetta, A. Rosso, R. Sanfilippo, L. Sion, and A. Tursi. 2015. Macro- and
279 megafauna recorded in the submarine Bari Canyon (southern Adriatic, Mediterranean Sea) using different tools.
280 *Mediterranean Marine Science* 16:180–196.
- 281 Eliášová, H. 2005. Upper Cenomanian–Turonian two species of cyclostomaceous Bryozoa from the northern part
282 of the Bohemian Cretaceous basin, Czech Republic. *Journal of the Czech Geological Society* 50:59–62.
- 283 Ernst, A. 2000. Permian Bryozoans of the NW-Tethys. *Facies* 43:79–102.
- 284 Ernst, A. 2009. *Petaloporella* (Cryptostoma, Bryozoa) from the Lower Devonian of central Bohemia. *Bulletin of*
285 *Geosciences* 84:767–770.

- 286 Ernst, A. 2013. Diversity dynamics and evolutionary patterns of Devonian Bryozoa. *Palaeobiodiversity and*
287 *Palaeoenvironments* 93:45–63.
- 288 Ernst, A., and M. Carrera. 2008. Cryptostomid bryozoans from the Sassito Formation, Upper Ordovician
289 cool-water carbonates of the Argentinean Precordillera. *Palaeontology* 51:1117–1127.
- 290 Ernst, A., and P. Königshof. 2008. The role of bryozoans in fossil reefs—an example from the Middle Devonian of
291 the Western Sahara. *Facies* 54:613–620.
- 292 Ernst, A., and P. Königshof. 2010. Bryozoan fauna and microfacies from a Middle Devonian reef complex (Western
293 Sahara, Morocco). *Abhandlungen der Senckenberg Gesellschaft Naturforschung* 568:1–91.
- 294 Ernst, A., P. Königshof, P. D. Taylor, and J. Bohaty. 2011. Microhabitat complexity—an example from Middle
295 Devonian bryozoan-rich sediments in the Blankenheim Syncline (northern Eifel, Rheinisches Schiefergebirge).
296 *Palaeobiodiversity and Palaeoenvironments* 91:257–284.
- 297 Ernst, A., and A. May. 2012. Bryozoan fauna from the Lower Devonian (Middle Lochkovian) of Sierra de
298 Guadarrama, Spain. *Journal of Paleontology* 86:60–80.
- 299 Ernst, A., A. May, and S. Marks. 2012. Bryozoans, corals, and microfacies of Lower Eifelian (Middle Devonian)
300 limestones at Kierspe, Germany. *Facies* 58:727–758.
- 301 Ernst, A., and E. Minwegen. 2006. Late Carboniferous bryozoans from La Hermida, Spain. *Acta Palaeontologica*
302 *Polonica* 51:569–588.
- 303 Ernst, A., A. Munnecke, and I. Oswald. 2015a. Exceptional bryozoan assemblage of a microbial-dominated reef
304 from the early Wenlock of Gotland, Sweden. *GFF* 0:1–24.
- 305 Ernst, A., P. Schäfer, and J. A. Grant-Mackie. 2015b. New Caledonian Triassic Bryozoa. *Journal of Paleontology*
306 89:730–747.
- 307 Ernst, A., P. Schäfer, and J. J. G. Reijmer. 2005. Stenolaemate Bryozoa from the Upper Carboniferous of the
308 Cantabrian Basin, northern Spain. *Senckenbergiana lethaea* 85:301–317.
- 309 Ernst, A., B. Senowbari-Daryan, and K. Rashidi. 2009. Bryozoa from the Surmaq Formation (Permian) of the
310 Hambast Mountains, south of Abadeh, central Iran. *Facies* 55:595–608.
- 311 Ernst, A., O. Weidlich, and P. Schäfer. 2008. Stenolaemate Bryozoa from the Permian of Oman (Aseelah Unit,
312 Batain coast). *Journal of Paleontology* 82:676–716.
- 313 Ernst, A., and C. F. Winkler Prins. 2008. Pennsylvanian bryozoans from the Cantabrian Mountains (northwestern
314 Spain). *Scripta Geologica* 137:1–123.
- 315 Ferrario, J., J.-L. D’Hondt, A. Marchini, and A. Occhipinti-Ambrogi. 2015. From the Pacific Ocean to the
316 Mediterranean Sea: *Watersipora arcuata*, a new non-indigenous bryozoan in Europe. *Marine Biology Research*
317 11:909–919.
- 318 Fordinál, K., K. Zágorsek, and A. Zlinská. 2006. Early Sarmatian biota in the northern part of the Danube Basin
319 (Slovakia). *Geologica Carpathica* 57:123–130.
- 320 Frazier, J. G., J. E. Winston, and C. A. Ruckdeschel. 1992. Epizoan communities on marine turtles. III. Bryozoa.
321 *Bulletin of Marine Science* 51:1–8.
- 322 Fritz, M. A. 1944. Upper Devonian Bryozoa from New Mexico. *Journal of Paleontology* 18:31–41.
- 323 Fritz, M. A. 1946. Permian(?) Bryozoa from Sustut Lake, British Columbia. *Journal of Paleontology* 20:86–87.
- 324 Galle, A., and R. L. Parsley. 2005. Epibiont relationships on hyolithids demonstrated by Ordovician trepostomes
325 (Bryozoa) and Devonian tabulates (Anthozoa). *Bulletin of Geosciences* 80:125–138.
- 326 Gautier, T. G., P. N. Wyse Jackson, and F. K. McKinney. 2013. *Adlatipora*, a distinctive new Acanthocladiid
327 Bryozoan from the Permian of the Glass Mountains, Texas, U.S.A., and its bearing on fenestrate astogeny and
328 growth. *Journal of Paleontology* 87:444–455.
- 329 Gazdzicki, A., A. Tatur, U. Hara, and R. A. del Valle. 2004. The Weddell Sea Formation: post-Late Pliocene
330 terrestrial glacial deposits on Seymour Island, Antarctic Peninsula. *Polish Polar Research* 25:189–204.
- 331 Gili, J.-M., R. Coma, C. Orejas, P. J. López-González, and M. Zabala. 2001. Are Antarctic suspension-feeding
332 communities different from those elsewhere in the world? *Polar Biology* 24:473–485.

- 333 Gilmour, E. H. 2007. New Carboniferous Bryozoa of the Bird Spring Formation, southern Nevada. *Journal of*
334 *Paleontology* 81:581–587.
- 335 Gilmour, E. H., and M. E. McColloch. 1995. Fenestrada and Rhabdomesida (Bryozoa) of the Otter Formation
336 (Viséan), Central Montana. *Journal of Paleontology* 69:813–830.
- 337 Gilmour, E. H., M. E. McColloch, and B. R. Wardlaw. 1997. Bryozoa of the Murdock Mountain Formation
338 (Wordian, Permian), Leach Mountains, Northeastern Nevada. *Journal of Paleontology* 71:214–236.
- 339 Gilmour, E. H., and E. M. Snyder. 2000. Bryozoa of the Mission Argillite (Permian) Northeastern Washington.
340 *Journal of Paleontology* 74:545–570.
- 341 Gontar, V. I. 2009. New species from the Order Cyclostomata (Bryozoa) from the northern and middle Kurile
342 Islands. *Proceedings of the Zoological Institute RAS* 313:41–57.
- 343 Gordon, D. P. 1988. The bryozoan families Sclerodomidae, Bifaxariidae, and Urceoliporidae and a novel type of
344 frontal wall. *New Zealand Journal of Zoology* 15:249–290.
- 345 Gordon, D. P. 1989. The marine fauna of New Zealand: Bryozoa: Gymnolaemata (Cheilostomatida Ascophorina)
346 from the western South Island continental shelf and slope. *New Zealand Oceanographic Institute Memoir*
347 *97:1–158.*
- 348 Gordon, D. P. 1993a. Bryozoa: The ascophorine infraorders Cribriomorpha, Hippothoomorpha and Umbonulo-
349 morpha mainly from New Caledonian waters. *Mémoires du Muséum National d’Histoire Naturelle* 158:299–347.
- 350 Gordon, D. P. 1993b. Bryozoan frontal shields: studies on umbonulomorphs and impacts on classification.
351 *Zoologica Scripta* 22:203–221.
- 352 Gordon, D. P. 2000. First fossil occurrence of the austral bryozoan family Urceoliporidae. *New Zealand Journal*
353 *of Geology and Geophysics* 43:385–389.
- 354 Gordon, D. P. 2009a. New names for some bryozoan homonyms. *Zootaxa* 2133:64–68.
- 355 Gordon, D. P. 2009b. *Baudina* gen. nov., constituting the first record of Pasytheidae from Australia, and
356 Sinoflustridae fam. nov., with a checklist of Bryozoa and Pterobranchia from Beagle Gulf. *The Beagle, Records*
357 *of the Museums and Art Galleries of the Northern Territory* 25:43–54.
- 358 Gordon, D. P. 2009c. New bryozoan taxa from a new marine conservation area in New Zealand, with a checklist
359 of Bryozoa from Greater Cook Strait. *Zootaxa* 1987:39–60.
- 360 Gordon, D. P. 2014. Apprehending novel biodiversity—fifteen new genera of Zealandian Bryozoa. *Journal of the*
361 *Marine Biological Association of the United Kingdom* 94:1597–1628.
- 362 Gordon, D. P., and A. V. Grischenko. 1994. Bryozoan frontal shields: the types species of *Desmacystis*,
363 *Ramphostomella*, *Ramphosmittina*, *Ramphostomellina*, and new genus *Arctonula*. *Zoologica Scripta* 23:61–72.
- 364 Gordon, D. P., S. F. Mawatari, and H. Kajihara. 2002. New taxa of Japanese and New Zealand Eurystomellidae
365 (Phylum Bryozoa) and their phylogenetic relationships. *Zoological Journal of the Linnean Society* 136:199–216.
- 366 Gordon, D. P., and W. B. Rudman. 2006. *Integripelta acanthus* n. sp. (Bryozoa: Eurystomellidae) — a tropical
367 prey species of *Okenia hiroi* (Nudibranchia). *Zootaxa* 1229:41–48.
- 368 Gordon, D. P., and M. E. Spencer Jones. 2013. The amathiiiform Ctenostomata (phylum Bryozoa) of New Zealand
369 - including four new species, two of them of probable alien origin. *Zootaxa* 3647:75–95.
- 370 Gordon, D. P., I. G. Stuart, and J. D. Collen. 1994. Bryozoan fauna of the Kaipuke Siltstone, northwest Nelson:
371 a Miocene homologue of the modern Tasman Bay coralline bryozoan grounds. *New Zealand Journal of Geology*
372 *and Geophysics* 37:239–247.
- 373 Gordon, D. P., and P. D. Taylor. 2008. Systematics of the bryozoan genus *Macropora* (Cheilostomata). *Zoological*
374 *Journal of the Linnean Society* 153:115–146.
- 375 Gordon, D. P., and P. D. Taylor. 2010. New seamount- and ridge-associated cyclostome Bryozoa from New
376 Zealand. *Zootaxa* 2533:43–68.
- 377 Gordon, D. P., and P. D. Taylor. 2015. Bryozoa of the Early Eocene Tumaio Limestone, Chatham Island, New
378 Zealand. *Journal of Systematic Palaeontology* 13:983–1070.
- 379 Gordon, D. P., P. D. Taylor, and F. P. Bigey. 2009. Phylum Bryozoa - moss animals, sea mats, lace corals. Pages

- 380 271–297 in D. P. Gordon, editor. The New Zealand inventory of biodiversity. Volume 1. Kingdom Animalia.
381 Radiata, Lophotrochozoa, and Deuterostomia. Canterbury University Press, Christchurch.
- 382 Gorjunova, R. V., and A. V. Koromyslova. 2008. A new genus of the Arenigian bryozoans, *Lynnopora*, and its
383 systematic position in the family Revalotrypidae Gorjunova, 1988. *Paleontological Journal* 42:491–499.
- 384 Gómez, A., P. J. Wright, D. H. Lunt, J. M. Cancino, G. R. Carvalho, and R. N. Hughes. 2007. Mating trials
385 validate the use of DNA barcoding to reveal cryptic speciation of a marine bryozoan taxon. *Proceedings of the*
386 *Royal Society of London, B* 274:199–207.
- 387 Grange, K. R., and D. P. Gordon. 2005. The spread and potential impact of the introduced bryozoan *Biflustra*
388 *grandicella*. National Institute of Water and Atmospheric Research Final Research Report 31.
- 389 Greeley, R. 1969. Basally "uncalcified" zoaria of lunulitiform Bryozoa. *Journal of Paleontology* 43:252–256.
- 390 Gregory, J. W. 1896. Catalogue of the fossil Bryozoa in the Department of Geology, British Museum (Natural
391 History). The Jurassic Bryozoa. Pages 1–239. Trustees of the British Museum, London.
- 392 Grischenko, A. V., M. A. Dick, and S. F. Mawatari. 2007. Diversity and taxonomy of intertidal Bryozoa
393 (Cheilostomata) at Akkeshi Bay, Hokkaido, Japan. *Journal of Natural History* 41:1047–1161.
- 394 Grischenko, A. V., D. P. Gordon, A. Nojo, M. Kawamura, N. Kaneko, and S. F. Mawatari. 2004. New cheilostome
395 bryozoans from the Middle Miocene of Hokkaido, Japan, and the first fossil occurrences of *Kubaninella* and
396 *Hayamiellina* gen. nov. *Paleontological Research* 8:167–179.
- 397 Grischenko, A. V., S. F. Mawatari, and P. D. Taylor. 2000. Systematics and phylogeny of the cheilostome
398 bryozoan *Doryporella*. *Zoologia Scripta* 29:247–264.
- 399 Grischenko, A. V., J. E. Seo, and B. S. Min. 2010. *Flustrellidra armata* (Bryozoa: Ctenostomatida) — a new
400 species from the southern shoreline of Korea. *Zootaxa* 2684:25–35.
- 401 Guha, A. K. 2013. Tertiary Bryozoa from western Kachchh, Gujarat - a review. *Journal of the Palaeontological*
402 *Society of India* 58:3–15.
- 403 Guha, A. K., and K. Gopikrishna. 2005. New smittinid (Bryozoa, Cheilostomata) species from Tertiary sequences
404 of western Kachchh, Gujarat, India. *Freiberger Forschungshefte C507*:11–25.
- 405 Guha, A. K., and K. Gopikrishna. 2007a. Some fossil ascophoran bryozoan taxa from Tertiary sequences of
406 Western Kachchh, Gujarat. *Journal of the Paleontological Society of India* 52:195–222.
- 407 Guha, A. K., and K. Gopikrishna. 2007b. New calloporid (Bryozoa, Cheilostomata) species from Tertiary
408 sequences of western Kachchh, Gujarat. *Journal of the Geological Society of India* 70:121–130.
- 409 Guha, A. K., and K. Gopikrishna. 2007c. New celleporid (Bryozoa, Cheilostomata) species from Tertiary deposits
410 of western Kachchh, Gujarat, India. *Paläontologische Zeitschrift* 81:83–92.
- 411 Hageman, S. J. 1991a. *Worthenopora*: an unusual cryptostome (Bryozoa) that looks like a cheilostome. *Journal of*
412 *Paleontology* 65:648–661.
- 413 Hageman, S. J. 1991b. Approaches to systematic and evolutionary studies of perplexing groups: an example using
414 fenestrate Bryozoa. *Journal of Paleontology* 65:630–647.
- 415 Hageman, S. J. 1993. Effects of nonnormality on studies of morphological variation of a rhabdomesine bryozoan,
416 *Streblotrypa* (*Streblascopora*) *prisca* (Gabb and Horn). *The University of Kansas Paleontological Contributions*
417 4:1–13.
- 418 Hageman, S. J., P. E. Bock, Y. Bone, and B. McGowran. 1998. Bryozoan growth habits: Classification and
419 analysis. *Journal of Paleontology* 72:418–436.
- 420 Hageman, S. J., J. Lukasik, B. McGowran, and Y. Bone. 2003. Paleoenvironmental significance of Celleporaria
421 (Bryozoa) from modern and Tertiary cool-water carbonates of southern Australia. *Palaios* 18:510–527.
- 422 Hageman, S. J., and F. K. McKinney. 2010. Discrimination of fenestrate bryozoan genera in morphospace.
423 *Palaeontologia Electronica* 13:1–43.
- 424 Hageman, S. J., P. N. Wyse Jackson, A. R. Abernethy, and M. Steinhorsdottir. 2011. Calender scale, environ-
425 mental variation preserved in the skeletal phenotype of a fossil bryozoan (*Rhombopora blakei* n.sp.), from the
426 Mississippian of Ireland. *Journal of Paleontology* 85:853–870.

- 427 Hao, J., C. Li, X. Sun, and Q. Yang. 2005. Phylogeny and divergence time estimation of cheilostome bryozoans
428 based on mitochondrial 16S rRNA sequences. *Chinese Science Bulletin* 50:1205–1211.
- 429 Hara, U. 2001. Bryozoans from the Eocene of Seymour Island, Antarctic Peninsula. *Palaeontological Results of*
430 *the Polish Antarctic Expeditions. Part III. Palaeontologia Polonica* 60:33–156.
- 431 Hara, U. 2002. A new macroporid bryozoan from Eocene of Seymour Island, Antarctic Peninsula. *Polish Polar*
432 *Research* 23:213–225.
- 433 Hara, U., A. Ernst, and Z. Mikolajewski. 2009. Permian trepostome bryozoans from the Zechstein Main Dolomite
434 (Ca2) of Western Poland and NE Germany. *Geological Quartelry* 53:249–254.
- 435 Hara, U., and M. Jasionowski. 2012. The Early Sarmatian bryozoan *Celleporina medoborensis* sp. nov. from the
436 Medobory reefs of western Ukraine (Central Paratethys). *Geological Quarterly* 56:895–906.
- 437 Harmelin, J.-G. 1988. Les Bryozoaires, de bons indicateurs bathymétriques en paléocéologie? *Géologie Méditer-*
438 *ranéenne* 15:49–63.
- 439 Harmelin, J.-G. 2014. *Monoporella bouchardii* (Audouin & Savigny, 1826) (Bryozoa, Cheilostomata): a forgotten
440 taxon redescribed from Eastern Mediterranean material. *Cahiers de Biologie Marine* 55:91–99.
- 441 Harmelin, J.-G., G. Bitar, and H. Zibrowius. 2009. Smittinidae (Bryozoa, Cheilostomata) from coastal habitats of
442 Lebanon (Mediterranean Sea), including new and non-indigenous species. *Zoosystema* 31:163–187.
- 443 Harmelin, J.-G., A. N. Ostrovsky, J. P. Cáceres-Chamizo, and J. Sanner. 2011. Bryodiversity in the tropics:
444 taxonomy of *Microporella* species (Bryozoa, Cheilostomata) with personate maternal zooids from Indian Ocean,
445 Red Sea and southeast Mediterranean. *Zootaxa* 2798:1–30.
- 446 Harmer, S. F. 1900. A revision of the genus *Steganoporella*. *Quarterly Journal of Microscopical Science* 43:225–297.
- 447 Harmer, S. F. 1902. On the morphology of the Cheilostomata. *Quarterly Journal of Microscopical Science*
448 46:263–350.
- 449 Harmer, S. F. 1915. The Polyzoa of the Siboga Expedition, Part I. Entoprocta, Ctenostomata and Cyclostomata.
450 *Siboga Expeditie* 28a:1–180.
- 451 Harzhauser, M., A. Kroh, O. Mandic, W. E. Piller, U. Göhlich, M. Reuter, and B. Berning. 2007. Biogeographic
452 responses to geodynamics: a key study all around the Oligo–Miocene Tethyan Seaway. *Zoologischer Anzeiger*
453 246:241–256.
- 454 Harzhauser, M., and W. E. Piller. 2004. The Early Sarmatian - hidden seesaw changes. *Cour. Forsch.-Inst.*
455 *Senckenberg* 246:89–111.
- 456 Hastings, A. B. 1949. On the Polyzoan *Cellepora pulchra* Manzoni and the genus *Hippopleurifera*. *Journal of the*
457 *Linnean Society of London, Zoology* 41:521–528.
- 458 Hastings, A. B. 1964. The cheilostomatous Polyzoa *Neoethyris woosteri* (MacGillivray) and *Reginella doliaris*
459 (Maplestone). *Bulletin of the British Museum (Natural History), Zoology* 11:245–262.
- 460 Hayward, P. J. 1974. Studies on the cheilostome bryozoan fauna of the Aegean island of Chios. *Journal of Natural*
461 *History* 8:369–402.
- 462 Hayward, P. J. 1978. Bryozoa from the west European continental slope. *Journal of Zoology, London* 184:207–224.
- 463 Hayward, P. J. 1993. New species of cheilostomate Bryozoa from Antarctica and the Subantarctic southwest
464 Atlantic. *Journal of Natural History* 27:1409–1430.
- 465 Hayward, P. J., and K. B. Hansen. 1999. Three newly recognized cheilostomate bryozoans from the British sea
466 area. *Journal of the Marine Biological Association* 79:917–921.
- 467 Hayward, P. J., and J. S. Ryland. 1985. Systematic note on some British Cyclostomata (Bryozoa). *Journal of*
468 *Natural History* 19:1073–1078.
- 469 Hayward, P. J., and J. S. Ryland. 1990. Some Antarctic and Subantarctic species of Microporellidae (Bryozoa:
470 Cheilostomata). *Journal of Natural History* 24:1263–1287.
- 471 Hayward, P. J., and P. D. Taylor. 1984. Fossil and Recent Cheilostomata (Bryozoa) from the Ross Sea, Antarctica.
472 *Journal of Natural History* 18:71–94.
- 473 Hayward, P. J., and J. P. Thorpe. 1988a. Species of *Chaperiopsis* (Bryozoa, Cheilostomata) collected by Discovery

- 474 Investigations. *Journal of Natural History* 22:45–69.
- 475 Hayward, P. J., and J. P. Thorpe. 1988b. Species of *Arachnopusia* (Bryozoa: Cheilostomata) collected by
476 Discovery Investigations. *Journal of Natural History* 22:773–799.
- 477 Hayward, P. J., and J. E. Winston. 2011. Bryozoa collected by the United States Antarctic Research Program:
478 new taxa and new records. *Journal of Natural History* 45:2259–2338.
- 479 Herrera-Cubilla, A., M. H. Dick, J. Sanner, and J. B. C. Jackson. 2006. Neogene Cupuladriidae of tropical America.
480 I: Taxonomy of Recent Cupuladria from opposite sides of the Isthmus of Panama. *Journal of Paleontology*
481 80:245–263.
- 482 Herrera-Cubilla, A., M. H. Dick, J. Sanner, and J. B. C. Jackson. 2008. Neogene Cupuladriidae of tropical
483 America. II: Taxonomy of Recent Discoporella from opposite sides of the Isthmus of Panama. *Journal of*
484 *Paleontology* 82:279–298.
- 485 Herrera-Cubilla, A., and J. B. Jackson. 2014. Phylogeny of genus *Cupuladria* (Bryozoa, Cheilostomata) in the
486 Neogene of tropical America. *Journal of Paleontology* 88:851–894.
- 487 Hillmer, G. 2006. Erhard Voigt, 28. Juli 1905 - 22. November 2004. *Paläontologische Zeitschrift* 80:1–11.
- 488 Hinds, R. W. 1975. Growth mode and homeomorphism in cyclostome Bryozoa. *Journal of Paleontology* 49:875–910.
- 489 Holcová, K., and K. Zágorsek. 2008. Bryozoa, foraminifera and calcareous nannoplankton as environmental proxies
490 of the “bryozoan event” in the Middle Miocene of the Central Paratethys (Czech Republic). *Palaeogeography,*
491 *Palaeoclimatology, Palaeoecology* 267:216–234.
- 492 Holcová, K., K. Zágorsek, V. Jasková, and T. Lehotsky. 2007. The oldest Miocene Bryozoa from the Carpathian
493 Foredeep (boreholes Premyslovice). *Scripta Facultatis Scientiarum Naturalium Universitatis Masarykianae*
494 *Brunensis* 36:47–55.
- 495 Horowitz, A. S., and J. F. Pachut. 1993. Specific, generic, and familial diversity of Devonian bryozoans. *Journal*
496 *of Paleontology* 67:42–52.
- 497 Hunter, E., K. Shimizu, and N. Fusetani. 1999. Role of protein in larval swimming and metamorphosis of *Bugula*
498 *neritina* (Bryozoa: Cheilostomatida). *Marine Biology* 133:701–707.
- 499 Hurlbut, C. J. 1991. Community recruitment: settlement and juvenile survival of seven co-occurring species of
500 sessile marine invertebrates. *Marine Biology* 109:507–515.
- 501 Illies, G. 1981. Evolutionary trends in budding patterns of stomatopodid cyclostomatous Bryozoa and *Stomato-*
502 *poropsis* n. gen. *Oberrheinische geologische Abhandlungen* 30:27–42.
- 503 Illies, G. 1987. A new uniserial *Stomatopora* (Bryozoa Cyclostomata) from the Upper Cretaceous of northern
504 Germany. Pages 129–136 in J. R. P. Ross, editor. *Bryozoa: Present and past*. Western Washington University,
505 Bellingham.
- 506 Jablonski, D., S. Lidgard, and P. D. Taylor. 1997. Comparative ecology of bryozoan radiations: origin of novelties
507 in cyclostomes and cheilostomes. *Palaaios* 12:505–523.
- 508 Jackson, J. B. C., and A. O’Dea. 2013. Timing of the oceanographic and biological isolation of the Caribbean Sea
509 from the tropical eastern Pacific Ocean. *Bulletin of Marine Science* 89:779–800.
- 510 Jackson, J. B. C., J. E. Winston, and A. G. Coates. 1985. Niche breadth, geographic range, and extinction of
511 caribbean reef-associated cheilostome Bryozoa and Scleractinia. *Proceedings of the Fifth International Coral*
512 *Reef Congress* 4:151–158.
- 513 Jackson, P. N. W., M. M. Key, and S. P. Coakley. 2014. Epizoozoan trepostome bryozoans on nautiloids from the
514 Upper Ordovician (Katian) of the Cincinnati Arch region, USA: An assessment of growth, form, and water flow
515 dynamics. *Journal of Paleontology* 88:475–487.
- 516 Jagadeeshan, S., and A. O’Dea. 2012. Integrating fossils and molecules to study cupuladriid evolution in an
517 emerging isthmus. *Evolutionary Ecology* 26:337–355.
- 518 James, D. W., M. S. Foster, and J. O’Sullivan. 2006. Bryoliths (Bryozoa) in the Gulf of California. *Pacific Science*
519 60:117–124.
- 520 James, N. P., Y. Bone, L. B. Collins, and T. K. Kyser. 2001. Surficial sediments of the Great Australian Bight:

- 521 facies dynamics and oceanography on a vast cool-water carbonate shelf. *Journal of Sedimentary Research*
522 71:549–567.
- 523 James, N. P., D. A. Feary, C. Betzler, Y. Bone, A. E. Holbourn, Q. Li, H. Machiyama, J. A. T. Simo, and F.
524 Surlyk. 2004. Origin of late Pleistocene bryozoan reef mounds: Great Australian Bight. *Journal of Sedimentary*
525 *Research* 74:20–48.
- 526 Jelly, E. C. 1889. A synonymic catalogue of the Recent marine Bryozoa. Including fossil synonyms. Dulau &
527 Company, London.
- 528 Jiménez-Sánchez, A., P. D. Taylor, and J. B. Gómez. 2013. Palaeogeographical patterns in Late Ordovician
529 bryozoan morphology as proxies for temperature. *Bulletin of Geosciences* 88:417–426.
- 530 Johnson, M. E., R. S. Ramalho, B. G. Baarli, M. Cachão, C. M. da Silva, E. Mayoral, and A. Santos. 2014.
531 Miocene–Pliocene rocky shores on São Nicolau (Cape Verde Islands): contrasting windward and leeward biofacies
532 on a volcanically active oceanic island. *Palaeogeography, Palaeoclimatology, Palaeoecology* 395:131–143.
- 533 Jürgensen, T. 1968. Cheilostome Bryozoa from the Danian deposits at Vallengbaek, Denmark. *Meddelelser fra*
534 *Dansk Geologisk Forening* 18:187–204.
- 535 Jürgensen, T. 1971. *Tornipora*, a new cheilostome bryozoan from the Maastrichtian and Danian of Denmark.
536 *Bulletin of the Geological Society of Denmark* 20:390–396.
- 537 Kahle, J., G. Liebezeit, and G. Gerdes. 2003. Growth aspects of *Flustra foliacea* (Bryozoa, Cheilostomata) in
538 laboratory culture. *Hydrobiologia* 503:237–244.
- 539 Kaiser, S., H. J. Griffiths, D. K. A. Barnes, S. N. Brandão, A. Brandt, and P. E. O’Brien. 2011. Is there a
540 distinct continental slope fauna in the Antarctic? *Deep-Sea Research Part II: Topical Studies in Oceanography*
541 58:91–104.
- 542 Karklins, O. L. 1983. Ptilodictyoid Cryptostomata Bryozoa from the Middle and Upper Ordovician rocks of
543 central Kentucky. *The Paleontological Society Memoir* 14:1–31.
- 544 Keij, A. J. 1972. *Sylonika* and *Kylonisa*, two new Palaeogene bryozoan genera (Cheilostomata, Skytoniidae).
545 *Scripta Geologica* 11:1–15.
- 546 Key, M. M. 2002. Presentation of the Paleontological Society Medal to Alan H. Cheetham. *Journal of Paleontology*
547 76:783.
- 548 Key, M. M., P. N. Wyse Jackson, and S. H. Felton. 2016. Intracolony variation in colony morphology in reassembled
549 fossil ramose stenolaemate bryozoans from the Upper Ordovician (Katian) of the Cincinnati Arch region, USA.
550 *Journal of Paleontology* 90:400–412.
- 551 Key, M. M., P. N. Wyse Jackson, E. Håkansson, W. P. Patterson, and M. D. Moore. 2005. Gigantism in Permian
552 trepostomes from Greenland: testing the algal symbiosis hypothesis using $\delta^{13}\text{C}$ and $\delta^{18}\text{O}$ values. Pages 141–151
553 *in* H. I. Moyano, J. M. Cancino, and P. N. Wyse Jackson, editors. *Bryozoan studies 2005*. Balkema, Leiden.
- 554 Key, M. M., P. N. Wyse Jackson, and L. J. Vitiello. 2011. Stream channel network analysis applied to colony-wide
555 feeding structures in a Permian bryozoan from Greenland. *Paleobiology* 37:287–302.
- 556 Kidwell, S. M., and E. D. Gyllenhaal. 1998. Symbiosis, competition, and physical disturbance in the growth
557 histories of Pliocene cheilostome bryoliths. *Lethaia* 31:221–239.
- 558 Kieppura, M. 1962. Bryozoa from the Ordovician erratic boulders of Poland. *Acta Palaeontologica Polonica*
559 7:347–428.
- 560 Knowles, T., M. J. Leng, M. Williams, P. D. Taylor, H. J. Sloane, and B. Okamura. 2010. Interpreting seawater
561 temperature range using oxygen isotopes and zooid size variation in *Pentapora foliacea* (Bryozoa). *Marine*
562 *Biology* 157:1171–1180.
- 563 Koçak, F. 2007. A new alien bryozoan *Celleporaria brunnea* (Hincks, 1884) in the Aegean Sea (eastern Mediter-
564 ranean). *Scientia Marina* 71:191–195.
- 565 Koçak, F., and A. Önen. 2014. Checklist of Bryozoa on the coasts of Turkey. *Turkish Journal of Zoology*
566 38:880–891.
- 567 Koenig, J. W. 1958. Fenestrate Bryozoa in the Chouteau Group of central Missouri. *Journal of Paleontology*
568 32:126–143.

- 569 Koromyslova, A. V. 2014a. Morphological features and systematic position of the bryozoans *Onychocella rowei*
570 and *O. Mimosa* (Cheilostomata) from a Campanian erratic block (Belarus). *Paleontological Journal* 48:275–286.
- 571 Koromyslova, A. V. 2014b. The earliest calcified opercula of bryozoans of the order Cheilostomata. *Paleontological*
572 *Journal* 48:587–593.
- 573 Koromyslova, A. V., and A. V. Pakhnevich. 2016. New species of *Pachydermopora* Gordon, 2002 and *Beisselina*
574 *Canu*, 1913 (Bryozoa: Cheilostomida) from a Campanian erratic block (Belarus) and their micro-CT investigation.
575 *Paleontological Journal* 50:41–53.
- 576 Koromyslova, A. V., and E. A. Shcherbinina. 2015. New data on the morphology and age of the bryozoans of
577 the genus *Onychocella* from the Campanian-Maastrichtian of Turkmenistan and Uzbekistan. *Paleontological*
578 *Journal* 49:474–484.
- 579 Kuklinski, P. 2013. Biodiversity and abundance patterns of rock encrusting fauna in a temperate fjord. *Marine*
580 *Environmental Research* 87-88:61–72.
- 581 Kuklinski, P., A. V. Grischenko, and S. C. Jewett. 2015. Two new species of the cheilostome bryozoan *Cheilopora*
582 from the Aleutian Islands. *Zootaxa* 3963:434–442.
- 583 Kuklinski, P., B. Gulliksen, O. J. Lønne, and J. M. Weslawski. 2006. Substratum as a structuring influence on
584 assemblages of Arctic bryozoans. *Polar Biology* 29:652–661.
- 585 Lagaaij, R. 1959. Some species of Bryozoa new to the Bowden Beds, Jamaica, B.W.I. *Micropaleontology* 5:482–486.
- 586 Landing, E., J. B. Antcliffe, M. D. Brasier, and A. B. English. 2015. Distinguishing Earth’s oldest known
587 bryozoan (*Pywackia*, late Cambrian) from pennatulacean octocorals (MesozoicRecent). *Journal of Paleontology*
588 89:292–317.
- 589 Landing, E., A. English, and J. D. Keppie. 2010. Cambrian origin of all skeletalized metazoan phyla - discovery
590 of Earth’s oldest bryozoans (Upper Cambrian, southern Mexico). *Geology* 38:547–550.
- 591 Larwood, G. P. 1962. The morphology and systematics of some Cretaceous cribrimorph Polyzoa (Pelmatoporinae).
592 *Bulletin of the British Museum (Natural History)*, *Geology* 6:1–285.
- 593 Larwood, G. P. 1969. Frontal calcification and its function on some Cretaceous and Recent cribrimorph and other
594 cheilostome Bryozoa. *Bulletin of the British Museum (Natural History)*, *Zoology* 18:171–182.
- 595 Larwood, G. P., A. W. Medd, D. E. Owen, and R. Tavener-Smith. 1967. Bryozoa. Pages 295–379 *in* *The Fossil*
596 *Record*. Geological Society of London, London.
- 597 Lidgard, S., M. C. Carter, M. H. Dick, D. P. Gordon, and A. N. Ostrovsky. 2011. Division of labor and recurrent
598 evolution of polymorphisms in a group of colonial animals. *Evolutionary Ecology* 26:233–257.
- 599 Lidgard, S., and J. B. Jackson. 1989. Growth in encrusting cheilostome bryozoans: I. Evolutionary trends.
600 *Paleobiology* 15:255–282.
- 601 Livingstone, A. A. 1926. Studies on Australian Bryozoa. No. 4. *Records of the Australian Museum* 15:167–176.
- 602 Lodola, A., J. Ferrario, and A. Occhipinti-Ambrogi. 2015. Further Mediterranean expansion of the non-indigenous
603 bryozoan *Celleporaria brunnea*: multiple records along the Italian coasts. *Scientia Marina* 79:263–274.
- 604 Lombardi, C., S. Cocito, K. Hiscock, A. Occhipinti Ambrogi, M. Setti, and P. D. Taylor. 2008. Influence of
605 seawater temperature on growth bands, mineralogy and carbonate production in a bioconstructional bryozoan.
606 *Facies* 54:333–342.
- 607 Lombardi, C., S. Cocito, A. Occhipinti Ambrogi, and K. Hiscock. 2006. The influence of seawater temperature on
608 zooid size and growth rate in *Pentapora fascialis* (Bryozoa: Cheilostomata). *Marine Biology* 149:1103–1109.
- 609 Lombardi, C., M. C. Gambi, C. Vasapollo, P. D. Taylor, and S. Cocito. 2011. Skeletal alterations and polymorphism
610 in a Mediterranean bryozoan at natural CO₂ vents. *Zoomorphology* 130:135–145.
- 611 Lombardi, C., P. D. Taylor, and S. Cocito. 2010. Systematics of the Miocene–Recent bryozoan genus *Pentapora*
612 (Cheilostomata). *Zoological Journal of the Linnean Society* 160:17–39.
- 613 Lonsdale, W. 1845a. Corals from the Tertiary formations of North America. I. Account of ten species of *Polyparia*
614 obtained from the Miocene Tertiary formations of North America. *Quarterly Journal of the Geological Society*
615 of London 1:495–509.

- 616 Lonsdale, W. 1845b. Account of six species of *Polyparia* obtained from Timber Creek, New Jersey. Quarterly
617 Journal of the Geological Society of London 1:65–75.
- 618 López de la Cuadra, C. M., and J. C. García-Gómez. 2000. The cheilostomate Bryozoa (Bryozoa: Cheilostatida)
619 collected by the Spanish 'Antártida 8611' Expedition to the Scotia Arc and South Shetland Islands. Journal of
620 Natural History 34:755–772.
- 621 López Gappa, J. J. 1989. Overgrowth competition in an assemblage of encrusting bryozoans settled on artificial
622 substrata. Marine Ecology Progress Series 51:121–130.
- 623 Ma, J., P. D. Taylor, F.-S. Xia, and R. Zhan. 2015. The oldest known bryozoan: *Prophyllodictya* (Cryptosto-
624 mata) from the lower Tremadocian (Lower Ordovician) of Liujiachang, south-western Hubei, central China.
625 Palaeontology 58:925–934.
- 626 Machiyama, H., T. Yamada, N. Kaneko, Y. Iryo, K. Odawara, R. Asami, H. Matsuda, S. F. Mawatari, Y. Bone,
627 and N. P. James. 2003. Carbon and oxygen isotopes of cool-water bryozoans from the Great Australian Bight
628 and their paleoenvironmental significance. Pages 1–29 in A. C. Hine, D. A. Feary, and M. J. Malone, editors.
629 Proceedings of the ocean drilling programme, scientific results.
- 630 Mackie, J. A. 2003. A molecular analysis of bryozoan dispersal. PhD thesis, University of Melbourne, Melbourne.
- 631 Mackie, J. A., J. A. Darling, and J. B. Geller. 2012. Ecology of cryptic invasions: latitudinal segregation among
632 *Watersipora* (Bryozoa) species. Scientific Reports 2:1–10.
- 633 Malecki, J. 1964. On two new genera of Bryozoa Cheilostomata from the Tortonian of Poland. Acta Palaeontologica
634 Polonica 9:499–512.
- 635 Marintsch, E. J. 1981. Taxonomic reevaluation of *Prasopora simulatrix* Ulrich (Bryozoa: Trepostomata). Journal
636 of Paleontology 55:957–961.
- 637 Maturo, F. J. S., and T. J. M. Schopf. 1968. Ectoproct and entoproct type material: reexamination of species
638 from New England and Bermuda named by A.E. Verrill, J.W. Dawson and E. Desor. Postilla 120:1–95.
- 639 McColloch, M. E., E. H. Gilmour, and E. M. Snyder. 1994. The order Fenestrata (Bryozoa) of the Toroweap
640 Formation (Permian), southern Nevada. Journal of Paleontology 68:746–762.
- 641 McGhee, G. R., and F. K. McKinney. 2002. A theoretical morphologic analysis of ecomorphologic variation in
642 *Archimedes* helical colony form. Palaios 17:556–570.
- 643 McKinney, F. K. 1983. Ectoprocta (Bryozoa) from the Permian Kaibab Formation, Grand Canyon National Park,
644 Arizona. Fieldiana Geology, N.S. 13:1–17.
- 645 McKinney, F. K. 2009. Bryozoan-hyroid symbiosis and a new ichnogenus, *Caupokeras*. Ichnos 16:193–201.
- 646 McKinney, F. K., and D. W. Burdick. 2001. A rare, larval-founded colony of the bryozoan *Archimedes* from the
647 Carboniferous of Alabama. Palaeontology 44:855–859.
- 648 McKinney, F. K., and A. Jaklin. 2001. Sediment accumulation in a shallow-water meadow carpeted by a small
649 erect bryozoan. Sedimentary Geology 145:397–410.
- 650 McKinney, F. K., and J. Kríz. 1986. Lower Devonian Fenestrata (Bryozoa) of the Prague Basin, Barrandian Area,
651 Bohemia, Czechoslovakia. Fieldiana 1368.
- 652 McKinney, F. K., M. R. A. Listokin, and C. D. Phifer. 1986. Flow and polypide distribution in the cheilostome
653 bryozoan *Bugula* and their inference in *Archimedes*. Lethaia 19:81–93.
- 654 McKinney, F. K., and G. R. McGhee. 2003. Evolution of erect helical colony form in the Bryozoa: phylogenetic,
655 functional, and ecological factors. Biological Journal of the Linnean Society 80:235–260.
- 656 McKinney, F. K., and M. J. McKinney. 1994. Preferences for settlement conditions by larvae of *Schizotheca*
657 *serratimargo* (Hincks, 1886), an erect bryozoan from protected habitats. Pages 119–124 in P. J. Hayward, J. S.
658 Ryland, and P. D. Taylor, editors. Biology and Palaeobiology of Bryozoans. Olsen & Olsen, Fredensborg.
- 659 McKinney, F. K., M. J. McKinney, and M. R. A. Listokin. 1987. Erect bryozoans are more than baffling: enhanced
660 sedimentation rate by a living unilaminate branched bryozoan and possible implications for fenestrate bryozoan
661 mudmounds. Palaios 2:41–47.
- 662 McKinney, F. K., and P. D. Taylor. 2001. Bryozoan generic extinctions and originations during the last one

- 663 hundred million years. *Palaeontologia Electronica* 4:1–26.
- 664 McKinney, F. K., and P. D. Taylor. 2003. Palaeoecology of free-lying domal bryozoan colonies from the Upper
665 Eocene of southeastern USA. *Acta Palaeontologica Polonica* 48:447–462.
- 666 McKinney, F. K., P. D. Taylor, and V. A. Zullo. 1993. Lyre-shaped hornerid bryozoan colonies: homeomorphy in
667 colony form between Paleozoic Fenestrata and Cenozoic Cyclostomata. *Journal of Paleontology* 67:343–354.
- 668 McKinney, F. K., and P. N. Wyse Jackson. 2010. *Phylloporina* Ulrich in Foerste, 1887 (Bryozoa, Fenestrata,
669 Phylloporinina): proposed designation of *Retepora trentoniensis* Nicholson, 1875 as the type species. *Bulletin*
670 *of Zoological Nomenclature* 67:38–43.
- 671 McKinney, F. K., and P. N. Wyse Jackson. 2015. Part G, Revised, Volume 2, Chapter 8H: Names of indeterminate,
672 unrecognizable, or excluded genera previously assigned to Fenestrata. *Treatise Online* 64:1–5.
- 673 McShea, D. W., and E. P. Venit. 2002. Testing for bias in the evolution of coloniality: a demonstration in
674 cyclostome bryozoans. *Paleobiology* 28:308–327.
- 675 Menzies, R. J. 1963. Abyssal Bryozoa collected by expeditions of the Lamont Geological Observatory. 1.
676 Bicelliariellidae (Bugulidae of authors), Kinetoskias. *American Museum Novitates* 2130:1–8.
- 677 Mesentseva, O. P. 2016. Cystoporida (Bryozoa) from the Emsian Stage (Lower Devonian) of the Salair Ridge and
678 Gorny and Rudny Altai. *Paleontological Journal* 50:376–387.
- 679 Messina, C., M. A. Rosso, F. Sciuto, I. Di Geronimo, W. Nemeč, T. Di Dio, R. Di Geronimo, R. Maniscalco, and
680 R. Sanfilippo. 2007. Anatomy of a transgressive systems tract revealed by integrated sedimentological and
681 palaeoecological study: the Barcellona Pozzo di Gotto Basin, northeastern Sicily, Italy. Pages 367–400 *in* G.
682 Nichols, E. A. Williams, and C. Paola, editors. *Sedimentary Processes, Environments and Basins – A Tribute*
683 *to Peter Friend*. Blackwell Publishing, Malden, MA, USA.
- 684 Moissette, P. 1996. The cheilostomate bryozoan *Batopora rosula* (Reuss, 1848): a paleobathymetric indicator in
685 the Mediterranean Neogene. Pages 193–198 *in* D. P. Gordon, A. M. Smith, and J. A. Grant-Mackie, editors.
686 *Bryozoans in space and time*. NIWA, Wellington.
- 687 Moissette, P. 2000. Changes in bryozoan assemblages and bathymetric variations. Examples from the Messinian
688 of northwest Algeria. *Palaeogeography, Palaeoclimatology, Palaeoecology* 155:305–326.
- 689 Moissette, P., J.-J. Cornée, and E. Koskeridou. 2010. Pleistocene rolling stones or large bryozoan nodules in a
690 mixed siliciclastic-carbonate environment (Rhodes, Greece). *Palaios* 25:24–39.
- 691 Moissette, P., A. Dulai, G. Escarguel, M. Kázmér, P. Müller, and J.-P. Saint Martin. 2007. Mosaic of environments
692 recorded by bryozoan faunas from the Middle Miocene of Hungary. *Palaeogeography, Palaeoclimatology,*
693 *Palaeoecology* 252:530–556.
- 694 Moissette, P., A. Dulai, and P. Müller. 2006. Bryozoan faunas in the Middle Miocene of Hungary: biodiversity
695 and biogeography. *Palaeogeography, Palaeoclimatology, Palaeoecology* 233:300–314.
- 696 Moissette, P., and S. Pouyet. 1987. Bryozoan faunas and the Messinian salinity crisis. *Annales Instituti Geologici*
697 *Publici Hungarici* 70:447–453.
- 698 Moissette, P., and J.-P. Saint-Martin. 1995. Bryozoaires des milieux récifaux miocènes du sillon sud-rifain au
699 Maroc. *Lethaia* 28:271–283.
- 700 Montoya-Cadavid, E., P. Flórez-Romero, and J. E. Winston. 2007. Checklist of the marine Bryozoa of the
701 Colombian Caribbean. *Biota Colombiana* 8:159–184.
- 702 Morozova, I. P., O. B. Weis, and G. Eacki. 2002. Emergence and extinction of the Givetian to Frasnian bryozoan
703 faunas in the Kostomloty facies zone, Holy Cross Mountains, Poland. *Acta Palaeontologica Polonica* 47:307–317.
- 704 Moyano, H. I. 1983. Southern Pacific Bryozoa: a general view with emphasis on Chilean species. *Gayana Zoologia*
705 46:1–45.
- 706 Moyano, H. I. 1999. Magellan Bryozoa: a review of the diversity and of the Subantarctic and Antarctic
707 zoogeographical links. *Scientia Marina* 63:219–226.
- 708 Mukai, H., K. Terakado, and C. G. Reed. 1997. Bryozoa. Pages 45–206 *in* F. W. Harrison and R. M. Woollacott,
709 editors. *Microscopic Anatomy of Invertebrates*. Wiley-Liss, New York.

- 710 Nakrem, H. A., and N. Spjeldnaes. 1995. *Ramipora hochstetteri* Toulou, 1875 (Bryozoa, Cystoporata), from the
711 Permian of Svalbard. *Journal of Paleontology* 69:831–838.
- 712 Newton, G. B. 1971. Rhabdomesid bryozoans of the Wreford megacyclothem (Wolfcampian, Permian) of Nebraska,
713 Kansas, and Oklahoma. *University of Kansas Paleontological Contributions* 56:1–71.
- 714 Nikulina, E. A. 2006. *Electra korobokkura* sp. n., a new species of cheilostome bryozoan from the Pacific coast of
715 Hokkaido, Japan. *Invertebrate Zoology* 3:23–31.
- 716 Nikulina, E. A. 2007. *Einhornia*, a new genus for electrids formerly classified as the *Electra crustulenta* species
717 group (Bryozoa, Cheilostomata). *Schriften des Naturwissenschaftlichen Vereins für Schleswig-Holstein* 69:29–40.
- 718 Nikulina, E. A. 2008a. Taxonomy and ribosomal DNA-based phylogeny of the *Electra crustulenta* species group
719 (Bryozoa: Cheilostomata) with revision of Borg’s varieties and description of *Electra moskvikvendi* sp. nov.
720 from the western Baltic Sea. *Organisms, Diversity & Evolution* 8:215–229.
- 721 Nikulina, E. A. 2008b. *Electra scuticifera* sp. nov.: redescription of *Electra pilosa* from New Zealand as a
722 new species (Bryozoa, Cheilostomata). *Schriften des Naturwissenschaftlichen Vereins für Schleswig-Holstein*
723 70:91–98.
- 724 Nikulina, E. A. 2010. Three new genera of Electridae (Bryozoa): *Arbopercula*, *Osburnea*, and *Arbocuspis*. *Schriften*
725 *des Naturwissenschaftlichen Vereins für Schleswig-Holstein* 72:25–28.
- 726 Nikulina, E. A., A. N. Ostrovsky, and M. Claereboudt. 2012. A new species of the genus *Electra* (Bryozoa,
727 Cheilostomata) from southern Oman, Arabian Sea. Pages 203–216 in A. Ernst, P. Schäfer, and J. Scholz,
728 editors. *Bryozoan studies 2010*. Springer, Berlin.
- 729 Nikulina, E. N., R. Hanel, and P. Schäfer. 2007. Cryptic speciation and paraphyly in the cosmopolitan bryozoan
730 *Electra pilosa* - impact of the Tethys closing on species evolution. *Molecular Phylogenetics and Evolution*
731 45:765–776.
- 732 Nitecki, M. H., and F. K. McKinney. 1975. *Cucumulites* Gurley 1884: a bryozoan, not a sponge. *Journal of*
733 *Paleontology* 49:561–563.
- 734 Novak, V., N. Santodomingo, A. Rösler, E. Di Martino, J. C. Braga, P. D. Taylor, K. G. Johnson, and W.
735 Renema. 2013. Environmental reconstruction of a late Burdigalian (Miocene) patch reef in deltaic deposits
736 (East Kalimantan, Indonesia). *Palaeogeography, Palaeoclimatology, Palaeoecology* 374:110–122.
- 737 Nye, O. B., and D. V. Lemone. 1978. Multilaminar growth in *Reptomulticava texana*, a new species of cyclostome
738 Bryozoa. *Journal of Paleontology* 52:830–845.
- 739 Okamura, B., A. O’Dea, P. D. Taylor, and A. Taylor. 2013. Evidence of El Niño/La Niña–Southern Oscillation
740 variability in the Neogene-Pleistocene of Panama revealed by a new bryozoan assemblage-based proxy. *Bulletin*
741 *of Marine Science* 89:857–867.
- 742 Olempska, E. 2012. Exceptional soft-tissue preservation in boring ctenostome bryozoans and associated “fungal”
743 borings from the Early Devonian of Podolia, Ukraine. *Acta Palaeontologica Polonica* 57:925–940.
- 744 Osburn, R. C. 1919. Bryozoa of the Crocker Land Expedition. *Bulletin of the American Museum of Natural*
745 *History* 41:603–624.
- 746 Osburn, R. C. 1950. Bryozoa of the Pacific Coast of America. Part 1, Cheilostomata-Anasca. *Allan Hancock*
747 *Pacific Expeditions* 14:1–269.
- 748 Osburn, R. C. 1952. Bryozoa of the Pacific Coast of America. Part 2, Cheilostomata Ascophora. *Allan Hancock*
749 *Pacific Expeditions* 14:271–611.
- 750 Osburn, R. C. 1953. Bryozoa of the Pacific Coast of America. Part 3, Cyclostomata, Ctenostomata, Entoprocta,
751 and Addenda. *Report of the Allan Hancock Pacific Expeditions* 14:613–841.
- 752 Ostrovsky, A. N. 1998. Variability of oeciostome shape in Antarctic idmidroniform bryozoans (Bryozoa:
753 Cyclostomatida). *Zoologischer Anzeiger* 237:97–106.
- 754 Ostrovsky, A. N. 2004. Brood chambers (ovicells) of cheilostome bryozoans (Bryozoa: Gymnolaemata): structure,
755 research history, and modern problematics. *Russian Journal of Marine Biology* 30:43–55.
- 756 Ostrovsky, A. N. 2008. External versus internal and self- versus cross-fertilization in Bryozoa: transformation of
757 the view and evolutionary considerations. Pages 103–115 in P. N. Wyse Jackson and M. E. Spencer Jones,

- 758 editors. *Annals of Bryozoology: Aspects of the History of Research on Bryozoans*. International Bryozoology
759 Association, Dublin.
- 760 Ostrovsky, A. N. 2013. Evolution of Sexual Reproduction in Marine Invertebrates. Example of Gymnolaemate
761 Bryozoans. Page 356. Springer, Dordrecht.
- 762 Ostrovsky, A. N., D. P. Gordon, and S. Lidgard. 2009. Independent evolution of matrotrophy in the major classes
763 of Bryozoa: transitions among reproductive patterns and their ecological background. *Marine Ecology Progress
764 Series* 378:113–124.
- 765 Ostrovsky, A. N., A. V. Grischenko, P. D. Taylor, P. E. Bock, and S. F. Mawatari. 2006. Comparative anatomical
766 study of internal brooding in three anascan bryozoans (Cheilostomata) and its taxonomic and evolutionary
767 implications. *Journal of Morphology* 267:739–749.
- 768 Ostrovsky, A. N., P. Schäfer, and D. P. Gordon. 2003. Ultrastructure and development of the ooecial walls in
769 some calloporid bryozoans (Gymnolaemata: Cheilostomata). *Zoologischer Anzeiger* 242:223–240.
- 770 Ostrovsky, A. N., and N. N. Shunatova. 2002. Colonial behaviour and group zooidal reactions in Bryozoa: history
771 of research. Pages 185–199 *in* P. N. Wyse Jackson and M. E. Spencer Jones, editors. *Annals of Bryozoology:
772 Aspects of the History of Research on Bryozoans*. International Bryozoology Association, Dublin.
- 773 Ostrovsky, A. N., N. N. Shunatova, and I. I. Antipenko. 2002. Historical review on individual autozooidal
774 behaviour and feeding mechanisms in Bryozoa. Pages 201–227 *in* P. N. Wyse Jackson and M. E. Spencer Jones,
775 editors. *Annals of Bryozoology: Aspects of the History of Research on Bryozoans*. International Bryozoology
776 Association, Dublin.
- 777 Ostrovsky, A. N., and P. D. Taylor. 2004. Systematics of Upper Cretaceous calloporid bryozoans with primitive
778 spinose ovicells. *Palaeontology* 47:775–793.
- 779 Ostrovsky, A. N., and P. D. Taylor. 2005. Brood chambers constructed from spines in fossil and Recent cheilostome
780 bryozoans. *Zoological Journal of the Linnean Society* 144:317–361.
- 781 Ostrovsky, A. N., P. D. Taylor, M. H. Dick, and S. F. Mawatari. 2008a. Pre-Cenomanian cheilostome Bryozoa:
782 current state of knowledge. Pages 69–74 *in* H. Okada, S. F. Mawatari, N. Suzuki, and P. Gautam, editors.
783 *Origin and Evolution of Natural Diversity*. 21st Century COE for Neo-Science of Natural History, Hokkaido
784 University.
- 785 Ostrovsky, A. N., N. Vávra, and J. S. Porter. 2008b. Sexual reproduction in gymnolaemate Bryozoa: History and
786 perspectives of the research. *in* P. N. Wyse Jackson and M. E. Spencer Jones, editors. *Annals of Bryozoology 2:
787 Aspects of the History of Research on Bryozoans*. International Bryozoology Association, Dublin.
- 788 O’Dea, A. 2006. Asexual propagation in the marine bryozoan *Cupuladria exfragminis*. *Journal of Experimental
789 Marine Biology and Ecology* 335:312–322.
- 790 O’Dea, A. 2009. Relation of form to life habit in free-living cupuladriid bryozoans. *Aquatic Biology* 7:1–18.
- 791 O’Dea, A., E. Håkansson, P. D. Taylor, and B. Okamura. 2011. Environmental change prior to the K–T
792 boundary inferred from temporal variation in the morphology of cheilostome bryozoans. *Palaeogeography,
793 Palaeoclimatology, Palaeoecology* 308:502–512.
- 794 O’Dea, A., A. Herrera-Cubilla, H. Fortunato, and J. B. C. Jackson. 2004. Life history variation in cupuladriid
795 bryozoans from either side of the Isthmus of Panama. *Marine Ecology Progress Series* 280:145–161.
- 796 O’Dea, A., and J. B. C. Jackson. 2002. Bryozoan growth mirrors contrasting seasonal regimes across the Isthmus
797 of Panama. *Palaeogeography, Palaeoclimatology, Palaeoecology* 185:77–94.
- 798 O’Dea, A., J. B. C. Jackson, P. D. Taylor, and F. Rodríguez. 2008. Modes of reproduction in Recent and fossil
799 cupuladriid bryozoans. *Palaeontology* 51:847–864.
- 800 O’Dea, A., and B. Okamura. 2000. Cheilostome bryozoans as indicators of seasonality in the Neogene epicontinental
801 seas of western Europe. Pages 316–320 *in* A. Herrera Cubilla and J. B. C. Jackson, editors. *Proceedings of
802 the 11th International Bryozoology Association Conference*. Smithsonian Tropical Research Institute, Balboa,
803 Republic of Panama, Panama City.
- 804 O’Dea, A., A. N. Ostrovsky, and F. Rodríguez. 2010. Embryonic brooding and clonal propagation in tropical
805 eastern Pacific cupuladriid bryozoans. *Journal of the Marine Biological Association of the United Kingdom*

- 806 90:291–299.
- 807 Pathmanaban, O. N., J. S. Porter, and I. R. White. 2005. Dogger Bank itch in the eastern English Channel:
808 a newly described geographical distribution of an old problem. *Clinical and Experimental Dermatology*
809 30:622–626.
- 810 Perez, F. M., and W. C. Banta. 1996. How does *Cellaria* get out of its box? A new cheilostome hydrostatic
811 mechanism (Bryozoa: Cheilostomata). *Invertebrate Biology* 115:162–169.
- 812 Pérez, L. M., M. Griffin, G. Pastorino, J. J. López Gappa, and M. O. Manceñido. 2015. Redescription and
813 palaeoecological significance of the bryozoan *Hippoporidra patagonica* (Pallaroni, 1920) in the San Julián
814 Formation (late Oligocene) of Santa Cruz province, Argentina. *Alcheringa* 39:1–7.
- 815 Pérez, L. M., J. J. López-Gappa, and M. Griffin. 2015. New and little-known bryozoans from Monte León
816 Formation (early Miocene, Argentina) and their paleobiogeographic relationships. *Journal of Paleontology*
817 89:956–965.
- 818 Pirrie, D., J. A. Crame, J. B. Riding, A. R. Butcher, and P. D. Taylor. 1997. Miocene glaciomarine sedimentation
819 in the northern Antarctic Peninsula region: the stratigraphy and sedimentology of the Hobbs Glacier Formation,
820 James Ross Island. *Geological Magazine* 136:745–762.
- 821 Pizzaferrri, C. 2010. New specimens of *Prenantia cheilostoma* (Manzoni) from the Pliocene of Castell'Arquato
822 area (Western Emilia Region, N Italy). (Bryozoa Gymnolaemata Cheilostomata Smittinidae). *Quaderno di*
823 *Studi e Notizie di Storia Naturale della Romagna* 31:89–117.
- 824 Pizzaferrri, C., and B. Berning. 2007. Taxonomic notes on some cheilostome Bryozoa from the Pliocene of the
825 Western Emilia region (N Italy). *Rivista Italiana di Paleontologia e Stratigrafia* 113:97–108.
- 826 Pizzaferrri, C., and G. Braga. 2000. Nuove osservazioni sullo sviluppo astogenetico di *Batopora rosula* (Reuss),
827 Bryozoa Cheilostomatida del Miocene del Pedepennino parmense. *Annali del Museo Civico di Rovereto*
828 14:55–88.
- 829 Pohowsky, R. A. 1974. Notes on the study and nomenclature of boring Bryozoa. *Journal of Paleontology*
830 48:557–564.
- 831 Poignant, A. 1991. *Unifissurinella*, new genus of Microproblematica from the Upper Eocene of southwestern
832 France. *Micropaleontology* 37:95–97.
- 833 Porter, J. S., J. S. Ryland, and G. R. Carvalho. 2002. Micro- and macrogeographic genetic structure in bryozoans
834 with different larval strategies. *Journal of Experimental Marine Biology and Ecology* 272:119–130.
- 835 Powers, C. M., and J. F. Pachut. 2008. Diversity and distribution of Triassic bryozoans in the aftermath of the
836 end-Permian mass extinction. *Journal of Paleontology* 82:362–371.
- 837 Pushkin, V. I., and L. E. Popov. 1999. Early Ordovician bryozoans from north-western Russia. *Palaeontology*
838 42:171–189.
- 839 Pushkin, V. I., and L. E. Popov. 2005. Two enigmatic bryozoans from the Middle Ordovician of the East Baltic.
840 *Palaeontology* 48:1065–1074.
- 841 Ramalho, L. V., G. Muricy, and P. D. Taylor. 2009. Cyclostomata (Bryozoa, Stenolaemata) from Rio de Janeiro
842 State, Brazil. *Zootaxa* 2057:32–52.
- 843 Reverter Gil, O., and E. Fernández Pulpeiro. 1995. *Chaperiopsis cristata* (Busk, 1884) (Bryozoa, Cheilostomata)
844 is *Chaperiopsis annulus* (Manzoni, 1870). The occurrence of this species from the European Atlantic coast.
845 *Journal of Natural History* 29:1057–1065.
- 846 Reverter Gil, O., and E. Fernández Pulpeiro. 2005. A new genus of cyclostome bryozoan from the European
847 Atlantic coast. *Journal of Natural History* 39:2379–2387.
- 848 Reverter-Gil, O., and E. Fernández-Pulpeiro. 2007. Species of genus *Schizotheca* Hincks (Bryozoa, Cheilostomata)
849 described in the Atlantic-Mediterranean region, with notes on some species of *Parasmittina* Osburn. *Journal of*
850 *Natural History* 41:1929–1953.
- 851 Reverter-Gil, O., and J. Souto. 2015. Redescription of some species of Bryozoa described by J. Jullien and L.
852 Calvet in the NE Atlantic. *European Journal of Taxonomy* 157:1–17.
- 853 Reverter-Gil, O., J. Souto, and E. Fernández Pulpeiro. 2009. Three new species of Iberian cheilostomate Bryozoa.

- 854 Journal of the Marine Biological Association of the United Kingdom 89:1499–1506.
- 855 Reverter-Gil, O., J. Souto, and E. Fernández-Pulpeiro. 2012. New and little known species of Bryozoa from
856 Iberian Atlantic waters. *Zoosystema* 34:157–170.
- 857 Reverter-Gil, O., J. Souto, and E. Fernández-Pulpeiro. 2014. Annotated checklist of Recent marine Bryozoa from
858 continental Portugal. *Nova Acta Cientifica Compostelana (Biología)* 21:1–55.
- 859 Rogick, M. D. 1956. Studies on marine Bryozoa. VIII. *Exochella longirostris* Jullien 1888. *Biological Bulletin*
860 111:123–128.
- 861 Rogick, M. D. 1960. Studies on marine Bryozoa. XIII. Two new genera and new species from Antarctica. *Biological*
862 *Bulletin* 119:479–493.
- 863 Ross, J. R. P. 1960. Larger cryptostome Bryozoa of the Ordovician and Silurian, Anticosti Island, Canada - Part
864 I. *Journal of Paleontology* 34:1057–1076.
- 865 Ross, J. R. P. 1961. Larger cryptostome Bryozoa of the Ordovician and Silurian, Anticosti Island, Canada - Part
866 II. *Journal of Paleontology* 35:331–344.
- 867 Ross, J. R. P. 1963. New Ordovician species of Chazyan trepostome and cryptostome Bryozoa. *Journal of*
868 *Paleontology* 37:57–63.
- 869 Ross, J. R. P. 1964. Champlainian cryptostome Bryozoa from New York State. *Journal of Paleontology* 38:1–32.
- 870 Ross, J. R. P. 1967. Champlainian Ectoprocta (Bryozoa), New York State. *Journal of Paleontology* 41:632–648.
- 871 Ross, J. R. P. 1969. Champlainian (Ordovician) Ectoprocta (Bryozoa), New York State, Part II. *Journal of*
872 *Paleontology* 43:257–284.
- 873 Ross, J. R. P. 1970. Distribution, paleoecology and correlation of Champlainian Ectoprocta (Bryozoa), New York
874 State, Part III. *Journal of Paleontology* 44:346–382.
- 875 Rosso, A. 1999. Recent and fossil species of *Characodoma* Maplestone, 1900 (Bryozoa) from the Mediterranean
876 with description of two new species. *Journal of Natural History* 33:415–437.
- 877 Rosso, A. 2002. *Amphiblestrum* Gray, 1848 (Bryozoa, Cheilostomata) from the Atlantic-Mediterranean area, with
878 description of a new species. *Journal of Natural History* 36:1489–1508.
- 879 Rosso, A. 2003. Bryozoan diversity in the Mediterranean Sea. *Biogeographia* 24:227–250.
- 880 Rosso, A. 2004. Two new species of *Phylactella* (Bryozoa, Cheilostomatida) from the Mediterranean area belonging
881 to the *P. labrosa* (Busk) complex of species. *Journal of Natural History* 38:2655–2668.
- 882 Rosso, A. 2008. Mediterranean setoselliniforms and their exploitation of small sized substrata. Pages 261–268 *in*
883 S. J. Hageman, M. M. Key, and J. E. Winston, editors. *Bryozoan studies 2007*. Virginia Museum of Natural
884 History, Martinsville, VA, USA.
- 885 Rosso, A. 2009. The first catenicellid (Bryozoa, Ascophora) from Mediterranean shallow waters: a hidden resident
886 or a new immigrant? *Journal of Natural History* 43:2209–2226.
- 887 Rosso, A., and I. Di Geronimo. 1998. Deep-sea Pleistocene Bryozoa of southern Italy. *Geobios* 30:303–317.
- 888 Rosso, A., and M. Novosel. 2010. The genus *Adeonella* (Bryozoa, Ascophora) in the Mediterranean, with
889 description of two new living species and rediscovery of a fossil one. *Journal of Natural History* 44:1697–1727.
- 890 Rosso, A., and R. Sanfilippo. 2005. Bryozoans and serpulideans in skeletobiont communities from the Pleistocene
891 of Sicily: spatial utilisation and competitive interaction. *Annali dell'Università degli Studi di Ferrara, Museologia*
892 *Scientifica e Naturalistica*, Volume speciale:17 pp.
- 893 Rosso, A., F. Sciuto, and A. Sinagra. 2010. *Bertonsonidra* n. gen. (Bryozoa, Cheilostomata) for *Tremoporaprenanti*
894 *Gautier*, 1955, a rare species from the Mediterranean. *Zoosystema* 32:457–467.
- 895 Rosso, A., and P. D. Taylor. 2002. A new anaskan cheilostome bryozoan from Icelandic deep waters and its
896 uniserial colony growth pattern. *Sarsia* 87:35–46.
- 897 Rucker, J. B., and R. E. Carver. 1969. A survey of the carbonate mineralogy of cheilostome Bryozoa. *Journal of*
898 *Paleontology* 43:791–799.
- 899 Ryland, J. S. 1969. A nomenclatural index to "A History of the British Marine Polyzoa" by T. Hincks (1880).

- 900 Bulletin of the British Museum (Natural History), Zoology 17:205–260.
- 901 Ryland, J. S. 1970. Bryozoans. Hutchinson University Library, London.
- 902 Ryland, J. S. 2001. Convergent colonial organization and reproductive function in two bryozoan species epizoic on
903 gastropod shells. *Journal of Natural History* 35:1085–1101.
- 904 Ryland, J. S., J. D. D. Bishop, H. De Blauwe, A. El Nagar, D. Minchin, C. A. Wood, and A. L. E. Yunnie. 2011.
905 Alien species of *Bugula* (Bryozoa) along the Atlantic coasts of Europe. *Aquatic Invasions* 6:17–31.
- 906 Ryland, J. S., H. De Blauwe, R. Lord, and J. A. Mackie. 2009. Recent discoveries of alien *Watersipora* (Bryozoa)
907 in Western Europe, with redescrptions of species. *Zootaxa* 2093:43–59.
- 908 Ryland, J. S., and J. S. Porter. 2003. The identity of *Alcyonidium gelatinosum* (Linnaeus, 1761) (Bryozoa:
909 Ctenostomatida). *Journal of Natural History* 37:2179–2189.
- 910 Sakagami, S. 1970. On the Paleozoic Bryozoa of Japan and Thai-Malayan districts. *Journal of Paleontology*
911 44:680–692.
- 912 Sakagami, S. 1999. Permian bryozoans from some localities in the Khao Hin Kling area near Phetchabun,
913 north-central Thailand. *Bulletin of the Kitakyushu Museum of Natural History* 18:77–103.
- 914 Sakagami, S., D. Sciunnach, and E. Garzanti. 2006. Late Paleozoic and Triassic bryozoans from the Tethys
915 Himalaya (N India, Nepal and S Tibet). *Facies* 52:279–298.
- 916 Sandberg, P. 1962. New cheilostome Bryozoa from the Miocene of the Dominican Republic. *Micropaleontology*
917 8:61–66.
- 918 Santagata, S., and W. C. Banta. 1996. Origin of brooding and ovicells in cheilostome bryozoans: interpretive
919 morphology of *Scrupocellaria ferox*. *Invertebrate Biology* 115:170–180.
- 920 Santana, F. T., L. V. Ramalho, and C. P. Gumarães. 2009. A new species of *Metrarabdotos* (Bryozoa, Ascophora)
921 from Brazil. *Zootaxa* 2222.
- 922 Schmid, H. P., M. Harzhauser, and A. Kroh. 2001. Hypoxic events on a Middle Miocene carbonate platform of the
923 Central Paratethys (Austria, Badenian, 14 Ma). *Annalen des Naturhistorischen Museums in Wien* 102 A:1–50.
- 924 Schmidt, R. 2007. Australian Cenozoic Bryozoa, 2: free-living Cheilostomata of the Eocene St. Vincent Basin,
925 S.A., including *Bonellina* gen. nov. *Alcheringa* 31:67–84.
- 926 Schmidt, R., and Y. Bone. 2003. Biogeography of Eocene bryozoans from the St. Vincent Basin, South Australia.
927 *Lethaia* 36:345–356.
- 928 Schneider, S., B. Berning, M. A. Bitner, R.-P. Carriol, M. Jäger, J. Kriwet, A. Kroh, and W. Werner. 2009.
929 A parautochthonous shallow marine fauna from the Late Burdigalian (early Ottnangian) of Gurlarn (Lower
930 Bavaria, SE Germany): macrofaunal inventory and paleoecology. *Neues Jahrbuch für Geologie und Paläontologie,*
931 *Abhandlungen* 254:63–103.
- 932 Schopf, T. J. M. 1969. Paleoecology of ectoprocts (bryozoans). *Journal of Paleontology* 43:234–244.
- 933 Schopf, T. J. M., and F. T. Manheim. 1967. Chemical composition of Ectoprocta (Bryozoa). *Journal of*
934 *Paleontology* 41:1197–1225.
- 935 Sendino, C., and P. D. Taylor. 2011. Sir Charles Lyell’s fossil bryozoans from Gran Canaria. Pages 123–142 *in*
936 P. N. Wyse Jackson and M. E. Spencer Jones, editors. *Annals of Bryozoology 3: Aspects of the History of*
937 *Research on Bryozoans*. International Bryozoology Association, Dublin.
- 938 Shapiro, D. F. 1996. Size-dependent neural integration between genetically different colonies of a marine bryozoan.
939 *The Journal of Experimental Biology* 199:1229–1239.
- 940 Shier, D. E. 1964. Marine Bryozoa from northwest Florida. *Bulletin of Marine Science of the Gulf and Caribbean*
941 14:603–662.
- 942 Shunatova, N. N., and A. N. Ostrovsky. 2002. Group autozooidal behaviour and chimneys in marine bryozoans.
943 *Marine Biology* 140:503–518.
- 944 Smith, A. M. 1995. Palaeoenvironmental interpretation using bryozoans: a review. Pages 231–243 *in* D. W. J.
945 Bosence and P. A. Allison, editors. *Marine Palaeoenvironmental Analysis from Fossils*. Geological Society
946 Special Publication 83.

- 947 Smith, A. M. 2014. Growth and calcification of marine bryozoans in a changing ocean. *Biological Bulletin*
948 226:203–210.
- 949 Smith, A. M., and E. Girvan. 2010. Understanding a bimineralic bryozoan: Skeletal structure and carbonate
950 mineralogy of *Odontionella cyclops* (Foveolariidae: Cheilostomata: Bryozoa) in New Zealand. *Palaeogeography,*
951 *Palaeoclimatology, Palaeoecology* 289:113–122.
- 952 Smith, A. M., M. M. Key, and D. P. Gordon. 2006. Skeletal mineralogy of bryozoans: taxonomic and temporal
953 patterns. *Earth-Science Reviews* 78:287–306.
- 954 Smith, A. M., B. Stewart, M. M. Key, and C. M. Jamet. 2001. Growth and carbonate production by *Adeonellopsis*
955 (Bryozoa: Cheilostomata) in Doubtful Sound, New Zealand. *Palaeogeography, Palaeoclimatology, Palaeoecology*
956 175:201–210.
- 957 Sogot, C. E., E. M. Harper, and P. D. Taylor. 2014. The Lilliput effect in colonial organisms: cheilostome
958 bryozoans at the Cretaceous–Paleogene mass extinction. *Public Library of Science: ONE* 9:e87048.
- 959 Sonar, M. A., and R. M. Badve. 2008. Fossil cyclostome bryozoans from the Holocene rocks of west coast of
960 Maharashtra and Goa, India. *Gondwana Geological Magazine* 23:69–76.
- 961 Sonar, M. A., R. M. Badve, and S. G. Gaikwad. 2010. Usefulness of Bryozoa to deduce palaeoenvironment: a
962 case study from the Holocene of west coast of Maharashtra and Goa. *Gondwana Geological Magazine* 25:69–80.
- 963 Sonar, M. A., and S. G. Gaikwad. 2013a. Cyclostome Bryozoa from the Cenozoic sediments of western Kachchh,
964 Gujarat, India. *Journal of the Palaeontological Society of India* 58:195–203.
- 965 Sonar, M. A., and S. G. Gaikwad. 2013b. Fossil neocheilostomine bryozoans from the Tertiary rocks of Western
966 Kachchh, Gujarat. *Journal of the Geological Society of India* 81:665–676.
- 967 Sonar, M. A., and S. G. Gaikwad. 2013c. Fossil steginoporellid (Cheilostomata: Neocheilostomina), Bryozoa from
968 the Tertiary sediments of Western Kachchh, Gujarat, India. *Journal of Earth System Science* 122:149–161.
- 969 Souaya, F. J. 1965. On the Bryozoa of Gebel Gharra (Cairo-Suez Road) and other Miocene sections in Egypt.
970 *Journal of Paleontology* 39:1129–1144.
- 971 Soule, J. D. 1973. Histological and histochemical studies on the bryozoan-substrate interface. Pages 343–347 *in* G.
972 P. Larwood, editor. *Living and Fossil Bryozoa*. Academic Press, London.
- 973 Soule, J. D., and D. F. Soule. 1969. Systematics and biogeography of burrowing bryozoans. *American Zoologist*
974 9:791–802.
- 975 Soule, J. D., and D. F. Soule. 1970. New species of *Thalamoporella* (Ectoprocta) from Hawaii, examined by
976 scanning electron microscopy. *American Museum Novitates* 2417:1–18.
- 977 Souto, J., O. Reverter-Gil, and E. Fernández-Pulpeiro. 2010. Bryozoa from detritic bottoms in the Menorca
978 Channel (Balearic Islands, western Mediterranean), with notes on the genus *Cribellopora*. *Zootaxa* 2536:36–52.
- 979 Souto, J., O. Reverter-Gil, and E. Fernández-Pulpeiro. 2011. Redescription of some bryozoan species originally
980 described by J. Jullien from Iberian waters. *Zootaxa* 2827:31–53.
- 981 Souto, J., O. Reverter-Gil, and A. N. Ostrovsky. 2014. New species of Bryozoa from Madeira associated with
982 rhodoliths. *Zootaxa* 3795:135–151.
- 983 Stach, L. W. 1936. Studies on Recent Petraliidae (Bryozoa). *Records of the Australian Museum* 19:355–379.
- 984 Steinhorsdottir, M., S. Lidgard, and E. Håkansson. 2006. Fossils, sediments, tectonics. Reconstructing
985 palaeoenvironments in a Pliocene–Pleistocene Mediterranean microbasin. *Facies* 52:361–380.
- 986 Suárez-Andrés, J. L., and A. Ernst. 2015. Lower–Middle Devonian Fenestellidae (Bryozoa) of NW Spain:
987 implications for fenestrate palaeobiogeography. *Facies* 61:415.
- 988 Suárez-Andrés, J. L., and P. N. Wyse Jackson. 2014. *Ernstipora mackinneyi*, a new unique fenestrate bryozoan
989 genus and species with an encrusting growth habit from the Emsian (Devonian) of NW Spain. *Neues Jahrbuch*
990 *für Geologie und Paläontologie, Abhandlungen* 271:229–242.
- 991 Suárez-Andrés, J. L., and P. N. Wyse Jackson. 2015. Feeding currents: a limiting factor for disparity of Palaeozoic
992 fenestrate bryozoans. *Palaeogeography, Palaeoclimatology, Palaeoecology* 433:219–232.
- 993 Suttner, T. J., and A. Ernst. 2007. Upper Ordovician bryozoans of the Pin Formation (Spiti Valley, northern

- 994 India). *Palaeontology* 50:1485–1518.
- 995 Szczechura, J. 1994. A segmented Paleocene cheilostomatous bryozoan and its possible relationships with
996 pseudarcellids. *Acta Palaeontologica Polonica* 39:223–231.
- 997 Sørensen, A. M., E. Håkansson, and L. Stemmerik. 2007. Faunal migration into the Late Permian Zechstein Basin
998 – evidence from bryozoan palaeobiogeography. *Palaeogeography, Palaeoclimatology, Palaeoecology* 251:198–209.
- 999 Tang, S., and R. J. Cuffey. 1998. *Inconobotopora lichenoporoides* a new genus and species of cystoporate bryozoan
1000 from the Silurian of Gotland, and its evolutionary implication. *Journal of Paleontology* 72:2562–64.
- 1001 Taylor, P. D. 1978. The spiral bryozoan *Terebellaria* from the Jurassic of southern England and Normandy.
1002 *Palaeontology* 21:357–391.
- 1003 Taylor, P. D. 1982. Probable predatory borings in Late Cretaceous bryozoans. *Lethaia* 15:67–74.
- 1004 Taylor, P. D. 1986. The ancestrula and early growth pattern in two primitive cheilostome bryozoans: *Pyripora*
1005 *catenularia* (Fleming) and *Pyriporopsis portlandensis* Pohowsky. *Journal of Natural History* 20:101–110.
- 1006 Taylor, P. D. 2001. Preliminary systematics and diversity patterns of cyclostome bryozoans from the Neogene of
1007 the Central American Isthmus. *Journal of Paleontology* 75:578–589.
- 1008 Taylor, P. D. 2008. Late Cretaceous cheilostome bryozoans from California and Baja California. *Journal of*
1009 *Paleontology* 82:823–834.
- 1010 Taylor, P. D. 2009. Bryozoans from the Middle Jurassic of Balin, Poland: a revision of material described by A.E.
1011 Reuss (1867). *Annalen des Naturhistorischen Museums in Wien* 110A:17–54.
- 1012 Taylor, P. D., B. Berning, and M. A. Wilson. 2013. Reinterpretation of the Cambrian 'bryozoan' *Pywackia* as an
1013 octocoral. *Journal of Paleontology* 87:984–990.
- 1014 Taylor, P. D., and P. L. Cook. 1981. *Hippoporidra edax* (Busk 1859) and a revision of some fossil and living
1015 *Hippoporidra* (Bryozoa). *Bulletin of the British Museum (Natural History), Geology* 35:243–251.
- 1016 Taylor, P. D., and E. Di Martino. 2014. Why is the tropical Cenozoic fossil record so poor for bryozoans? *Studi*
1017 *Trentini di Scienze Naturali* 94:249–257.
- 1018 Taylor, P. D., M. H. Dick, D. Clements, and S. F. Mawatari. 2012. A diverse bryozoan fauna from Pleistocene
1019 marine gravels at Kuromatsunai, Hokkaido, Japan. Pages 367–383 *in* A. Ernst, P. Schäfer, and J. Scholz,
1020 editors. *Bryozoan Studies 2010*. Springer, Berlin.
- 1021 Taylor, P. D., and A. Ernst. 2008. Bryozoans in transition: the depauperate and patchy Jurassic biota.
1022 *Palaeogeography, Palaeoclimatology, Palaeoecology* 263:9–23.
- 1023 Taylor, P. D., and D. P. Gordon. 1997. *Fenestulipora*, gen. nov., an unusual cyclostome bryozoan from New
1024 Zealand and Indonesia. *Invertebrate Taxonomy* 11:689–703.
- 1025 Taylor, P. D., and D. P. Gordon. 2002. Alcide d'Orbigny's work on Recent and fossil bryozoans. *Comptes Rendus*
1026 *Palevol* 1:533–547.
- 1027 Taylor, P. D., and D. P. Gordon. 2003. Endemic new cyclostome bryozoans from Spirits Bay, a New Zealand
1028 marine-biodiversity "hotspot". *New Zealand Journal of Marine and Freshwater Research* 37:653–669.
- 1029 Taylor, P. D., D. P. Gordon, and P. B. Batson. 2004. Bathymetric distributions of modern populations of some
1030 common Cenozoic Bryozoa from New Zealand, and paleodepth estimation. *New Zealand Journal of Geology*
1031 *and Geophysics* 47:57–69.
- 1032 Taylor, P. D., and N. P. James. 2013. Secular changes in colony-forms and bryozoan carbonate sediments through
1033 geological history. *Sedimentology* 60:1184–1212.
- 1034 Taylor, P. D., N. P. James, Y. Bone, P. Kuklinski, and T. K. Kyser. 2009a. Evolving mineralogy of cheilostome
1035 bryozoans. *Palaios* 24:440–452.
- 1036 Taylor, P. D., P. Kuklinski, and D. P. Gordon. 2007. Branch diameter and depositional depth in cyclostome
1037 bryozoans: testing a potential paleobathymetric tool. *Palaios* 22:220–224.
- 1038 Taylor, P. D., D. G. Lazo, and M. B. Aguirre-Ureta. 2009b. Lower Cretaceous bryozoans from Argentina: a
1039 'by-catch' fauna from the Agrío Formation (Neuquén Basin). *Cretaceous Research* 30:193–203.
- 1040 Taylor, P. D., and F. K. McKinney. 1996. An Archimedes-like cyclostome bryozoan from the Eocene of North

- 1041 Carolina. *Journal of Paleontology* 70:218–229.
- 1042 Taylor, P. D., and F. K. McKinney. 2000. Reinterpretation of *Stictostega* Shaw, 1967, an Upper Cretaceous
1043 cheilostome bryozoan from Arkansas. *Journal of Paleontology* 74:1–6.
- 1044 Taylor, P. D., and K. S. Schindler. 2004. A new Eocene species of the hermit-crab symbiont *Hippoporidra*
1045 (Bryozoa) from the Ocala Limestone of Florida. *Journal of Paleontology* 78:790–794.
- 1046 Taylor, P. D., and C. Sendino. 2010. Latitudinal distribution of bryozoan-rich sediments in the Ordovician.
1047 *Bulletin of Geosciences* 85:565–572.
- 1048 Taylor, P. D., O. Vinn, and M. A. Wilson. 2010. Evolution of biomineralization in "lophophorates". *Palaeontology*
1049 84:317–333.
- 1050 Taylor, P. D., and A. Waeschenbach. 2015. Phylogeny and diversification of bryozoans. *Palaeontology* 58:585–599.
- 1051 Taylor, P. D., A. Waeschenbach, and W. K. Florence. 2011. Phylogenetic position and systematics of the bryozoan
1052 *Tennysonia*: further evidence for convergence and plasticity in skeletal morphology among cyclostome bryozoans.
1053 *Zootaxa* 3010:58–68.
- 1054 Taylor, P. D., and M. A. Wilson. 2002. A new terminology for marine organisms inhabiting hard substrates.
1055 *Palaios* 17:522–525.
- 1056 Taylor, P. D., and M. A. Wilson. 2003. Palaeoecology and evolution of marine hard substrate communities.
1057 *Earth-Science Reviews* 62:1–103.
- 1058 Taylor, P. D., M. A. Wilson, and R. G. Bromley. 1999. A new ichnogenus for etchings made by cheilostome
1059 bryozoans into calcareous substrates. *Palaeontology* 42:595–604.
- 1060 Taylor, P. D., and P. M. Zaborski. 2002. A Late Cenomanian bryozoan biostrome from north-eastern Nigeria.
1061 *Cretaceous Research* 23:241–253.
- 1062 Taylor, P. D., and M. Zaton. 2008. Taxonomy of the bryozoan genera *Oncousoecia*, *Microeciella* and *Eurystroto*
1063 (Cyclostomata: Oncousoeciidae). *Journal of Natural History* 42:2557–2574.
- 1064 Taylor, P. D., and K. Zágorský. 2011. Operculate cyclostome bryozoans (Eleidae) from the Bohemian Cretaceous.
1065 *Paläontologische Zeitschrift* 85:407–432.
- 1066 Tenison-Woods, J. E. 1880. *Palaeontology of New Zealand. Part IV. Corals and Bryozoa of the Neozoic Period in*
1067 *New Zealand. Page 34. G. Didsbury, Government Printer, Wellington.*
- 1068 Thomson, W. T. C. 1852. Notes on some Scotch Zoophytes and Polyzoa. *Annals and Magazine of Natural History,*
1069 *series 2* 9:403–404.
- 1070 Tilbrook, K. J. 2001. Indo-West Pacific species of the genus *Stylopoma* Levinsen, 1909 (Bryozoa: Cheilostatida).
1071 *Zoological Journal of the Linnean Society* 131:1–34.
- 1072 Tilbrook, K. J. 2011. New genus for a unique species of Indo-West Pacific bryozoan. *Zootaxa* 3134:63–67.
- 1073 Tilbrook, K. J., and D. P. Gordon. 2015. Bryozoa from the Straits of Johor, Singapore, with the description of a
1074 new species. *Raffles Bulletin of Zoology* 31:255–263.
- 1075 Tilbrook, K. J., and A. V. Grischenko. 2004. New sub-Arctic species of the tropical genus *Antropora* (Bryozoa:
1076 Cheilostomata): a gastropod-pagurid crab associate. *Journal of the Marine Biological Association of the United*
1077 *Kingdom* 84:1001–1004.
- 1078 Tilbrook, K. J., P. J. Hayward, and D. P. Gordon. 2001. Cheilostomatous Bryozoa from Vanuatu. *Zoological*
1079 *Journal of the Linnean Society* 131:35–109.
- 1080 Tolokonnikova, Z. A., and A. Ernst. 2010. Palaeobiogeography of Famennian (Late Devonian) bryozoans.
1081 *Palaeogeography, Palaeoclimatology, Palaeoecology* 298:360–369.
- 1082 Tompsett, S., J. S. Porter, and P. D. Taylor. 2009. Taxonomy of the fouling cheilostome bryozoans *Schizoporella*
1083 *unicornis* (Johnston) and *Schizoporella errata* (Waters). *Journal of Natural History* 43:2227–2243.
- 1084 Toots, H., and J. F. Cutler. 1962. Bryozoa from the "Mesaverde" Formation (Upper Cretaceous) of southeastern
1085 Wyoming. *Journal of Paleontology* 36:81–86.
- 1086 Toscano, F., and P. D. Taylor. 2008. A new Early Miocene bryozoan, *Favosipora ichnusae* sp. nov. (Cyclostomata),
1087 from the Isili Limestone of Sardinia, Italy. *Neues Jahrbuch für Geologie und Paläontologie Abhandlungen*

- 1088 248:301–308.
- 1089 Tuckey, M. E. 1990. Distributions and extinctions of Silurian Bryozoa. Pages 197–206 *in* W. S. McKerrow and C.
1090 R. Scotese, editors. *Palaeozoic Palaeogeography and Biogeography*. Geological Society, London.
- 1091 Urbanek, A. 2004. Morphogenetic gradients in graptolites and bryozoans. *Acta Palaeontologica Polonica*
1092 49:485–504.
- 1093 Utgaard, J. 1981. *Lunaferamita*, a new genus of Constellariidae (Bryozoa) with strong cystoporate affinities.
1094 *Journal of Paleontology* 55:1058–1070.
- 1095 Vávra, N. 1980. Tropische Faunenelemente in den Bryozoenfaunen des Badenien (Mittelmiozän) der Zen-
1096 tralen Paratethys. *Sitzungsberichte der Österreichischen Akademie der Wissenschaften, Mathematisch-*
1097 *naturwissenschaftliche Klasse, Abteilung 1* 189:49–63.
- 1098 Vieira, L. M., and D. P. Gordon. 2010. *Eutaleola*, a replacement name for the homonym *Euteleia* (Bryozoa:
1099 Pasytheidae). *Zoologia* 27:646–648.
- 1100 Vieira, L. M., D. P. Gordon, and M. D. Correia. 2007. First record of a living ditaxiporine catenicellid in the
1101 Atlantic, with a description of *Vasignyella oviceolata* n. sp. (Bryozoa). *Zootaxa* 1582:49–58.
- 1102 Vieira, L. M., D. P. Gordon, F. B. C. Souza, and M. A. Haddad. 2010. New and little-known cheilostomatous
1103 Bryozoa from the south and southeastern Brazilian continental shelf and slope. *Zootaxa* 2722:1–53.
- 1104 Vieira, L. M., M. E. Spencer Jones, and J. E. Winston. 2013. *Cradoscrupocellaria*, a new bryozoan genus for
1105 *Scrupocellaria bertholletii* (Audouin) and related species (Cheilostomata, Candidae): taxonomy, biodiversity
1106 and distribution. *Zootaxa* 3707:1–63.
- 1107 Vieira, L. M., M. E. Spencer Jones, J. E. Winston, A. E. Migotto, and A. C. Marques. 2014. Evidence for
1108 polyphyly of the genus *Scrupocellaria* (Bryozoa: Candidae) based on a phylogenetic analysis of morphological
1109 characters. *Public Library of Science: ONE* 9:e95296.
- 1110 Vieira, L. M., and S. N. Stampar. 2014. A new *Fenestulina* (Bryozoa, Cheilostomata) commensal with
1111 tube-dwelling anemones (Cnidaria, Ceriantharia) in the tropical southwestern Atlantic. *Zootaxa* 3780:365–374.
- 1112 Vine, G. R. 1884. Polyzoa (bryozoa) found in the boring at richmond, surrey referred to by prof. Judd, frs.
1113 *Quarterly Journal of the Geological Society* 40:784–794.
- 1114 Viskova, L. A., and A. V. Koromyslova. 2012. *Tamanicella* gen. nov., a new genus of bryozoans forming the Late
1115 Miocene bioherms of Cape Panagia in the Taman Peninsula (Russia). *Paleontological Journal* 46:29–43.
- 1116 Voigt, E. 1964. A bryozoan fauna of Dano-Montian age from Boryszew and Sochaczew in Central Poland. *Acta*
1117 *Palaeontologica Polonica* 9:419–498.
- 1118 Voigt, E. 1968. On the Cretaceous age of the so-called Jurassic cheilostomatous Polyzoa (Bryozoa). A contribution
1119 to the knowledge of the Polyzoa-fauna of the Maastrichtian in the Cotentin (Manche). *Bulletin of the British*
1120 *Museum (Natural History), Geology* 17:1–45.
- 1121 Waeschenbach, A., J. S. Porter, and R. N. Hughes. 2012. Molecular variability in the *Celleporella hyalina*
1122 (Bryozoa; Cheilostomata) species complex: evidence for cryptic speciation from complete mitochondrial
1123 genomes. *Molecular Biology Reports* 39:8601–8614.
- 1124 Wass, R. E. 1972. Permian Bryozoa from South Africa. *Journal of Paleontology* 46:871–873.
- 1125 Wass, R. E. 1977. Branching patterns and phylogeny of the family Vittaticellidae (Bryozoa: Cheilostomata).
1126 *Australian Journal of Zoology* 25:103–119.
- 1127 Wass, R. E., and J. J. Yoo. 1975. Bryozoa from Site 282 West Tasmania. Pages 809–831 *in* J. P. Lemmet, R.
1128 E. Houtz, and E. al., editors. *Initial Reports of the Deep Sea Drilling Project, Volume 29*. U.S Government
1129 Printing Office, Washington, USA.
- 1130 Wass, R. E., and J. J. Yoo. 1983. Cheilostome Bryozoa from the southern Australian continental shelf. *Australian*
1131 *Journal of Marine and Freshwater Research* 34:303–354.
- 1132 Waters, A. W. 1881. On fossil chilostomatous Bryozoa from south-west Victoria, Australia. *Quarterly Journal of*
1133 *the Geological Society of London* 37:309–347.
- 1134 Waters, A. W. 1882. On chilostomatous Bryozoa from Bairnsdale (Gippsland). *Quarterly Journal of the Geological*

- 1135 Society of London 38:502–513.
- 1136 Waters, A. W. 1883. On fossil chilostomatous Bryozoa from Muddy Creek, Victoria. Quarterly Journal of the
1137 Geological Society of London 39:423–443.
- 1138 Waters, A. W. 1884a. Fossil cyclostomatous Bryozoa from Australia. Quarterly Journal of the Geological Society
1139 of London 40:674–697.
- 1140 Waters, A. W. 1884b. Closure of the cyclostomatous Bryozoa. Journal of the Linnean Society of London, Zoology
1141 17:400–404.
- 1142 Waters, A. W. 1885. Chilostomatous Bryozoa from Aldinga and the River-Murray Cliffs, South Australia.
1143 Quarterly Journal of the Geological Society of London 41:279–310.
- 1144 Waters, A. W. 1888. Supplementary Report on the Polyzoa collected by H.M.S. 'Challenger' during the years
1145 1873-1876. Report on the Scientific Results of the Voyage of the H.M.S. 'Challenger'. Zoology Part LXXIX
1146 31:1–41.
- 1147 Waters, A. W. 1889. Bryozoa from New South Wales. Part IV. Annals and Magazine of Natural History, series 6
1148 4:1–24.
- 1149 Waters, A. W. 1898. Observations on Membraniporidae. Journal of the Linnean Society (Zoology) 26:654–693.
- 1150 Waters, A. W. 1904a. Bryozoa. Résultats du Voyage du S.Y. Belgica en 1897-1898-1899, Zoologie 4:1–114.
- 1151 Waters, A. W. 1904b. Bryozoa from Franz-Josef Land, collected by the Jackson-Harmsworth expedition, 1896-1897.
1152 Part 2, Cyclostomata, Ctenostomata, and Entoprocta. Journal of the Linnean Society (Zoology) 29:161–184.
- 1153 Waters, A. W. 1905. Bryozoa from near Cape Horn. Journal of the Linnean Society (Zoology) 29:230–251.
- 1154 Waters, A. W. 1907. *Tubucellaria*: its species and ovicells. Journal of the Linnean Society (Zoology) 30:126–133.
- 1155 Weedon, M. J., and P. D. Taylor. 1995. Calcitic nacreous ultrastructures in bryozoans: implications for comparative
1156 biomineralization of lophophorates and molluscs. Biological Bulletin 188:281–292.
- 1157 Weedon, M. J., and P. D. Taylor. 1996. Skeletal ultrastructures in some cerioporine cyclostome bryozoans. Acta
1158 Zoologica 77:249–265.
- 1159 West, R. R., F. K. McKinney, J. A. Fagerstrom, and J. Vacelet. 2011. Biological interactions among extant and
1160 fossil clonal organisms. Facies 57:351–374.
- 1161 Williams, J. D., and J. J. McDermott. 2004. Hermit crab biocoenoses: a worldwide review of the diversity and
1162 natural history of hermit crab associates. Journal of Experimental Marine Biology and Ecology 305:1–128.
- 1163 Williams, M., A. M. Haywood, E. M. Harper, A. L. A. Johnson, T. Knowles, M. J. Leng, D. J. Lunt, B. Okamura, P.
1164 D. Taylor, and J. Zalasiewicz. 2009. Pliocene climate and seasonality in North Atlantic shelf seas. Philosophical
1165 Transactions of the Royal Society of London, Ser. A 367:85–108.
- 1166 Wilson, M. A., and P. D. Taylor. 2001. Palaeoecology of hard substrate faunas from the Cretaceous Qahlah
1167 Formation of the Oman Mountains. Palaeontology 44:21–41.
- 1168 Wilson, M. A., and P. D. Taylor. 2006. Predatory drill holes and partial mortality in Devonian colonial metazoans.
1169 Geology 34:565–568.
- 1170 Winston, J. E. 1977. Feeding in marine bryozoans. Pages 233–271 in R. M. Woollacott and R. L. Zimmer, editors.
1171 Biology of Bryozoans. Academic Press, New York.
- 1172 Winston, J. E., and S. E. Beaulieu. 1999. *Striatodoma dorothea* (Cheilostatida: Tessaradomidae), a new genus
1173 and species of bryozoan from deep water off California. Proceedings of the Biological Society of Washington
1174 112:313–318.
- 1175 Winston, J. E., and B. F. Heimberg. 1986. Bryozoans from Bali, Lombok, and Komodo. American Museum
1176 Novitates 2847:1–49.
- 1177 Winston, J. E., and J. B. C. Jackson. 1984. Ecology of cryptic coral reef communities. IV. Community development
1178 and life histories of encrusting cheilostome Bryozoa. Journal of Experimental Marine Biology and Ecology
1179 76:1–21.
- 1180 Winston, J. E., and L. M. Vieira. 2013. Systematics of interstitial encrusting bryozoans from southeastern Brazil.
1181 Zootaxa 3710:101–146.

- 1182 Winston, J. E., L. M. Vieira, and R. M. Woollacott. 2014. Scientific results of the Hassler Expedition. Bryozoa No.
1183 2. Brazil. *Bulletin of the Museum of Comparative Zoology* 161:139–239.
- 1184 Winston, J. E., and R. M. Woollacott. 2009. Scientific results of the Hassler Expedition. Bryozoa. No. 1 Barbados.
1185 *Bulletin of the Museum of Comparative Zoology* 159:239–300.
- 1186 Wood, A. C. L., P. K. Probert, A. A. Rowden, and A. M. Smith. 2012. Complex habitat generated by marine
1187 bryozoans: a review of its distribution, structure, diversity, threats and conservation. *Aquatic Conservation:*
1188 *Marine and Freshwater Ecosystems* 22:547–563.
- 1189 Wyse Jackson, P. N., and C. J. Buttler. 2015. Part G, Revised, Volume 2, Chapter 3: Preparation, imaging, and
1190 conservation of Paleozoic bryozoans for study. *Treatise Online* 63:1–15.
- 1191 Wyse Jackson, P. N., E. Cross, and R. Swift. 2009. *Ptilofenestella*, a distinctive rare fenestrate bryozoan from the
1192 Mississippian of Ireland: new records and an extension of its stratigraphical range. *Irish Naturalist's Journal*
1193 30:76–77.
- 1194 Wyse Jackson, P. N., and M. M. Key. 2007. Borings in trepostome bryozoans from the Ordovician of Estonia:
1195 two ichnogenera produced by a single maker, a case of host morphology control. *Lethaia* 40:237–252.
- 1196 Wyse Jackson, P. N., and M. M. Key. 2014. Epizoic bryozoans on cephalopods through the Phanerozoic: a review.
1197 *Studi Trentini di Scienze Naturali* 94:283–291.
- 1198 Wyse Jackson, P. N., and F. K. McKinney. 2013. A micro-computed tomography and scanning electron microscopy
1199 investigation of the structure of *Polyfenestella* Bancroft, 1986 (Bryozoa: Fenestrata), from the Mississippian of
1200 Scotland: revealing the nature of its heteromorphs. *Irish Journal of Earth Sciences* 31:1–6.
- 1201 Wyse Jackson, P. N., F. K. McKinney, and A. Ernst. 2010. *Retepora undata* M'Coy, 1844, type species of
1202 *Phylloporella* Frederiks, 1916, a junior synonym of *Chainodictyon* Foerste, 1877: a phylloporinid bryozoan from
1203 the Mississippian of Ireland redescribed. *Irish Journal of Earth Sciences* 28:53–60.
- 1204 Wyse Jackson, P. N., C. M. Reid, and F. K. McKinney. 2011. The status of *Protoretepora* de Koninck, 1878
1205 (Fenestrata: Bryozoa), and description of *P. crockfordae* sp. nov. and *P. wassi* sp. nov. from the Permian of
1206 Australia. *Alcheringa* 35:539–552.
- 1207 Wyse Jackson, P. N., C. M. Reid, and F. K. McKinney. 2012. Fixation of the type species of the genus *Protoretepora*
1208 de Koninck, 1878 (Bryozoa, Fenestrata). *Alcheringa: An Australasian Journal of Palaeontology* 36:137–138.
- 1209 Xia, F.-S., S.-G. Zhang, and Z.-Z. Wang. 2007. The oldest bryozoans: new evidence from the Late Tremadocian
1210 (Early Ordovician) of East Yangtze Gorges in China. *Journal of Paleontology* 81:1308–1326.
- 1211 Yagunova, E. B., and A. N. Ostrovsky. 2008. Encrusting bryozoan colonies on stones and algae: variability of
1212 zooidal size and its possible causes. *Journal of the Marine Biological Association of the United Kingdom*
1213 88:901–908.
- 1214 Zabala, M., and P. Maluquer. 1988. *Illustrated Keys for the Classification of Mediterranean Bryozoa*. Ajuntament
1215 de Barcelona, Barcelona.
- 1216 Zaton, M., and P. D. Taylor. 2009. Middle Jurassic cyclostome bryozoans from the Polish Jura. *Acta Palaeontolo-*
1217 *gica Polonica* 54:267–288.
- 1218 Zágorsek, K. 2003. Upper Eocene Bryozoa from Waschberg Zone (Austria). *Beiträge zur Paläontologie* 28:101–263.
- 1219 Zágorsek, K. 2010. Bryozoa from the Langhian (Miocene) of the Czech Republic. Part II: Systematic description
1220 of the suborder Ascophora Levinsen, 1909 and paleoecological reconstruction of the studied paleoenvironment.
1221 *Acta Musei Nationalis Pragae, Series B - Historia Naturalis* 66:139–255.
- 1222 Zágorsek, K., S. Filipescu, and K. Holcová. 2010. New Middle Miocene Bryozoa from Gârbova de Sus (Romania)
1223 and their relationship to the sedimentary environment. *Geologica Carpathica* 61:495–512.
- 1224 Zágorsek, K., and K. Fordinál. 2006. Lower Sarmatian Bryozoa from brackish sediment in the northern part of
1225 the Danube Basin (Dubová, Slovakia). *Linzer biologische Beiträge* 38:93–99.
- 1226 Zágorsek, K., and D. P. Gordon. 2013. Late Tortonian bryozoans from Mut Basin, Central Anatolian Plateau,
1227 southern Turkey. *Acta Palaeontologica Polonica* 58:595–607.
- 1228 Zágorsek, K., and D. P. Gordon. 2014. Revision of the Oligocene bryozoan taxa described by Stoliczka (1862),
1229 with the description of a new genus of Bryocryptellidae. *Geodiversitas* 36:541–564.

- 1230 Zágorsek, K., K. Holcová, S. Nehyba, A. Kroh, and S. Hladilová. 2009. The invertebrate fauna of the Middle
1231 Miocene (lower Badenian) sediments of Kralice nad Oslavou (Central Paratethys, Moravian part of the
1232 Carpathian Foredeep). *Bulletin of Geosciences* 84:465–496.
- 1233 Zágorsek, K., A. N. Ostrovsky, and N. Vávra. 2011. The new cheilostome bryozoan *Metrarabdotos nehybai* from
1234 the Middle Miocene of Moravia (Czech Republic): palaeofaunistic, taxonomic and ontogenetic aspects. *Neues
1235 Jahrbuch für Geologie und Paläontologie, Abhandlungen* 260:21–31.
- 1236 Zágorsek, K., U. Radwanska, and A. Radwanski. 2012. Bryozoa from the Korytnica Basin (Middle Miocene; Holy
1237 Cross Mountains, Central Poland). *Bulletin of Geosciences* 87:201–218.
- 1238 Zágorsek, K., L. V. Ramalho, B. Berning, and V. A. Távora. 2014. A new genus of the family Jaculinidae
1239 (Cheilostomata, Bryozoa) from the Miocene of the tropical western Atlantic. *Zootaxa* 3838:98–112.
- 1240 Zágorsek, K., L. Silye, and B. Szabó. 2008. New Bryozoa from the Sarmatian (Middle Miocene) deposits of the
1241 Cerna-Strei Depression Romania. *Studia Universitatis Babes-Bolyai, Geologia* 53:25–29.
- 1242 Zágorsek, K., N. Vávra, and K. Holcová. 2007. New and unusual Bryozoa from the Badenian (Middle Miocene) of
1243 the Moravian part of the Vienna Basin (Central Paratethys, Czech Republic). *Neues Jahrbuch für Geologie
1244 und Paläontologie, Abhandlungen* 243:201–215.
- 1245 Zágoršek, K., D. P. Gordon, and N. Vávra. 2015. Revision of Chlidoniopsidae Harmer, 1957 (Bryozoa: Cheilosto-
1246 mata) including a description of *Celiopsis vici* gen. and sp. nov. *Journal of Paleontology* 89:140–147.
- 1247 Zenetos, A., M. E. Çınar, M. A. Pancucci-Papadopoulou, J.-G. Harmelin, G. Furnari, F. Andaloro., N. Bellou, N.
1248 Streftaris, and H. Zibrowius. 2005. Annotated list of marine alien species in the Mediterranean with records of
1249 the worst invasive species. *Mediterranean Marine Science* 6:63–118.
- 1250 Zenetos, A., S. Gofas, C. Morri, A. Rosso, and et Al. 2012. Alien species in the Mediterranean Sea by 2012. A
1251 contribution to the application of European Union’s Marine Strategy Framework Directive (MSFD). Part 2.
1252 Introduction trends and pathways. *Mediterranean Marine Science* 13:328–352.
- 1253 Zhang, S.-G., F.-S. Xia, H.-J. Yan, and Z.-Z. Wang. 2009. Horizon of the oldest known bryozoans (Ordovician).
1254 *Palaeoworld* 18:67–73.
- 1255 Ziko, A., Y. A. El Safori, A. El Sorogy, M. Abd El-Wahab, N. El Dera, and W. Sheata. 2012. Bryozoa from
1256 northern Red Sea, Egypt: 1 *Crisia* (Cyclostomata). *Historical Biology* 24:113–119.