

Interface Focus

A model for the optimization of anti-inflammatory treatment with Chemerin

Simao Laranjeira, Daniel Regan-Komito, Asif J. Iqbal, David R. Greaves, Stephen J. Payne¹, Piotr Orlowski

S2 Table: Measurement of the mean and standard deviation of the number of neutrophils and monocytes at different times during the zymosan induced peritonitis experiment.

Time [hours]	Neutrophil [cell $\times 10^6$]	Monocytes [cell $\times 10^6$]
0	$1.08 \times 10^{-1} \pm 1.35 \times 10^{-1}$	$7.73 \times 10^{-2} \pm 9.39 \times 10^{-2}$
2	5.13 ± 1.3	$2.5 \times 10^{-2} \pm 7.92 \times 10^{-3}$
4	7.09 ± 2.68	$4.91 \times 10^{-1} \pm 1.66 \times 10^{-1}$
8	$7.28 \pm 5.27 \times 10^{-1}$	$3.26 \pm 8.86 \times 10^{-1}$
16	5.51 ± 1.59	$2.81 \pm 6.8 \times 10^{-3}$
20	3.73 ± 1.99	2.56 ± 1.12
48	$8.12 \times 10^{-1} \pm 6.29 \times 10^{-1}$	$7.06 \times 10^{-1} \pm 1.46 \times 10^{-1}$
72	$1.06 \times 10^{-1} \pm 8.56 \times 10^{-2}$	$1.13 \pm 3.56 \times 10^{-1}$
96	$2.29 \times 10^{-1} \pm 4.06 \times 10^{-2}$	$1.13 \pm 3.46 \times 10^{-1}$