

**Table S3:** Pre-Post analyses: unadjusted means and standard errors for outcomes at all three study visits for CATCH study participants.

<b>Outcome Name, Instrument Acronym (value range)<sup>1</sup></b>	<b>Baseline</b>	<b>3-months</b>	<b>6-months</b>
Physical component score, SF36-PCS (0 to 100)	47.4 (0.7)	47.8 (0.8)	48.9 (0.8)
Mental component scores, SF36-MCS (0 to 100)	29.0 (0.9)	33.5 (0.9)	36.0 (1.0)
Mental health symptoms, CSI (14 to 70)	42.0 (0.7)	-	34.7 (0.8)
Quality of life, QOLI20 total score (20 to 140)	71.4 (1.5)	-	81.6 (1.7)
Alcohol component, ASI ACOMP (0 to 1)	0.19 (0.02)	-	0.15 (0.01)
Drug component, ASI DCOMP (0 to 1)	0.08 (0.01)	-	0.05 (0.01)
Number of hospital admissions in past 3 months	0.58 (0.04)	0.42 (0.06)	0.35 (0.06)
Number of ED Visits in past 3 months	2.07 (0.21)	2.00 (0.28)	1.71 (0.31)
Number of days homeless in past 3 months	52.11 (2.21)	54.73 (2.45)	51.01 (2.98)

<sup>1</sup> Value ranges are only available for standardized instruments. Higher values are associated with better outcomes for SF36-PCS, SF36-MCS and QOLI-20 total score, and with poorer outcomes for the ASI and the CSI..