

## Supplemental Data

**Table 1.** Reasons for exclusion in this meta-analysis

Reasons for exclusion	Study (year)
Study protocol (n=2)	Dobourg (2013) [1], Koziarz (2017) [2]
No extractable data (n=7)	Karakitsos (2006)[3], Girisgin (2007)[4], Skoloudik (2011)[5], Shirodkar(2014)[6] Aduayi (2015) [7], Komut (2016) [8], Tarzamni (2016) [9]
Absence of interest outcomes (n=2)	Salahuddin (2016) [10], Kazdal (2016) [11]

## References

1. Dubourg J, Messerer M, Karakitsos D, Rajajee V, Antonsen E, Javouhey E, et al. Individual patient data systematic review and meta-analysis of optic nerve sheath diameter ultrasonography for detecting raised intracranial pressure: Protocol of the onsd research group. *Syst Rev* 2013;2:62.
2. Koziarz A, Sne N, Kegel F, Alhazzani W, Nath S, Badhiwala JH, et al. Optic nerve sheath diameter sonography for the diagnosis of increased intracranial pressure: A systematic review and meta-analysis protocol. *BMJ Open* 2017;7:e016194.
3. Karakitsos D, Soldatos T, Gouliamos A, Armaganidis A, Poularas J, Kalogeromitros A, et al. Transorbital sonographic monitoring of optic nerve diameter in patients with severe brain injury. *Transplant Proc* 2006;38:3700-6.
4. Girisgin AS, Kalkan E, Kocak S, Cander B, Gul M, Semiz M. The role of optic nerve ultrasonography in the diagnosis of elevated intracranial pressure. *Emerg Med J* 2007;24:251-4.
5. Skoloudik D, Herzig R, Fadrna T, Bar M, Hradilek P, Roubec M, et al. Distal enlargement of the optic nerve sheath in the hyperacute stage of intracerebral haemorrhage. *Br J Ophthalmol* 2011;95:217-21.
6. Shirodkar CG, Rao SM, Mutkule DP, Harde YR, Venkategowda PM, Mahesh MU. Optic nerve sheath diameter as a marker for evaluation and prognostication of intracranial pressure in indian patients: An observational study. *Indian J Crit Care Med* 2014;18:728-34.
7. Aduayi OS, Asaleye CM, Adetiloye VA, Komolafe EO, Aduayi VA. Optic nerve sonography: A noninvasive means of detecting raised intracranial pressure in a resource-limited setting. *J Neurosci Rural*

Pract 2015;6:563-7.

8. Komut E, Kozaci N, Sonmez BM, Yilmaz F, Komut S, Yildirim ZN, et al. Bedside sonographic measurement of optic nerve sheath diameter as a predictor of intracranial pressure in ed. Am J Emerg Med 2016;34:963-7.
9. Tarzamni MK, Derakhshan B, Meshkini A, Merat H, Fouladi DF, Mostafazadeh S, et al. The diagnostic performance of ultrasonographic optic nerve sheath diameter and color doppler indices of the ophthalmic arteries in detecting elevated intracranial pressure. Clin Neurol Neurosurg 2016;141:82-8.
10. Salahuddin N, Mohamed A, Alharbi N, Ansari H, Zaza KJ, Marashly Q, et al. The incidence of increased icp in icu patients with non-traumatic coma as diagnosed by onsd and ct: A prospective cohort study. BMC Anesthesiol 2016;16:106.
11. Kazdal H, Kanat A, Findik H, Sen A, Ozdemir B, Baticik OE, et al. Transorbital ultrasonographic measurement of optic nerve sheath diameter for intracranial midline shift in patients with head trauma. World Neurosurg 2016;85:292-7.

**Figure 1.** Forest plot of diagnostic odds ratio (DOR) of optic nerve sheath diameter (cutoff > 5.0 mm) for detecting increased intracranial pressure seen on computed tomography in adults. Pooled DOR was 177.55 (95% confidence interval: 52.66–598.60) without heterogeneity.

