

Supplementary Table: List of variables used in the current study for the Frailty Index

Changes in everyday activities	Tiredness all the time	Seizures, partial complex
Facial bradykinesia	Clinical impression of depression	Seizures, generalized
Problems getting dressed	Sleep changes	Syncope or blackouts
Urinary incontinence	Restlessness	Breast problems
Toileting problems	Memory changes	Cerebrovascular problems
Gastrointestinal problems	Short-term memory impairment	History of stroke
Problems cooking	Long-term memory impairment	History of diabetes mellitus
Problems going out alone	Changes in general cognitive functioning	Arterial hypertension
Impaired mobility	Onset of cognitive symptoms	Peripheral pulses
Falls	Delirium	Cardiac problems
Musculoskeletal problems	Paranoid features	Myocardial infarction
Bradykinesia of the limbs	Cognitive impairment history	Arrhythmia
Poor muscle tone in limbs	Cognitive impairment family history	Congestive heart failure
Poor limb coordination	Impaired vibration	Lung problems
Poor coordination, trunk	Tremor at rest	Respiratory problems
Poor standing posture	Postural tremor	History of thyroid disease
Irregular gait pattern	Intention tremor	Thyroid problems
Mood problems	History of Parkinson's disease	Malignant disease
Feeling sad, depressed	Family history of degenerative disease	Presence of snout reflex
History of depressed mood	Swallowing problems	Presence of the palmomental reflex
		Other medical history