

Table 5. Glucagon-like peptide-1 (GLP-1; pmol/L)

	100-300μ	300-500μ	Sham	100-300μ vs. 300-500μ (<i>p</i>)	100-300μ vs. Sham (<i>p</i>)	300-500μ vs. Sham (<i>p</i>)
T0	36.61 \pm 21.89	42.56 \pm 58.23	42.75 \pm 60.15	0.43	0.44	0.49
T1	41.27 \pm 20.64	65.25 \pm 54.63	63.35 \pm 42.62	0.19	0.22	0.48
T2	34.57 \pm 30.79	20.5 \pm 12.47	66.97 \pm 91.25	0.14	0.35	0.24
T3	41.03 \pm 41.77	14.40 \pm 12.19	45.92 \pm 49.19	0.17	0.49	0.17
T4	14.72 \pm 1.88	19.06 \pm 9.28	41.88 \pm 33.56	0.15	0.3	0.22
T0 vs. T1 (<i>p</i>)	0.38	0.29	0.29			
T0 vs. T2 (<i>p</i>)	0.45	0.24	0.34			
T0 vs. T3 (<i>p</i>)	0.42	0.19	0.47			
T0 vs. T4 (<i>p</i>)	0.12	0.22	0.49			
T2 vs. T3 (<i>p</i>)	0.4	0.25	0.37			
T2 vs. T4 (<i>p</i>)	0.21	0.42	0.33			
T3 vs. T4 (<i>p</i>)	0.22	0.28	0.45			

GLP-1 levels are expressed in pmol/L. T0= before embolization; T1= after 1 hour; T2= after 1 month; T3= after 3 months; T4= after 6 months. Data are expressed as mean \pm SD (standard deviation). A *p* value <0.05 was considered statistically significant.