

Table 6. Peptide YY (pg/mL)

	100-300μ	300-500μ	Sham	100-300μ vs. 300-500μ (p)	100-300μ vs. Sham (p)	300-500μ vs. Sham (p)
T0	552.11±266.76	479.61±180.66	501.72±39.56	0.33	0.37	0.47
T1	435.68±32.059	388.47±242.95	463.51±75.85	0.48	0.21	0.22
T2	556.39±174.27	506.53±194.02	417.29±59.79	0.35	0.09	0.09
T3	445.89±205.77	361.67±150.60	422.20±58.42	0.2	0.24	0.38
T4	380.30±150.86	313.91±266	513.46±140.29	0.36	0.19	0.17
T0 vs. T1 (p)	0.33	0.1	0.33			
T0 vs. T2 (p)	0.3	0.38	0.02			
T0 vs. T3 (p)	0.17	0.02	0.05			
T0 vs. T4 (p)	0.25	0.23	0.45			
T2 vs. T3 (p)	0.13	0.037	0.39			
T2 vs. T4 (p)	0.23	0.13	0.24			
T3 vs. T4 (p)	0.35	0.39	0.24			

PYY levels are expressed in pmol/L. T0= before embolization; T1= after 1 hour; T2= after 1 month; T3= after 3 months; T4= after 6 months. Data are expressed as mean ± SD (standard deviation). A p value <0.05 was considered statistically significant.