



Let-7 microRNAs and opioid tolerance

Ying He and Zaijie Jim Wang*

Department of Biopharmaceutical Sciences, Cancer Center, University of Illinois, Chicago, IL, USA

Edited by:

Leonard Lipovich, Wayne State University, USA

Reviewed by:

Jun Yasuda, The JFCR-Cancer Institute, Japan
Heh-In Im, Korea Institute of Science and Technology, South Korea

***Correspondence:**

Zaijie Jim Wang, Department of Biopharmaceutical Sciences, Cancer Center, University of Illinois, 833 S Wood Street, MC865, Chicago, IL 60612, USA.
e-mail: zjwang@uic.edu

This chapter will focus on the role of microRNAs (miRs) in regulating the actions of opioid drugs through the opioid receptors. Opioids, such as morphine, are analgesics that are used for treating many forms of acute and chronic pain. However, their chronic use is limited by undesirable effects such as opioid tolerance. The μ opioid receptor (MOR) is the primary receptor responsible for opioids' analgesia and antinociceptive tolerance. The long 3'-untranslated region (3'-UTR) of MOR mRNA is of great interest since this region may contain elements for the post-transcriptional regulation of receptor expression, such as altering the stability of mRNA, influencing translational efficiency, and controlling mRNA transport. MicroRNAs are small non-coding RNA molecules that exert their functions through base-pairing with partially complementary sequences in the 3'-UTR of target mRNAs, resulting in decreased polypeptide formation from those mRNAs. Since the discovery of the first miR, lin-4 in *Caenorhabditis elegans*, hundreds of miRs have been identified from humans to viruses, which have provided a crucial and pervasive layer of post-transcriptional gene regulation. The nervous system is a rich source of miR expression, with a diversity of miR functions in fundamental neurobiological processes including neuronal development, plasticity, metabolism, and apoptosis. Recently, the let-7 family of miRs is found to be a critical regulator of MOR function in opioid tolerance. Let-7 is the first identified human miR. Its family members are highly conserved across species in sequence and function. In the review, we will present a brief review of the opioid receptors, their regulation, and opioid tolerance as well as an overview of miRs and a perspective how miRs may interact with MOR and serve as a regulator of opioid tolerance.

Keywords: opioid, miR, epigenetics, addiction, pain

INTRODUCTION

Since first isolated from opium poppy in the early nineteenth century, morphine and related opioids remain as the most powerful analgesics to treat many forms of acute and chronic pain. However, repeated and prolonged use of opioids leads to tolerance, of which, the analgesic tolerance is the most problematic that can lead to therapeutic failure (i.e., inadequate pain control; McQuay, 1999). In some cases, even the highest tolerable dose of an opioid cannot achieve the desirable analgesic effect in patients (Wang and Wang, 2006; Harden, 2008). Development of opioid tolerance not only limits the analgesic efficacy, but the increased doses of opioids in order to counter tolerance exacerbate another problem, namely opioid addiction that is a significant medical and public health problem. Extensive efforts have been made to elucidate the mechanisms underlying opioid tolerance and drug addiction (Kieffer and Evans, 2002). Increasing evidence implicates the contribution of transcriptional and epigenetic regulation in opioid tolerance and drug addiction, such as activation and inhibition of transcription factors (Carlezon et al., 2005; Zachariou et al., 2006), modification of chromatin and DNA structure (Renthal et al., 2008; Guo et al., 2011), and induction of non-coding RNAs including microRNAs (Pietrzykowski, 2011; Robison and Nestler, 2011).

MicroRNAs (miRs) are small non-coding RNA molecules that repress target gene expression through base-pairing with partially complementary sequences in the 3'-untranslated region (3'-UTR)

of target mRNAs. Owing to recent cloning, sequencing, and computational efforts, the numbers of known miRs has been rapidly increasing, and to date, there are a total of 21,643 mature miRs found across 103 species, of which 1921 miRs are found in humans (miRBase Release 18.0, November 2011¹). With the emerging identification of miRs from humans to viruses, a crucial and pervasive layer of post-transcriptional gene regulation by miRs has been elucidated (Ambros, 2004; Taft et al., 2010). It is generally accepted that miRs serve as an important class of epigenetic regulators that participate in a variety of cellular activities. In respect to their diverse functions, miRs play an integral role in fundamental neurobiological processes including neuronal development, plasticity, metabolism, and apoptosis (Kosik, 2006). As one form of long-lasting synaptic plasticity, opioid tolerance is a particularly interesting research area to study miR-mediated cellular adaptation. In this review, we summarize recent findings on miRs in opioid tolerance, with a focus on the role of let-7 miRs in regulating opioid tolerance.

μ OPIOID RECEPTOR AND OPIOID TOLERANCE

The μ opioid receptor (MOR), a member of the G-protein coupled receptor superfamily, is the primary receptor responsible for

¹<http://microrna.sanger.ac.uk/sequences>

opioids' analgesia and antinociceptive tolerance (Matthes et al., 1996; Sora et al., 1997). Opioid tolerance may be result from opioid receptor desensitization and trafficking, which include opioid receptor down-regulation, internalization, and uncoupling from G-proteins due to chronic exposure to opioid agonists (Bailey and Connor, 2005; Martini and Whistler, 2007; Koch and Hollt, 2008; Lopez-Gimenez and Milligan, 2010). Receptor down-regulation as one of mechanisms contributing to opioid tolerance has been previously proposed (Davis et al., 1979; Tao et al., 1987; Bhargava and Gulati, 1990; Bernstein and Welch, 1998; Diaz et al., 2000). Chronic morphine treatment produced a marked decrease in brain MOR density (Davis et al., 1979; Tempel et al., 1988). Down-regulation of the high-affinity MOR site in rats has also been reported following continuous infusion of morphine (*i.t.*; Wong et al., 1996) or etorphine (*s.c.*; Tao et al., 1987). Morphine-induced MOR down-regulation was also observed in SH-SY5Y cells, with or without differentiation (Zadina et al., 1993). In addition to receptor down-regulation, chronic treatment with morphine has been shown to significantly reduce MOR-signaling in sensory neurons and brainstem nuclei (Sim et al., 1996; Johnson et al., 2006), which are in agreement with the findings that reduced receptor number and resultant reduced MOR-signaling contribute to opioid tolerance. On the other hand, there have been reports that MOR expression was not altered (Dum et al., 1979) or even up-regulated (Lewis et al., 1984) in the brain by various opioids. Some of the discrepancies may be caused by uncontrolled variables (e.g., different opioids, doses, methods of opioid treatments, anatomical regions, and times samples were taken, integrality of tissue samples before assays, and opioid receptor subtypes studied), as well as detection methods. It has been suggested that MOR down-regulation is agonist selective and depends on the agonist's intrinsic efficacy (Nishino et al., 1990; Chakrabarti et al., 1997; Chan et al., 1997; Koch and Hollt, 2008). The purity and selectivity of radiolabeled ligands – a problem not only for some early studies, but also in more recent reports employing questionable materials – used in studies are the other potential culprits for conflicting findings.

DIRECT INTERACTION BETWEEN let-7 FAMILY miRs AND MOR

The long 3'-UTR of MOR mRNA (Ide et al., 2005; Han et al., 2006) suggests that this region may contain physiologically relevant elements for regulating receptor expression by mechanisms such as miR targeting. Indeed, early research on 3'-UTR of human MOR mRNA suggested that MOR expression was increased after a 712-bp segment, immediately downstream of the stop codon, was removed (Zollner et al., 2000). Comparative bioinformatics predicted potential miRs that may interact with the human and mouse MOR (Table 1). Let-7 family of miRs was identified as a top candidate according to the number of putative target sites and alignment pattern. Our group experimentally validated the *in silico* prediction that members of the let-7 miR family can interact with the 3'-UTR of MOR mRNA at the predicted positions (He et al., 2010). Furthermore, downregulating let-7 with specific LNA-modified antisense oligodeoxynucleotides (LNA-let-7-AS) was found to increase MOR expression in SH-SY5Y cells, a human neuroblastoma cell line, suggesting that (1) MOR is a target of let-7; (2) expression of MOR is under constitutive suppression by let-7.

Table 1 | MicroRNA targets predicted by miRanda (<http://www.microma.org/microma/getGeneForm.do>) and ranked according to the alignment scores.

miRNA	Query target sites	Alignment score	PhastCons score
HUMAN OPRM1 3'-UTR			
hsa-miR-600	291	169	0.666615
hsa-miR-384	83	157	0.619627
hsa-miR-105	172	156	0.619627
hsa-let-7b	381	154	0.607749
hsa-let-7c	383	154	0.607749
hsa-let-7g	386	154	0.607749
hsa-let-7i	386	154	0.607749
hsa-miR-1272	261	154	0.643121
hsa-miR-642	151	154	0.619627
hsa-miR-98	386	154	0.607749
hsa-let-7f	383	153	0.607749
hsa-let-7d	386	152	0.607749
hsa-miR-18a	401	152	0.607749
hsa-miR-198	124	152	0.619627
hsa-let-7a	385	151	0.607749
hsa-miR-302a	169	151	0.619627
hsa-miR-1324	286	150	0.666615
hsa-miR-192	102	150	0.619627
hsa-miR-215	102	150	0.619627
hsa-miR-648	144	150	0.619627
hsa-miR-18b	399	149	0.607749
hsa-miR-383	389	149	0.607749
hsa-miR-526b	52	149	0.619627
hsa-miR-659	233	149	0.619627
hsa-let-7e	385	147	0.607749
hsa-miR-302b	170	147	0.619627
hsa-miR-136	166	146	0.619627
hsa-miR-431	220	145	0.619627
hsa-miR-557	216	145	0.619627
hsa-miR-1262	105	144	0.619627
hsa-miR-154	322	144	0.666615
hsa-miR-373	168	144	0.619627
hsa-miR-583	114	144	0.619627
hsa-miR-1259	271	143	0.643121
hsa-miR-135a	264	143	0.643121
hsa-miR-135b	264	143	0.643121
hsa-miR-493	4	143	0.6910845
hsa-miR-519c-5p	79	143	0.619627
hsa-miR-520d-3p	170	143	0.619627
hsa-miR-580	307	143	0.666615
hsa-miR-196a	382	142	0.607749
hsa-miR-520a-3p	172	142	0.619627
hsa-miR-577	302	142	0.666615
hsa-miR-142-3p	384	141	0.607749
hsa-miR-378	208	141	0.619627
hsa-miR-422a	207	141	0.619627
hsa-miR-1826	387	140	0.607749
hsa-miR-302c	170	140	0.619627

(Continued)

Table 1 | Continued

miRNA	Query target sites	Alignment score	PhastCons score
hsa-miR-302d	170	140	0.619627
hsa-miR-504	1	140	0.6910845
hsa-miR-521	89	140	0.619627
hsa-miR-9	234	140	0.619627
hsa-miR-942	333	140	0.666615
MOUSE OPRM1 3'-UTR			
mmu-miR-540-5p	278	168.0	0.62498
mmu-miR-540-3p	395	159.0	0.584342
mmu-miR-134	403	158.0	0.584342
mmu-miR-302c	165	158.0	0.586054
mmu-miR-302d	165	151.0	0.586054
mmu-let-7i	384	149.0	0.62498
mmu-miR-139-5p	363	149.0	0.62498
mmu-let-7g	388	148.0	0.62498
mmu-miR-199a-3p	193	148.0	0.586054
mmu-miR-199b	193	148.0	0.586054
mmu-miR-496	241	148.0	0.586054
mmu-miR-532-3p	63	148.0	0.586054
mmu-let-7a	388	147.0	0.62498
mmu-let-7f	388	147.0	0.62498
mmu-let-7c	386	146.0	0.62498
mmu-miR-486	214	146.0	0.586054
mmu-miR-741	512	146.0	0.567263
mmu-miR-383	93	144.0	0.586054
mmu-miR-695	404	144.0	0.584342
mmu-let-7b	388	143.0	0.62498
mmu-let-7e	388	143.0	0.62498
mmu-miR-188-3p	64	143.0	0.586054
mmu-miR-196b	386	143.0	0.62498
mmu-miR-302b	165	143.0	0.586054
mmu-miR-323-5p	233	143.0	0.586054
mmu-miR-466f-5p	138	143.0	0.586054
mmu-miR-98	387	143.0	0.62498
mmu-miR-298	391	142.0	0.584342
mmu-miR-339-3p	71	142.0	0.586054
mmu-miR-381	528	142.0	0.590822
mmu-miR-682	186	142.0	0.586054
mmu-let-7d	388	141.0	0.62498
mmu-miR-504	57	141.0	0.586054
mmu-miR-154	321	140.0	0.62498
mmu-miR-196a	387	140.0	0.62498
mmu-miR-340-5p	527	140.0	0.590822
mmu-miR-34b-3p	367	140.0	0.62498


In order to elucidate a physiological role of let-7 in regulating MOR and opioid tolerance, we further examined the *in vivo* relevance of let-7 miRNAs in cellular and animal models of opioid tolerance. For the former, SH-SY5Y cells were treated with morphine (1–10 μ M for 24–48 h) to induce cellular tolerance. The expression of let-7 miRNAs was significantly up-regulated by chronic treatment with morphine, while the expression level of MOR was reduced as determined by the western blotting method

(He et al., 2010) and the receptor radioligand binding using 3 H-DAMGO (He and Wang, unpublished data). Therefore, chronic morphine treatment caused the increase of let-7 and decrease of MOR expression during the development of opioid tolerance in SH-SY5Y. Furthermore, the regulation of let-7 expression by morphine occurred in a mouse model of opioid tolerance. Mice were implanted *s.c.* with a 75-mg morphine pellet (75 mg morphine pellet/mouse, *s.c.*). Brain expression of let-7 increased gradually over time after morphine treatment, temporally correlating with the development of antinociceptive tolerance to morphine. Interestingly, up-regulation of let-7 occurred in MOR-expressing cells, but not in MOR-negative cells in mice brain cortex region as determined by *in situ* hybridization (He et al., 2010). This was in agreement with the aforementioned finding that MOR is a direct target of let-7. In order to further examine a causative role of let-7 in leading to opioid tolerance, we directly targeted let-7 in the mouse model of opioid tolerance. Treatment with the let-7 inhibitor (*i.c.v.*) decreased brain let-7 levels and partially attenuated opioid antinociceptive tolerance (He et al., 2010).

Previous reports from a number of different cell lines or animal models (e.g., Brodsky et al., 1995; Johnson et al., 2006) indicated that MOR mRNA was not changed upon treatment with morphine. We also found that the total MOR transcripts were unchanged by morphine. For example, MOR mRNA remained the same in SH-SY5Y cells following chronic morphine treatment. It raised a question as to how let-7 miRNAs repress MOR in opioid tolerance. We confirmed that let-7 did not affect MOR mRNA stability; however, polysome-bound MOR transcript was significantly decreased. Where is MOR mRNA hiding? It turns out that it can be accumulated in P-bodies. So these intriguing data led us to propose the following regulatory mechanism where upon let-7 can regulate opioid tolerance: let-7 recruits and sequesters MOR mRNA to P-bodies that are deprived of translational machinery, effectively reducing polysome-bound MOR mRNA and leading to translation repression. A similar pathway was observed in HEK293 and HeLa cells (Pillai et al., 2005), thus translation repression may serve as a general mechanism by let-7 to regulate its target gene expression (Figure 1). Keep in mind that let-7 activity is turned up in opioid tolerance by up-regulation of their transcripts. Nature works in a wonder to ensure the activity of let-7, ultimately dampening the activity of opioids upon persistent activation.

POTENTIAL TARGET GENES OF let-7 IN OPIOID TOLERANCE

Let-7 was the first identified human miR. Its family members are highly conserved across species in sequence and function (Pasquinelli et al., 2000). Major roles of let-7 include the regulation of stem-cell differentiation, neuromuscular development, and cell proliferation & differentiation (Reinhart et al., 2000; Mansfield et al., 2004; Roush and Slack, 2008). Let-7 was initially identified as a heterochronic gene (Pasquinelli et al., 2000). In mammals, let-7 levels increase during embryogenesis and during brain development (Schulman et al., 2005; Wulczyn et al., 2007). Let-7 is undetectable in human and mouse embryonic stem cells, and the level of let-7 increases upon differentiation (Thomson et al., 2004, 2006; Wulczyn et al., 2007). This high expression of let-7 is then maintained in various adult tissues (Sempere et al., 2004; Thomson et al., 2004). Furthermore, let-7 is widely viewed



as a tumor suppressor miR (Boyerinas et al., 2010). There is a clear link between loss of let-7 expression and the development of poorly differentiated, aggressive cancers (Takamizawa et al., 2004; Dahiya et al., 2008; Childs et al., 2009). Using the computational algorithm TargetScan 6.0² to screen human 3'-UTR sequences containing let-7 family miRs complementary sites, 886 conserved targets, with a total of 989 conserved sites and 111 poorly conserved sites were predicted. High-mobility group AT-hook 2 (HMGA2), which is the top-scoring candidate gene on the list (Table 2), was confirmed as a direct let-7 target both *in vitro* and *in vivo* (Lee and Dutta, 2007; Mayr et al., 2007; Shell et al., 2007). HMGA2 is a chromatin-associated non-histone protein capable of modulating chromatin architecture and thus affecting transcription. In addition to abundant studies on HMGA2 in embryogenesis (Zhou et al., 1995; Sun et al., 2009) and tumorigenesis (Peng et al., 2008; Qian et al., 2009), it would be interesting to investigate its possible involvement in opioid tolerance as a functional target gene of let-7.

Seed pairing rules are widely used to predict functional miR target sites. miR-mRNA recognition requires the perfect complementarity of 6-nucleotide mRNA seed sites with the 5' end of miRs (positions 2–7; Bartel, 2009). However, when predictions based on such short complementary sequences are compared to experimental results from proteomic studies, the false-positive and false-negative rates appear to be above 50% (Easow et al., 2007; Baek et al., 2008; Mourelatos, 2008; Selbach et al., 2008). Several previous findings strongly indicated that a sizeable fraction of miR targets do not obey the “canonical” seed rule (Ha et al., 1996; Didi-ano and Hobert, 2006; Tay et al., 2008; Stefani and Slack, 2012). For let-7, biological studies clearly demonstrated that genetically verified let-7 targets in *Caenorhabditis elegans* contain only imperfect binding sites, with bulges or G-U wobble pairs in the seed region (Vella et al., 2004). A recent study identified a new class of miR target site nucleation bulges and an alternative mode of miR target recognition by a pivot-pairing rule (Chi et al., 2012). They proposed a transitional nucleation model in which a transitional nucleation state determines the binding of miRs to nucleation

bulge mRNAs. Therefore, the identification of non-canonical let-7-mRNA interactions may lead to important breakthroughs in discovering new let-7 targets and further decipher the functional role of let-7 in opioid tolerance.

With respect to let-7 target gene identification, another important issue need to be addressed is the redundancy of let-7 family members. There are 14 and 13 different let-7 family members in mouse and human, respectively (Roush and Slack, 2008). In human, these different members, let-7a-1, 7a-2, 7a-3, 7b, 7c, 7d, 7e, f7-1, 7f-2, 7g, 7i, miR-98, and miR-202 are located on nine different chromosomes (Ruby et al., 2006). While let-7 was initially viewed as one single activity, emerging data suggest that the let-7 family contains miRs with different activities (i.e., targets). For example, it has been reported that let-7b* was highly expressed but let-7e* was drastically reduced in malignant mesothelioma (Guled et al., 2009). So the question remains as whether individual let-7 family member with its own expression pattern exerts specialized function during opioid tolerance development.

CONCLUSION AND FUTURE DIRECTIONS

Opioid tolerance, even to the analgesic actions of these drugs, is likely not caused by a single mechanism; rather an intricate neuronal circuitry involving multiple mechanisms may ultimately lead to the expression of tolerance seen in humans. Regulation of opioid tolerance, through MOR or other mechanisms, represents one of these mechanisms. What is not known is how many miRs are involved. Existing literature on miR-related mechanisms of opioid tolerance is sparse. A typical study in miR research field tends to survey transcript levels of miRs in a disease state (Zheng et al., 2010). For example, morphine-induced up-regulation of miR-15b in human monocyte-derived macrophages (Dave and Khalili, 2010), while miR-133b was decreased by morphine in zebrafish embryos (Sanchez-Simon et al., 2010). It is often extremely difficult to identify a casual relationship. Furthermore, moving from a cell line to *in vivo* relevance is another big hurdle to overcome.

In the case of let-7 miRs, it was identified as a critical regulator of both human and mouse MOR. Moreover, it was demonstrated to be relevant for both cellular opioid tolerance as well as animal models of opioid tolerance (He et al., 2010). Let-7 represents one of several miRs contributing to opioid tolerance. For example, it

²<http://www.targetscan.org/>

Table 2 | Top 100 predicted targets of let-7 miRNAs ranked by total context score (TargetScan 6.0, <http://www.targetscan.org/>.)

Target gene (1–50)	Total context score	Aggregate P_{CT}	Target gene (51–100)	Total context score	Aggregate P_{CT}
Hmga2	−1.08	>0.99	Map4k3	−0.46	0.96
Fignl2	−1.07	>0.99	Tmem211	−0.46	<0.1
Lin28b	−0.98	>0.99	Zfp583	−0.46	0.96
Trim71	−0.98	>0.99	Dtx2	−0.46	0.99
Zc3hav11	−0.85	0.86	Diap2	−0.46	0.98
Fign	−0.84	>0.99	Pgrmc1	−0.46	0.9
Nr6a1	−0.8	>0.99	Pars2	−0.46	<0.1
2200002K05Rik	−0.7	>0.99	Cd200r1	−0.45	0.94
Skil	−0.67	>0.99	Abt1	−0.45	<0.1
Igdcc3	−0.65	>0.99	Acot11	−0.45	0.5
Thrsp	−0.62	<0.1	Kcnj11	−0.45	0.81
Prtg	−0.61	>0.99	Zfp248	−0.45	<0.1
Slc5a9	−0.6	0.83	Bcl2l13	−0.44	<0.1
Tgfb1	−0.58	>0.99	Gatm	−0.44	0.95
Yod1	−0.58	>0.99	Hic2	−0.44	>0.99
Smarcad1	−0.57	0.96	Ccnj	−0.44	0.97
Gab2	−0.57	0.6	Arid3a	−0.44	>0.99
Ngly1	−0.54	<0.1	Hand1	−0.44	0.98
Kctd21	−0.54	0.98	Igf1r	−0.44	>0.99
Dna2	−0.53	<0.1	Rpusd3	−0.43	0.59
Ppp1r15b	−0.53	>0.99	Pbx3	−0.43	0.99
Nphp3	−0.53	0.94	Zmat4	−0.43	0.97
Vezt	−0.52	<0.1	Tmem164	−0.43	<0.1
Suv39h2	−0.51	0.44	Bcat1	−0.43	0.99
Gdf6	−0.5	0.98	Pxt1	−0.43	<0.1
Brwd1	−0.5	<0.1	Als2cl	−0.43	<0.1
Coil	−0.49	0.97	9930012K11Rik	−0.42	0.94
Lrig3	−0.49	0.89	Atg10	−0.42	<0.1
Hoxb1	−0.49	0.98	Limd2	−0.42	0.94
Zcchc9	−0.48	<0.1	Zfp341	−0.42	<0.1
B3gnt7	−0.48	0.76	Fam103a1	−0.42	0.87
Entpd7	−0.48	<0.1	Med28	−0.42	0.56
Acvr1c	−0.48	0.99	Smug1	−0.42	<0.1
Lrig2	−0.48	0.99	Trpm3	−0.42	<0.1
Bzw1	−0.48	>0.99	Fras1	−0.42	0.96
Gnptab	−0.48	0.98	Uhrf2	−0.41	0.86
Wdr41	−0.48	<0.1	Cdc25a	−0.41	0.8
Cdc34	−0.47	0.99	Igf2bp1	−0.41	>0.99
Slc35d2	−0.47	0.98	Cep110	−0.41	0.94
Dclre1b	−0.47	0.12	Mrps33	−0.41	<0.1
Hif1an	−0.47	0.61	Tnfsf10	−0.41	<0.1
Zfp322a	−0.47	0.82	Arhgap28	−0.41	0.98
Adrb2	−0.47	0.98	Apbb3	−0.41	0.98
Tmprss11f	−0.47	0.92	Fndc3a	−0.41	0.97
Ddx19b	−0.47	0.98	Col1a2	−0.41	0.97
Ddx19a	−0.47	0.98	Gemin7	−0.41	<0.1
Galnt1	−0.47	>0.99	Wnt9a	−0.41	0.99
Greb1l	−0.47	0.99	Igf2bp2	−0.41	0.93
Erc6	−0.47	0.98	Tmem2	−0.41	0.98
Ddi2	−0.46	0.89	Zfp879	−0.41	0.35

was found miR-23b could interact with the MOR 3'-UTR via a K box motif (5'-UGUGAU-3') in SH-SY5Y and mouse P19 cells (Wu et al., 2008, 2009).

While MOR is involved in the development of morphine tolerance, it is not the only target for the action of miRs in opioid tolerance (Matthes et al., 1996; Sora et al., 1997). With the increasing understanding of miRs in epigenetic regulation, further research is needed to identify other target genes of miRs (including let-7). A number of receptors, ion channels, and protein kinases, which have

been implicated in opioid tolerance, such as the NMDA receptors, PKC, and CaMKII (Kieffer and Evans, 2002; Wang and Wang, 2006; Ueda and Ueda, 2009) may become potential targets for let-7.

In summary, miR-mediated mechanisms provide a novel direction toward unraveling the complex mechanisms involved in the development of opioid tolerance. These studies will enrich our knowledge on basic principles of neuronal and behavioral adaptation in opioid tolerance, and eventually lead to novel design and development of pharmacological treatments for opioid tolerance.

REFERENCES

- Ambros, V. (2004). The functions of animal microRNAs. *Nature* 431, 350–355.
- Baek, D., Villen, J., Shin, C., Camargo, F. D., Gygi, S. P., and Bartel, D. P. (2008). The impact of microRNAs on protein output. *Nature* 455, 64–71.
- Bailey, C. P., and Connor, M. (2005). Opioids: cellular mechanisms of tolerance and physical dependence. *Curr. Opin. Pharmacol.* 5, 60–68.
- Bartel, D. P. (2009). MicroRNAs: target recognition and regulatory functions. *Cell* 136, 215–233.
- Bernstein, M. A., and Welch, S. P. (1998). mu-Opioid receptor down-regulation and cAMP-dependent protein kinase phosphorylation in a mouse model of chronic morphine tolerance. *Brain Res. Mol. Brain Res.* 55, 237–242.
- Bhargava, H. N., and Gulati, A. (1990). Down-regulation of brain and spinal cord mu-opiate receptors in morphine tolerant-dependent rats. *Eur. J. Pharmacol.* 190, 305–311.
- Boyerinas, B., Park, S. M., Hau, A., Murmann, A. E., and Peter, M. E. (2010). The role of let-7 in cell differentiation and cancer. *Endocr. Relat. Cancer* 17, F19–F36.
- Brodsky, M., Elliott, K., Hynansky, A., and Inturrisi, C. E. (1995). CNS levels of mu opioid receptor (MOR-1) mRNA during chronic treatment with morphine or naltrexone. *Brain Res. Bull.* 38, 135–141.
- Carlezon, W. A. Jr., Duman, R. S., and Nestler, E. J. (2005). The many faces of CREB. *Trends Neurosci.* 28, 436–445.
- Chakrabarti, S., Yang, W., Law, P. Y., and Loh, H. H. (1997). The mu-opioid receptor down-regulates differently from the delta-opioid receptor: requirement of a high affinity receptor/G protein complex formation. *Mol. Pharmacol.* 52, 105–113.
- Chan, K. W., Duttory, A., and Yoburn, B. C. (1997). Magnitude of tolerance to fentanyl is independent of mu-opioid receptor density. *Eur. J. Pharmacol.* 319, 225–228.
- Chi, S. W., Hannon, G. J., and Darnell, R. B. (2012). An alternative mode of microRNA target recognition. *Nat. Struct. Mol. Biol.* 19, 321–327.
- Childs, G., Fazzari, M., Kung, G., Kawachi, N., Brandwein-Gensler, M., McLemore, M., Chen, Q., Burk, R. D., Smith, R. V., Prystowsky, M. B., Belbin, T. J., and Schlecht, N. F. (2009). Low-level expression of microRNAs let-7d and miR-205 are prognostic markers of head and neck squamous cell carcinoma. *Am. J. Pathol.* 174, 736–745.
- Dahiya, N., Sherman-Baust, C. A., Wang, T. L., Davidson, B., Shih, Ie, M., Zhang, Y., Wood, W. III, Becker, K. G., and Morin, P. J. (2008). MicroRNA expression and identification of putative miRNA targets in ovarian cancer. *PLoS ONE* 3, e2436. doi:10.1371/journal.pone.0002436
- Dave, R. S., and Khalili, K. (2010). Morphine treatment of human monocyte-derived macrophages induces differential miRNA and protein expression: impact on inflammation and oxidative stress in the central nervous system. *J. Cell. Biochem.* 110, 834–845.
- Davis, M. E., Akera, T., and Brody, T. M. (1979). Reduction of opiate binding to brainstem slices associated with the development of tolerance to morphine in rats. *J. Pharmacol. Exp. Ther.* 211, 112–119.
- Diaz, A., Pazos, A., Florez, J., and Hurler, M. A. (2000). Autoradiographic mapping of mu-opioid receptors during opiate tolerance and supersensitivity in the rat central nervous system. *Naunyn Schmiedebergs Arch. Pharmacol.* 362, 101–109.
- Didiano, D., and Hobert, O. (2006). Perfect seed pairing is not a generally reliable predictor for miRNA-target interactions. *Nat. Struct. Mol. Biol.* 13, 849–851.
- Dum, J., Meyer, G., Holtt, V., and Herz, A. (1979). In vivo opiate binding unchanged in tolerant/dependent mice. *Eur. J. Pharmacol.* 58, 453–460.
- Easow, G., Teleman, A. A., and Cohen, S. M. (2007). Isolation of microRNA targets by miRNP immunoprecipitation. *RNA* 13, 1198–1204.
- Guled, M., Lahti, L., Lindholm, P. M., Salmenkivi, K., Bagwan, I., Nicholson, A. G., and Knuutila, S. (2009). CDKN2A, NF2, and JUN are dysregulated among other genes by miRNAs in malignant mesothelioma - A miRNA microarray analysis. *Genes Chromosomes Cancer* 48, 615–623.
- Guo, J. U., Su, Y., Zhong, C., Ming, G. L., and Song, H. (2011). Hydroxylation of 5-methylcytosine by TET1 promotes active DNA demethylation in the adult brain. *Cell* 145, 423–434.
- Ha, I., Wightman, B., and Ruvkun, G. (1996). A bulged lin-4/lin-14 RNA duplex is sufficient for *Caenorhabditis elegans* lin-14 temporal gradient formation. *Genes Dev.* 10, 3041–3050.
- Han, W., Kasai, S., Hata, H., Takahashi, T., Takamatsu, Y., Yamamoto, H., Uhl, G. R., Sora, I., and Ikeda, K. (2006). Intracisternal a-particle element in the 3' noncoding region of the mu-opioid receptor gene in CXBK mice: a new genetic mechanism underlying differences in opioid sensitivity. *Pharmacogenet. Genomics* 16, 451–460.
- Harden, R. N. (2008). Chronic pain and opiates: a call for moderation. *Arch. Phys. Med. Rehabil.* 89, S72–S76.
- He, Y., Yang, C., Kirkmire, C. M., and Wang, Z. J. (2010). Regulation of opioid tolerance by let-7 family microRNA targeting the mu opioid receptor. *J. Neurosci.* 30, 10251–10258.
- Ide, S., Han, W., Kasai, S., Hata, H., Sora, I., and Ikeda, K. (2005). Characterization of the 3' untranslated region of the human mu-opioid receptor (MOR-1) mRNA. *Gene* 364, 139–145.
- Johnson, E. E., Chieng, B., Napier, I., and Connor, M. (2006). Decreased mu-opioid receptor signalling and a reduction in calcium current density in sensory neurons from chronically morphine-treated mice. *Br. J. Pharmacol.* 148, 947–955.
- Kieffer, B. L., and Evans, C. J. (2002). Opioid tolerance-in search of the holy grail. *Cell* 108, 587–590.
- Koch, T., and Holtt, V. (2008). Role of receptor internalization in opioid tolerance and dependence. *Pharmacol. Ther.* 117, 199–206.
- Kosik, K. S. (2006). The neuronal microRNA system. *Nat. Rev. Neurosci.* 7, 911–920.
- Lee, Y. S., and Dutta, A. (2007). The tumor suppressor microRNA let-7 represses the HMG2 oncogene. *Genes Dev.* 21, 1025–1030.
- Lewis, J. W., Lewis, M. E., Loomus, D. J., and Akil, H. (1984). Acute systemic administration of morphine selectively increases mu opioid receptor binding in the rat brain. *Neuropeptides* 5, 117–120.
- Lopez-Gimenez, J. F., and Milligan, G. (2010). Opioid regulation of mu receptor internalisation: relevance to the development of tolerance and dependence. *CNS Neurol. Disord. Drug Targets* 9, 616–626.
- Mansfield, J. H., Harfe, B. D., Nissen, R., Obenaus, J., Srinell, J., Chaudhuri, A., Farzan-Kashani, R., Zuker, M., Pasquinelli, A. E., Ruvkun, G., Sharp, P. A., Tabin, C. J., and McManus, M. T. (2004). MicroRNA-responsive 'sensor' transgenes uncover Hox-like and other developmentally regulated patterns of vertebrate microRNA expression. *Nat. Genet.* 36, 1079–1083.
- Martini, L., and Whistler, J. L. (2007). The role of mu opioid receptor desensitization and endocytosis in morphine tolerance and dependence. *Curr. Opin. Neurobiol.* 17, 556–564.
- Matthes, H. W., Maldonado, R., Simonin, F., Valverde, O., Slowe, S., Kitchen, I., Befort, K., Dierich, A., Le Meur, M., Dolle, P., Tzavara, E., Hanoune, J., Roques, B. P., and Kieffer, B. L. (1996). Loss of morphine-induced analgesia, reward effect and withdrawal symptoms in mice lacking the mu-opioid-receptor gene. *Nature* 383, 819–823.
- Mayr, C., Hemann, M. T., and Bartel, D. P. (2007). Disrupting the pairing between let-7 and Hmga2 enhances oncogenic transformation. *Science* 315, 1576–1579.
- McQuay, H. (1999). Opioids in pain management. *Lancet* 353, 2229–2232.
- Mourelatos, Z. (2008). Small RNAs: the seeds of silence. *Nature* 455, 44–45.

- Nishino, K., Su, Y. F., Wong, C. S., Watkins, W. D., and Chang, K. J. (1990). Dissociation of mu opioid tolerance from receptor down-regulation in rat spinal cord. *J. Pharmacol. Exp. Ther.* 253, 67–72.
- Pasquinelli, A. E., Reinhart, B. J., Slack, F., Martindale, M. Q., Kuroda, M. I., Maller, B., Hayward, D. C., Ball, E. E., Degnan, B., Muller, P., Spring, J., Srinivasan, A., Fishman, M., Finnerty, J., Corbo, J., Levine, M., Leahy, P., Davidson, E., and Ruvkun, G. (2000). Conservation of the sequence and temporal expression of let-7 heterochronic regulatory RNA. *Nature* 408, 86–89.
- Peng, Y., Laser, J., Shi, G., Mittal, K., Melamed, J., Lee, P., and Wei, J. J. (2008). Antiproliferative effects by Let-7 repression of high-mobility group A2 in uterine leiomyoma. *Mol. Cancer Res.* 6, 663–673.
- Pietrzykowski, A. Z. (2011). The role of microRNAs in drug addiction: a big lesson from tiny molecules. *Int. Rev. Neurobiol.* 91, 1–24.
- Pillai, R. S., Bhattacharyya, S. N., Artus, C. G., Zoller, T., Cougot, N., Basyuk, E., Bertrand, E., and Filipowicz, W. (2005). Inhibition of translational initiation by Let-7 MicroRNA in human cells. *Science* 309, 1573–1576.
- Qian, Z. R., Asa, S. L., Siomi, H., Siomi, M. C., Yoshimoto, K., Yamada, S., Wang, E. L., Rahman, M. M., Inoue, H., Itakura, M., Kudo, E., and Sano, T. (2009). Overexpression of HMGA2 relates to reduction of the let-7 and its relationship to clinicopathological features in pituitary adenomas. *Mod. Pathol.* 22, 431–441.
- Reinhart, B. J., Slack, F. J., Basson, M., Pasquinelli, A. E., Bettinger, J. C., Rougvie, A. E., Horvitz, H. R., and Ruvkun, G. (2000). The 21-nucleotide let-7 RNA regulates developmental timing in *Caenorhabditis elegans*. *Nature* 403, 901–906.
- Renthal, W., Carle, T. L., Maze, I., Covington, H. E. III, Truong, H. T., Alibhai, I., Kumar, A., Montgomery, R. L., Olson, E. N., and Nestler, E. J. (2008). Delta FosB mediates epigenetic desensitization of the c-fos gene after chronic amphetamine exposure. *J. Neurosci.* 28, 7344–7349.
- Robison, A. J., and Nestler, E. J. (2011). Transcriptional and epigenetic mechanisms of addiction. *Nat. Rev. Neurosci.* 12, 623–637.
- Roush, S., and Slack, F. J. (2008). The let-7 family of microRNAs. *Trends Cell Biol.* 18, 505–516.
- Ruby, J. G., Jan, C., Player, C., Axtell, M. J., Lee, W., Nusbaum, C., Ge, H., and Bartel, D. P. (2006). Large-scale sequencing reveals 21U-RNAs and additional microRNAs and endogenous siRNAs in *C. elegans*. *Cell* 127, 1193–1207.
- Sanchez-Simon, F. M., Zhang, X. X., Loh, H. H., Law, P. Y., and Rodriguez, R. E. (2010). Morphine regulates dopaminergic neuron differentiation via miR-133b. *Mol. Pharmacol.* 78, 935–942.
- Schulman, B. R., Esquela-Kerscher, A., and Slack, F. J. (2005). Reciprocal expression of lin-41 and the microRNAs let-7 and mir-125 during mouse embryogenesis. *Dev. Dyn.* 234, 1046–1054.
- Selbach, M., Schwanhauser, B., Thierfelder, N., Fang, Z., Khanin, R., and Rajewsky, N. (2008). Widespread changes in protein synthesis induced by microRNAs. *Nature* 455, 58–63.
- Sempere, L. F., Freemantle, S., Pitha-Rowe, I., Moss, E., Dmitrovsky, E., and Ambros, V. (2004). Expression profiling of mammalian microRNAs uncovers a subset of brain-expressed microRNAs with possible roles in murine and human neuronal differentiation. *Genome Biol.* 5, R13.
- Shell, S., Park, S. M., Radjabi, A. R., Schickel, R., Kistner, E. O., Jewell, D. A., Feig, C., Lengyel, E., and Peter, M. E. (2007). Let-7 expression defines two differentiation stages of cancer. *Proc. Natl. Acad. Sci. U.S.A.* 104, 11400–11405.
- Sim, L. J., Selley, D. E., Dworkin, S. I., and Childers, S. R. (1996). Effects of chronic morphine administration on mu opioid receptor-stimulated [35S]GTPgammaS autoradiography in rat brain. *J. Neurosci.* 16, 2684–2692.
- Sora, I., Takahashi, N., Funada, M., Ujike, H., Revay, R. S., Donovan, D. M., Miner, L. L., and Uhl, G. R. (1997). Opiate receptor knockout mice define mu receptor roles in endogenous nociceptive responses and morphine-induced analgesia. *Proc. Natl. Acad. Sci. U.S.A.* 94, 1544–1549.
- Stefani, G., and Slack, F. J. (2012). A 'pivotal' new rule for microRNA-mRNA interactions. *Nat. Struct. Mol. Biol.* 19, 265–266.
- Sun, T., Fu, M., Bookout, A. L., Kliewer, S. A., and Mangelsdorf, D. J. (2009). MicroRNA let-7 regulates 3T3-L1 adipogenesis. *Mol. Endocrinol.* 23, 925–931.
- Taft, R. J., Pang, K. C., Mercer, T. R., Dinger, M., and Mattick, J. S. (2010). Non-coding RNAs: regulators of disease. *J. Pathol.* 220, 126–139.
- Takamizawa, J., Konishi, H., Yanagisawa, K., Tomida, S., Osada, H., Endoh, H., Harano, T., Yatabe, Y., Nagino, M., Nimura, Y., Mitsudomi, T., and Takahashi, T. (2004). Reduced expression of the let-7 microRNAs in human lung cancers in association with shortened postoperative survival. *Cancer Res.* 64, 3753–3756.
- Tao, P. L., Law, P. Y., and Loh, H. H. (1987). Decrease in delta and mu opioid receptor binding capacity in rat brain after chronic etorphine treatment. *J. Pharmacol. Exp. Ther.* 240, 809–816.
- Tay, Y., Zhang, J., Thomson, A. M., Lim, B., and Rigoutsos, I. (2008). MicroRNAs to Nanog, Oct4 and Sox2 coding regions modulate embryonic stem cell differentiation. *Nature* 455, 1124–1128.
- Tempel, A., Habas, J., Paredes, W., and Barr, G. A. (1988). Morphine-induced downregulation of mu-opioid receptors in neonatal rat brain. *Brain Res.* 469, 129–133.
- Thomson, J. M., Newman, M., Parker, J. S., Morin-Kensicki, E. M., Wright, T., and Hammond, S. M. (2006). Extensive post-transcriptional regulation of microRNAs and its implications for cancer. *Genes Dev.* 20, 2202–2207.
- Thomson, J. M., Parker, J., Perou, C. M., and Hammond, S. M. (2004). A custom microarray platform for analysis of microRNA gene expression. *Nat. Methods* 1, 47–53.
- Ueda, H., and Ueda, M. (2009). Mechanisms underlying morphine analgesic tolerance and dependence. *Front. Biosci.* 14, 5260–5272.
- Vella, M. C., Choi, E. Y., Lin, S. Y., Reinert, K., and Slack, F. J. (2004). The *C. elegans* microRNA let-7 binds to imperfect let-7 complementary sites from the lin-41 3'UTR. *Genes Dev.* 18, 132–137.
- Wang, Z. J., and Wang, L. X. (2006). Phosphorylation: a molecular switch in opioid tolerance. *Life Sci.* 79, 1681–1691.
- Wong, C. S., Cherng, C. H., Luk, H. N., Ho, S. T., and Tung, C. S. (1996). Effects of NMDA receptor antagonists on inhibition of morphine tolerance in rats: binding at mu-opioid receptors. *Eur. J. Pharmacol.* 297, 27–33.
- Wu, Q., Law, P. Y., Wei, L. N., and Loh, H. H. (2008). Post-transcriptional regulation of mouse mu opioid receptor (MOR1) via its 3' untranslated region: a role for microRNA23b. *FASEB J.* 22, 4085–4095.
- Wu, Q., Zhang, L., Law, P. Y., Wei, L. N., and Loh, H. H. (2009). Long-term morphine treatment decreases the association of mu-opioid receptor (MOR1) mRNA with polysomes through miRNA23b. *Mol. Pharmacol.* 75, 744–750.
- Wulczyn, F. G., Smirnova, L., Rybak, A., Brandt, C., Kwizdzinski, E., Ninnemann, O., Strehle, M., Seiler, A., Schumacher, S., and Nitsch, R. (2007). Post-transcriptional regulation of the let-7 microRNA during neural cell specification. *FASEB J.* 21, 415–426.
- Zachariou, V., Bolanos, C. A., Selley, D. E., Theobald, D., Cassidy, M. P., Kelz, M. B., Shaw-Lutchman, T., Berton, O., Sim-Selley, L. J., Dileone, R. J., Kumar, A., and Nestler, E. J. (2006). An essential role for DeltaFosB in the nucleus accumbens in morphine action. *Nat. Neurosci.* 9, 205–211.
- Zadina, J. E., Chang, S. L., Ge, L. J., and Kastin, A. J. (1993). Mu opiate receptor down-regulation by morphine and up-regulation by naloxone in SH-SY5Y human neuroblastoma cells. *J. Pharmacol. Exp. Ther.* 265, 254–262.
- Zheng, H., Zeng, Y., Zhang, X., Chu, J., Loh, H. H., and Law, P. Y. (2010). mu-Opioid receptor agonists differentially regulate the expression of miR-190 and NeuroD. *Mol. Pharmacol.* 77, 102–109.
- Zhou, X., Benson, K. F., Ashar, H. R., and Chada, K. (1995). Mutation responsible for the mouse pygmy phenotype in the developmentally regulated factor HMGI-C. *Nature* 376, 771–774.
- Zollner, C., Johnson, P. S., Bei Wang, J., Roy, A. J. Jr., Layton, K. M., Min Wu, J., and Surratt, C. K. (2000). Control of mu opioid receptor expression by modification of cDNA 5'- and 3'-noncoding regions. *Brain Res. Mol. Brain Res.* 79, 159–162.

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Received: 20 March 2012; accepted: 29 May 2012; published online: 21 June 2012.

Citation: He Y and Wang ZJ (2012) Let-7 microRNAs and opioid tolerance. *Front. Gene.* 3:110. doi: 10.3389/fgene.2012.00110

This article was submitted to *Frontiers in Non-Coding RNA*, a specialty of *Frontiers in Genetics*.

Copyright © 2012 He and Wang. This is an open-access article distributed under the terms of the Creative Commons Attribution Non-Commercial License, which permits non-commercial use, distribution, and reproduction in other forums, provided the original authors and source are credited.