

PATIENT

DISEASE Lung adenocarcinoma
NAME
DATE OF BIRTH
SEX
MEDICAL RECORD #

PHYSICIAN

ORDERING PHYSICIAN
MEDICAL FACILITY
ADDITIONAL RECIPIENT
MEDICAL FACILITY ID
PATHOLOGIST

SPECIMEN

SPECIMEN SITE
SPECIMEN ID
SPECIMEN TYPE
DATE OF COLLECTION
SPECIMEN RECEIVED

CDx Associated Findings

GENOMIC FINDINGS DETECTED

EGFR exon 19 deletion (T751_I759>S)

FDA-APPROVED THERAPEUTIC OPTIONS

Gilotrif® (Afatinib)
 Iressa® (Gefitinib)
 Tarceva® (Erlotinib)

OTHER ALTERATIONS & BIOMARKERS IDENTIFIED

Results reported in this section are not prescriptive or conclusive for labeled use of any specific therapeutic product. See *professional services* section for additional information.

Microsatellite status MS-Stable§
Tumor Mutation Burden 5.04 Muts/Mb§
CDKN2A loss§
CDKN2B loss§
EGFR amplification§
ERBB3 P1212S

MET T263M
MTAP loss§
NFKBIA amplification§
NKX2-1 amplification§
TP53 R282G

§ Refer to appendix for limitation statements related to detection of any copy number alterations, gene rearrangements, MSI or TMB result in this section.

Please refer to appendix for Explanation of Clinical Significance Classification and for variants of unknown significance (VUS).

FoundationOne CDx™ (FICDx) is a next generation sequencing based in vitro diagnostic device for detection of substitutions, insertion and deletion alterations (indels), and copy number alterations (CNAs) in 324 genes and select gene rearrangements, as well as genomic signatures including microsatellite instability (MSI) and tumor mutational burden (TMB) using DNA isolated from formalin-fixed paraffin embedded (FFPE) tumor tissue specimens. The test is intended as a companion diagnostic to identify patients who may benefit from treatment with the targeted therapies listed in Table 1 in accordance with the approved therapeutic product labeling. Additionally, FICDx is intended to provide tumor mutation profiling to be used by qualified health care professionals in accordance with professional guidelines in oncology for patients with solid malignant neoplasms. The FICDx assay is a single-site assay performed at Foundation Medicine, Inc.

TABLE 1

INDICATIONS	BIOMARKER	THERAPY
Non-small cell lung cancer (NSCLC)	EGFR exon 19 deletions and EGFR exon 21 L858R alterations	Gilotrif® (Afatinib), Iressa® (Gefitinib), or Tarceva® (Erlotinib)
	EGFR exon 20 T790M alterations	Tagrisso® (Osimertinib)
	ALK rearrangements	Alecensa® (Alectinib), Xalkori® (Crizotinib), or Zykadia® (Ceritinib)
	BRAF V600E	Tafinlar® (Dabrafenib) in combination with Mekinist® (Trametinib)
Melanoma	BRAF V600E	Tafinlar® (Dabrafenib) or Zelboraf® (Vemurafenib)
	BRAF V600E or V600K	Mekinist® (Trametinib) or Cotellic® (Cobimetinib) in combination with Zelboraf® (Vemurafenib)
Breast cancer	ERBB2 (HER2) amplification	Herceptin® (Trastuzumab), Kadcyla® (Ado-trastuzumab emtansine), or Perjeta® (Pertuzumab)
Colorectal cancer	KRAS wild-type (absence of mutations in codons 12 and 13)	Erbix® (Cetuximab)
	KRAS wild-type (absence of mutations in exons 2, 3, and 4) and NRAS wild type (absence of mutations in exons 2, 3, and 4)	Vectibix® (Panitumumab)
Ovarian cancer	BRCA1/2 alterations	Rubraca® (Rucaparib)

ABOUT THE TEST FoundationOne CDx™ is the first FDA-approved broad companion diagnostic for solid tumors.

Interpretive content on this page and subsequent pages is provided as a professional service, and is not reviewed or approved by the FDA.

PATIENT

DISEASE **Lung adenocarcinoma**
NAME
DATE OF BIRTH
SEX
MEDICAL RECORD #

PHYSICIAN

ORDERING PHYSICIAN
MEDICAL FACILITY
ADDITIONAL RECIPIENT
MEDICAL FACILITY ID
PATHOLOGIST

SPECIMEN

SPECIMEN SITE
SPECIMEN ID
SPECIMEN TYPE
DATE OF COLLECTION
SPECIMEN RECEIVED

Biomarker Findings

Microsatellite Status - MS-Stable

Tumor Mutation Burden - TMB-Low (5 Muts/Mb)

Genomic Findings

For a complete list of the genes assayed, please refer to the Appendix.

EGFR amplification, exon 19 deletion (T751_I759>S)

ERBB3 P1212S

MET T263M - subclonal †

CDKN2A/B loss

MTAP loss

NFKBIA amplification

NKX2-1 amplification

TP53 R282G

6 Disease relevant genes with no reportable alterations: **KRAS, ALK, BRAF, RET, ERBB2, ROS1**

† See About the Test in appendix for details.

5 Therapies with Clinical Benefit in patient's tumor type

25 Clinical Trials

8 Therapies with Clinical Benefit in other tumor type

BIOMARKER FINDINGS

Microsatellite status - MS-Stable

Tumor Mutation Burden - TMB-Low (5 Muts/Mb)

GENOMIC FINDINGS

EGFR - amplification, exon 19 deletion (T751_I759>S)

10 Trials see p. 15

ERBB3 - P1212S

5 Trials see p. 18

MET - T263M

10 Trials see p. 19

ACTIONABILITY

No therapies or clinical trials. see Biomarker Findings section

No therapies or clinical trials. see Biomarker Findings section

THERAPIES WITH CLINICAL BENEFIT (IN PATIENT'S TUMOR TYPE)

Afatinib

Erlotinib

Gefitinib

Osimertinib

Afatinib

Crizotinib

THERAPIES WITH CLINICAL BENEFIT (IN OTHER TUMOR TYPE)

Cetuximab

Lapatinib

Panitumumab

Ado-trastuzumab emtansine

Lapatinib

Pertuzumab

Trastuzumab

Trastuzumab-dkst

Cabozantinib

GENOMIC FINDINGS AND BIOMARKERS WITH NO REPORTABLE THERAPEUTIC OR CLINICAL TRIAL OPTIONS

For more information regarding biological and clinical significance, including prognostic, diagnostic, germline, and potential chemosensitivity implications, see the Genomic Alterations section.

CDKN2A/B loss	p. 5	NKX2-1 amplification	p. 7
MTAP loss	p. 6	TP53 R282G	p. 7
NFKBIA amplification	p. 6		

Note: Genomic alterations detected may be associated with activity of certain FDA approved drugs; however, the agents listed in this report may have varied clinical evidence in the patient's tumor type. Neither the therapeutic agents nor the trials identified are ranked in order of potential or predicted efficacy for this patient, nor are they ranked in order of level of evidence for this patient's tumor type.

SAMPLE

BIOMARKER FINDINGS

BIOMARKER

Microsatellite status

CATEGORY

MS-Stable

POTENTIAL TREATMENT STRATEGIES

On the basis of clinical evidence, MSS tumors are significantly less likely than MSI-H tumors to respond to anti-PD-1 immune checkpoint inhibitors¹⁻³, including approved therapies nivolumab and pembrolizumab⁴⁻⁵. In a retrospective analysis of 361 patients with solid tumors treated with pembrolizumab, 3% were MSI-H and experienced a significantly higher ORR compared with non-MSI-H cases (70% vs. 12%, $p=0.001$)⁶. Pembrolizumab therapy resulted in a significantly lower objective

response rate (ORR) in MSS colorectal cancer (CRC) compared with MSI-H CRC (0% vs. 40%)⁵. Similarly, a clinical study of nivolumab, alone or in combination with ipilimumab, in patients with CRC reported a significantly higher response rate in patients with MSI-H tumors than those without⁴.

FREQUENCY & PROGNOSIS

MSI-high (MSI-H) has been reported at various frequencies in non-small cell lung cancer (NSCLC) as well as in small cell lung cancer⁷⁻¹². One study observed MSI-H in 0.8% (4/480) of lung adenocarcinoma cases; the MSI-H tumors occurred in patients with smoking history, and 3/4 MSI-H cases had nonsynchronous carcinomas in other organs, although none of the patients were diagnosed with Lynch syndrome⁷.

FINDING SUMMARY

Microsatellite instability (MSI) is a condition of genetic hypermutability that generates excessive amounts of short insertion/deletion mutations in the genome; it generally occurs at microsatellite DNA sequences and is caused by a deficiency in DNA mismatch repair (MMR) in the tumor¹³. Defective MMR and consequent MSI occur as a result of genetic or epigenetic inactivation of one of the MMR pathway proteins, primarily MLH1, MSH2, MSH6, or PMS2¹³⁻¹⁵. The tumor seen here is microsatellite-stable (MSS), equivalent to the clinical definition of an MSS tumor: one with mutations in none of the tested microsatellite markers¹⁶⁻¹⁸. MSS status indicates MMR proficiency and typically correlates with intact expression of all MMR family proteins^{13,15,17-18}.

BIOMARKER

Tumor Mutation Burden

CATEGORY

TMB-Low (5 Muts/Mb)

POTENTIAL TREATMENT STRATEGIES

On the basis of emerging clinical evidence, increased TMB may be associated with greater sensitivity to immunotherapeutic agents, including anti-CTLA-4¹⁹, anti-PD-L1²⁰⁻²², and anti-PD-1 therapies^{5,23-24}; FDA-approved agents include ipilimumab, atezolizumab, avelumab, durvalumab, pembrolizumab, and nivolumab. In multiple solid tumor types, higher mutational burden has corresponded with response and improved prognosis. Pembrolizumab improved progression-free survival (14.5 vs. 3.4-3.7 months) in patients with non-small cell lung cancer (NSCLC) and higher mutational load (greater than 200 nonsynonymous mutations; hazard ratio = 0.19)²⁴. In studies of patients with either NSCLC or colorectal cancer (CRC), patients whose tumors harbor elevated mutational burden reported higher overall response rates to pembrolizumab^{5,23-24}. Anti-PD-1 therapies have achieved clinical benefit for certain patients with high mutational burden, including 3 patients with endometrial

adenocarcinoma who reported sustained partial responses following treatment with pembrolizumab²⁵ or nivolumab²⁶, a patient with hypermutant glioblastoma who obtained clinical benefit from pembrolizumab²⁷, and two pediatric patients with biallelic mismatch repair deficiency (bMMRD)-associated ultrahypermutant glioblastoma who experienced clinically and radiologically significant responses to nivolumab²⁸. In patients with melanoma, mutational load was associated with long-term clinical benefit from ipilimumab^{19,29} and anti-PD-1/anti-PD-L1 treatments²¹. For patients with metastatic urothelial carcinoma, those who responded to atezolizumab treatment had a significantly increased mutational load [12.4 mutations (mut) per megabase (Mb)] compared to nonresponders (6.4 muts/Mb)²⁰, and mutational load of 16 muts/Mb or higher was associated with significantly longer overall survival²².

FREQUENCY & PROGNOSIS

Low TMB is observed more commonly in non-small cell lung carcinomas (NSCLC) harboring known driver mutations (EGFR, ALK, ROS1, or MET) with the exception of BRAF or KRAS mutations, which are observed in approximately half of intermediate-high TMB cases³⁰. Although some studies have reported a lack of association between smoking and

mutational burden in NSCLC³¹⁻³³, several other large studies did find a strong association with increased TMB³⁴⁻³⁷. A large study of Chinese patients with lung adenocarcinoma reported a shorter median overall survival (OS) for tumors with a higher number of mutations in a limited gene set compared with lower mutation number (48.4 vs. 61.0 months)³².

FINDING SUMMARY

Tumor mutation burden (TMB, also known as mutation load) is a measure of the number of somatic protein-coding base substitution and insertion/deletion mutations occurring in a tumor specimen. TMB is affected by a variety of causes, including exposure to mutagens such as ultraviolet light in melanoma³⁸⁻³⁹ and cigarette smoke in lung cancer^{24,40}, mutations in the proofreading domains of DNA polymerases encoded by the POLE and POLD1 genes⁴¹⁻⁴⁵, and microsatellite instability (MSI)^{41,44-45}. The tumor seen here harbors a low TMB. Compared to patients with tumors harboring higher TMB levels, patients with tumors harboring low TMB levels have experienced lower rates of clinical benefit from treatment with immune checkpoint inhibitors, including anti-CTLA-4 therapy in melanoma¹⁹, anti-PD-L1 therapy in urothelial carcinoma²⁰, and anti-PD-1 therapy in non-small cell lung cancer and colorectal cancer^{5,24}.

GENOMIC FINDINGS

GENE
EGFR

ALTERATION
amplification, exon 19
deletion (T751_I759>S)

POTENTIAL TREATMENT STRATEGIES

EGFR activating mutations or amplification may predict sensitivity to EGFR inhibitors including erlotinib, gefitinib, afatinib, osimertinib, cetuximab, panitumumab, and lapatinib⁴⁶⁻⁵⁰. Other EGFR-targeted therapies are also in clinical trials. A Phase 2 trial of the pan-ERBB inhibitor dacomitinib in patients with lung adenocarcinoma reported 98% (44/45) disease control [partial response (PR) or stable disease], including a 76% PR rate, in patients with EGFR exon 19 deletions or the L858R mutation; lower disease control and PR rates were reported in patients with other EGFR mutations, wild-type EGFR, or unknown EGFR status⁵¹. Third-generation EGFR inhibitors, such as osimertinib or rociletinib, selectively target mutated EGFR including the EGFR resistance variant T790M. Osimertinib is

FDA approved to treat patients with EGFR T790M-positive advanced NSCLC and disease progression on EGFR inhibitor therapy⁴⁹. Necitumumab is an anti-EGFR antibody that is FDA approved for the treatment of metastatic squamous NSCLC in combination with gemcitabine and cisplatin. Addition of necitumumab increased overall and progression-free survival in patients with squamous NSCLC relative to chemotherapy alone; however, it exhibited a poor tolerability profile in non-squamous NSCLC, and EGFR expression has not been demonstrated to be predictive of clinical benefit in NSCLC⁵²⁻⁵³. HSP90 inhibitors have been clinically evaluated for patients with EGFR-mutated NSCLC⁵⁴⁻⁵⁸ and have shown activity against NSCLC with certain EGFR mutations⁵⁹. The reovirus Reolysin, which targets cells that harbor activated RAS signaling due to alterations in RAS genes or upstream activators such as EGFR⁶⁰⁻⁶², is also in clinical trials in some tumor types. Reolysin has demonstrated mixed clinical efficacy, with the highest rate of response reported for head and neck cancer⁶³⁻⁷¹.

FREQUENCY & PROGNOSIS

Amplification of EGFR has been reported in 6-42% of non-small cell lung carcinoma (NSCLC) samples⁷²⁻⁷⁵. EGFR mutations have been reported in 12-36% of lung adenocarcinoma samples, with amplification found in 7% of cases^{34,73,76}, and EGFR protein expression/overexpression has been reported in up to 70% of NSCLC tumors⁷⁷. EGFR mutations were shown to predict survival advantage for patients with Stage 1-3 resected lung adenocarcinoma or NSCLC⁷⁸⁻⁷⁹.

FINDING SUMMARY

EGFR encodes the epidermal growth factor receptor, which belongs to a class of proteins called receptor tyrosine kinases. In response to signals from the environment, EGFR passes biochemical messages to the cell that stimulate it to grow and divide⁸⁰. Amplification of EGFR has been associated with increased expression of EGFR mRNA and protein in several cancer types^{72,81-82}. The mutation seen here is a deletion in exon 19, encoding a portion of the kinase domain of EGFR; such mutations have been shown to activate EGFR kinase activity and to confer sensitivity to inhibitors such as erlotinib and gefitinib⁸³⁻⁸⁵.

GENE
ERBB3

ALTERATION
P1212S

POTENTIAL TREATMENT STRATEGIES

ERBB3/HER3 possesses a low-activity kinase domain and requires other ERBB family members for efficient signaling⁸⁶⁻⁸⁸. Therefore, ERBB3 amplification or activating mutation may predict sensitivity to therapies targeting ERBB2, such as pertuzumab, trastuzumab, ado-trastuzumab, lapatinib, and afatinib. In a study of afatinib monotherapy for patients with metastatic urothelial carcinoma, patients with ERBB3 mutation or ERBB2

amplification had significantly improved overall survival compared to patients without alterations (6.6 months vs. 1.4 months)⁸⁹. A patient with HER2-negative breast cancer harboring an activating ERBB3 mutation had a partial response to the combination of trastuzumab and lapatinib⁹⁰. In preclinical studies, cells with ERBB3 activating mutations were reported to be sensitive to anti-ERBB2 inhibition⁸⁶. Antibodies targeting ERBB3 are also being studied in clinical trials. However, as the mutation reported here has not been characterized, it is not known if these therapeutic approaches would be relevant.

FREQUENCY & PROGNOSIS

ERBB3 mutations have been reported in up to 1% of lung adenocarcinomas^{73,86}. ERBB3

protein expression has been reported in 18% (9/51) of lung adenocarcinomas⁹¹. High-level expression of ERBB3 mRNA has been associated with distant site metastases and poor overall survival in non-small cell lung cancer (NSCLC) patients⁹².

FINDING SUMMARY

ERBB3, which is also known as HER3, encodes a member of the epidermal growth factor receptor (EGFR) family⁹³. This ERBB3 mutation has not been characterized and its effect on ERBB3 function is unknown; however, mutation at this position has been reported previously in the context of cancer, which may indicate biological significance.

GENOMIC FINDINGS

GENE
MET

ALTERATION
T263M

POTENTIAL TREATMENT STRATEGIES

Strong evidence suggests that MET activation may predict sensitivity to targeted therapies⁹⁴ such as crizotinib, FDA approved for the treatment of ALK-positive NSCLC⁹⁵, and cabozantinib, FDA approved for the treatment of metastatic medullary thyroid cancer⁹⁶. Sensitivity to crizotinib is suggested by extensive clinical data in patients with MET-amplified cancers, including non-small cell lung cancer (NSCLC)^{97-98,99-101}, gastric cancer¹⁰², gastroesophageal cancer¹⁰³, glioblastoma¹⁰⁴, and carcinoma of unknown primary¹⁰⁵, as well as in patients with MET-mutated cancers, including NSCLC¹⁰⁶⁻¹¹⁰, renal cell carcinoma (RCC)¹¹¹ and histiocytic sarcoma¹⁰⁶, and in patients with NSCLC¹¹² and pulmonary sarcomatoid carcinoma¹¹³ whose tumors harbored both MET mutation and amplification. Sensitivity to cabozantinib is suggested by clinical benefit derived by a patient with MET-amplified and MET-mutated

NSCLC in one case report¹⁰⁷, as well as by extensive preclinical data¹¹⁴⁻¹²⁰. Strong clinical data also suggest sensitivity of MET-altered tumors to various other MET inhibitors, with examples including AMG 337 in gastric, esophageal, or gastroesophageal junction cancer¹²¹⁻¹²², volitinib in RCC¹²³, tepotinib in colorectal cancer¹²⁴, capmatinib in NSCLC¹⁰⁶, MGCD265 in NSCLC¹²⁵, PF-04217903 in RCC¹²⁶, and foretinib in RCC¹²⁷. Furthermore, the MET-targeting antibodies onartuzumab and MetMab have elicited responses for patients with MET-amplified NSCLC¹²⁸ or gastric cancer¹²⁹. In addition, high MET expression has been suggested to predict patient response to therapy regimens involving rilotumumab, a monoclonal antibody that targets the MET ligand HGF¹³⁰. However, in cases of uncharacterized alterations, such as seen here, it is unclear whether these therapeutic approaches would be relevant.

FREQUENCY & PROGNOSIS

In one study of 4402 lung adenocarcinoma cases, MET mutations (primarily those affecting MET exon 14 splicing) have been reported in ~3% of samples¹⁰⁶. In the TCGA datasets, MET mutation has been observed in 8.3% of lung adenocarcinomas and 2.1% of lung

squamous cell carcinomas⁷³⁻¹³¹. MET amplification has been reported in up to 11% of non-small-cell lung cancer (NSCLC) cases and has been found to increase following treatment with EGFR inhibitors^{74,132-134}. The prognostic implications of MET amplification, gain, and/or protein overexpression in NSCLC have yielded conflicting results, with some studies reporting no association and others finding a correlation with decreased survival and poor prognosis^{74,132,135-138}.

FINDING SUMMARY

MET encodes a receptor tyrosine kinase, also known as c-MET or hepatocyte growth factor receptor (HGFR), that is activated by the ligand HGF; MET activation results in signaling mediated partly by the RAS-RAF-MAPK and PI3K pathways to promote proliferation^{94,139}. Alterations such as reported here have not been characterized and are of unclear functional significance; however, similar alterations have been reported in the context of cancer, which may indicate biological relevance. Multiple MET activating alterations have exhibited clinical sensitivity to a variety of MET inhibitors in multiple cancer types^{106-111,126-127}.

GENE
CDKN2A/B

ALTERATION
loss

POTENTIAL TREATMENT STRATEGIES

Preclinical data suggest that tumors with loss of p16INK4a and p15INK4b function may be sensitive to CDK4/6 inhibitors, such as abemaciclib, ribociclib, and palbociclib¹⁴⁰⁻¹⁴³. Although case studies have reported that patients with breast cancer or uterine leiomyosarcoma harboring CDKN2A loss responded to palbociclib treatment¹⁴⁴⁻¹⁴⁵, multiple other clinical studies have shown no significant correlation between p16INK4a loss or inactivation and therapeutic benefit of these agents^{146-147,148-152}; it is not known whether CDK4/6 inhibitors would be

beneficial in this case. Although preclinical studies have suggested that loss of p14ARF function may be associated with reduced sensitivity to MDM2 inhibitors¹⁵³⁻¹⁵⁴, the clinical relevance of p14ARF as a predictive biomarker is not clear.

FREQUENCY & PROGNOSIS

CDKN2A/B loss or mutation has been reported in 19% and 4% of lung adenocarcinomas, respectively⁷³. Loss of p16INK4a protein expression, through CDKN2A mutation, homozygous deletion, or promoter methylation, has been described in 49-68% of non-small cell lung cancer (NSCLC) samples, whereas low p14ARF protein expression has been detected in 21-43% of NSCLC samples¹⁵⁵⁻¹⁵⁹. Loss of p16INK4a protein as well as CDKN2A promoter hypermethylation correlate with poor survival in patients with NSCLC^{156,160-162}.

FINDING SUMMARY

CDKN2A encodes two different, unrelated tumor suppressor proteins, p16INK4a and p14ARF, whereas CDKN2B encodes the tumor suppressor p15INK4b¹⁶³⁻¹⁶⁴. Both p15INK4b and p16INK4a bind to and inhibit CDK4 and CDK6, thereby maintaining the growth-suppressive activity of the Rb tumor suppressor; loss or inactivation of either p15INK4b or p16INK4a contributes to dysregulation of the CDK4/6-cyclin-Rb pathway and loss of cell cycle control^{155,165}. The tumor suppressive functions of p14ARF involve stabilization and activation of p53, via a mechanism of MDM2 inhibition¹⁶⁶⁻¹⁶⁷. This alteration is predicted to inactivate p16INK4a¹⁶⁸⁻¹⁷¹, p15INK4b¹⁷², and p14ARF¹⁷³⁻¹⁷⁴.

GENE MTAP

ALTERATION loss

POTENTIAL TREATMENT STRATEGIES

Inactivation of MTAP is being explored for specific metabolic vulnerabilities. In preclinical cancer models, MTAP inactivation showed increased sensitivity to inhibitors of purine synthesis or purine analogs, especially upon addition of exogenous MTA, which is converted to adenine in normal cells, providing competition to purine poisons lacking in MTAP-deficient cells¹⁷⁵⁻¹⁸³. However, such combination approaches are not being clinically tested, and a Phase 2 study of L-alanosine, an inhibitor of adenine synthesis, as a monotherapy in 65 patients with MTAP-deficient cancers reported no responses and stable disease in 24% of patients¹⁸⁴. Other

approaches have been described in preclinical studies¹⁸⁵⁻¹⁸⁷, but these have not been clinically tested.

FREQUENCY & PROGNOSIS

MTAP loss/homozygous deletion as well as loss of expression has been reported in a wide variety of solid tumors and hematologic cancers¹⁸⁸⁻¹⁸⁹; such events have been correlated with poor prognosis in a variety of cancer types, including hepatocellular carcinoma¹⁹⁰, gastrointestinal stromal tumors¹⁹¹, mantle cell lymphoma (MCL)¹⁹², melanoma¹⁹³⁻¹⁹⁴, gastric cancer¹⁹⁵, myxofibrosarcoma¹⁹⁶, nasopharyngeal carcinoma¹⁹⁷, ovarian carcinoma¹⁸⁸ and non-small cell lung cancer¹⁹⁸. MTAP loss was not prognostic in pediatric B-cell acute lymphocytic leukemia¹⁹⁹ or in astrocytoma²⁰⁰. However, MTAP has also been reported to be overexpressed in colorectal cancer (CRC) samples²⁰¹, and MTAP retention is thought to be important for prostate cancer growth due to

continuous supply of SAM²⁰². Germline SNPs in MTAP have been correlated with the development of cutaneous melanoma²⁰³⁻²⁰⁴, esophageal cancer²⁰⁵⁻²⁰⁶, osteosarcoma²⁰⁷, and CRC²⁰⁸.

FINDING SUMMARY

MTAP encodes S-methyl-5'-thioadenosine (MTA) phosphorylase, a tumor suppressor involved in polyamine metabolism and methionine synthesis, although its enzymatic function is dispensable for its tumor suppressor activity²⁰⁹⁻²¹⁰. Decreased expression of MTAP leads to MTA accumulation within tumor cells and their microenvironment^{190,211-212}, thereby reducing intracellular arginine methylation¹⁸⁵⁻¹⁸⁷ and altering cell signaling²¹²⁻²¹³. MTAP is located at 9p21, adjacent to CDKN2A and CDKN2B, with which it is frequently co-deleted in various cancers. Other alterations in MTAP are rare and have not been extensively characterized.

GENE NFKBIA

ALTERATION amplification

POTENTIAL TREATMENT STRATEGIES

There are no therapies that directly target NFKBIA amplification or expression.

FREQUENCY & PROGNOSIS

In the TCGA datasets, amplification of NFKBIA has been reported with the highest incidence in

lung adenocarcinoma (11.7%)⁷³, esophageal carcinoma (3.8%), uterine carcinosarcoma (3.6%), lung squamous cell carcinoma (3.4%), and ovarian serous cystadenocarcinoma (2.6%) (cBioPortal, 2017). Amplification or increased expression of NFKBIA in EGFR-mutant lung cancer has been reported to predict improved response to EGFR tyrosine kinase inhibitors²¹⁴⁻²¹⁵. Certain NFKBIA polymorphisms, which may affect IκBα expression levels, have been studied as risk factors for some cancer types, although the data are mixed and conflicting²¹⁶⁻²¹⁸.

FINDING SUMMARY

NFKBIA encodes IκBα, an inhibitor of the NF-κappaB (NFκB)/REL complex. It has been reported to act as a tumor suppressor in Hodgkin's lymphoma²¹⁹⁻²²³ and in glioblastoma^{216,224-225}. NFKBIA has been reported to be amplified in cancer²²⁶ and may be biologically relevant in this context²²⁷⁻²²⁸. In contrast, truncating mutations that result in loss of the majority of the IκBα protein are predicted to be inactivating.

GENE

NKX2-1

ALTERATION

amplification

POTENTIAL TREATMENT STRATEGIES

There are no approved therapies or trials that target tumors with TTF-1 amplification or overexpression. Lung cancer cell lines that express both TTF-1 and NKX2-1, which is located in the same amplicon as NKX2-1, have demonstrated resistance to cisplatin therapy²²⁹, although conflicting data has also been reported²³⁰.

FREQUENCY & PROGNOSIS

Putative amplification of NKX2-1 has been reported with the highest incidence in lung cancer, and has been observed in 14% of adenocarcinomas⁷³ and 5% of squamous cell carcinomas (SCC)¹³¹ as well as other tumor types including prostate adenocarcinomas (6%)²³¹, and poorly differentiated and anaplastic thyroid cancers (4%)²³². NKX2-1 mutation has been observed in 9% of acinar cell carcinomas of the pancreas²³³, 5% of uterine carcinosarcomas²³⁴, and is infrequent in other tumor types (cBioPortal, COSMIC, 2017). TTF-1 is expressed in a majority of lung adenocarcinomas and small cell carcinomas, as well as in a subset of thyroid and CNS

tumors²³⁵⁻²³⁷. Cytoplasmic TTF-1 expression has been reported as an adverse prognostic factor in breast carcinoma²³⁸⁻²³⁹. However, whether amplification and/or expression status of NKX2-1 have prognostic implications for patients with lung cancer is controversial^{229-230,240-243}. TTF-1 has been observed to have tumor-promoting as well as anti-oncogenic roles²⁴⁴⁻²⁴⁵.

FINDING SUMMARY

NKX2-1 (NK2 homeobox 1) encodes the thyroid transcription factor TTF-1²⁴⁶. Amplification of NKX2-1 results in overexpression of TTF-1 and upregulated transcription of downstream target genes²⁴⁷.

GENE

TP53

ALTERATION

R282G

POTENTIAL TREATMENT STRATEGIES

There are no approved therapies to address TP53 mutation or loss. However, tumors with TP53 loss of function alterations may be sensitive to the WEE1 inhibitor AZD1775²⁴⁸⁻²⁵¹, therapies that reactivate mutant p53 such as APR-246²⁵²⁻²⁵⁵, or p53 gene therapy and immunotherapeutics such as SGT-53²⁵⁶⁻²⁶⁰ and ALT-801²⁶¹. In a Phase 1 study, AZD1775 in combination with gemcitabine, cisplatin, or carboplatin elicited partial response in 10% (17/176) and stable disease in 53% (94/176) of patients with solid tumors; the response rate was 21% (4/19) in patients with TP53 mutations versus 12% (4/33) in patients who were TP53-wild-type²⁶². Combination of AZD1775 with paclitaxel and carboplatin achieved significantly longer progression-free survival than paclitaxel and carboplatin alone in patients with TP53-mutant ovarian cancer²⁶³. Furthermore, AZD1775 in combination with carboplatin achieved a 27% (6/22) response rate and 41% (9/22) stable disease rate in patients with TP53-mutant

ovarian cancer refractory or resistant to carboplatin plus paclitaxel²⁶⁴. In a Phase 1b trial in patients with p53-positive high-grade serous ovarian cancer, APR-246 combined with carboplatin and pegylated liposomal doxorubicin achieved a 52% (11/21) response rate and 100% disease control rate²⁵². In a Phase 1b clinical trial of SGT-53 in combination with docetaxel in patients with solid tumors, 75% (9/12) of evaluable patients experienced clinical benefit, including two confirmed and one unconfirmed partial responses and two instances of stable disease with significant tumor shrinkage²⁶⁰. Additionally, the combination of a CHK1 inhibitor and irinotecan reportedly reduced tumor growth and prolonged survival in a TP53 mutant, but not TP53 wild-type, breast cancer xenotransplant mouse model²⁶⁵. Kevetrin has also been reported to activate p53 in preclinical studies and might be relevant in the context of mutant p53²⁶⁶. Clinical trials of these agents are under way for some tumor types for patients with a TP53 mutation.

FREQUENCY & PROGNOSIS

TP53 is one of the most commonly mutated genes in lung cancer. TP53 mutations have been reported in 43-80% of non-small cell lung cancers (NSCLCs)^{73,131,158,267-271}. Mutations in TP53 have been associated with lymph node

metastasis in patients with lung adenocarcinoma²⁷². In one study of 55 patients with lung adenocarcinoma, TP53 alterations correlated with immunogenic features including PD-L1 expression, tumor mutation burden and neoantigen presentation; likely as a consequence of this association TP53 mutations correlated with improved clinical outcomes to PD-1 inhibitors pembrolizumab and nivolumab in this study²³.

FINDING SUMMARY

Functional loss of the tumor suppressor p53, which is encoded by the TP53 gene, is common in aggressive advanced cancers²⁷³. Any alteration that results in the disruption or partial or complete loss of the region encoding the TP53 DNA-binding domain (DBD, aa 100-292) or the tetramerization domain (aa 325-356), such as observed here, is thought to dysregulate the transactivation of p53-dependent genes and is predicted to promote tumorigenesis²⁷⁴⁻²⁷⁶. Germline mutations in TP53 are associated with the very rare disorder Li-Fraumeni syndrome and the early onset of many cancers²⁷⁷⁻²⁸². Estimates for the prevalence of germline TP53 mutations in the general population range from 1:5,000²⁸³ to 1:20,000²⁸², and in the appropriate clinical context, germline testing of TP53 is recommended.

Afatinib

Assay findings association

EGFR
ERBB3
P1212S

APPROVED INDICATIONS

Afatinib is an irreversible kinase inhibitor that targets the kinase domains of EGFR, ERBB2/HER2, and ERBB4. It is FDA approved for the first-line treatment of patients with metastatic non-small cell lung cancer (NSCLC) and nonresistant EGFR mutations and for the treatment of patients with metastatic, squamous NSCLC after progression on platinum-based chemotherapy.

GENE ASSOCIATION

Activation or amplification of EGFR may indicate sensitivity to afatinib. In Phase 2 studies of afatinib, patients with EGFR-amplified NSCLC achieved an objective response rate of 20% (5/25) and a disease-control rate of 64% (16/25)²⁸⁴, and 2/5 patients with EGFR amplification in other solid tumors experienced stable disease²⁸⁵. ERBB3/HER3 possesses a low-activity kinase domain and requires other ERBB family members for efficient signaling, ERBB2/HER2 in particular^{86-88,286}; therefore, ERBB3 amplification or activating mutations may indicate sensitivity to therapies such as afatinib. Partial response (PR) or stable disease (SD) was elicited in 5/7 patients with urothelial carcinoma harboring ERBB3 mutations (V104M, R103G, or G284R) and/or HER2 copy number gain treated with afatinib⁸⁹.

SUPPORTING DATA

Phase 3 clinical trials have demonstrated that treatment with afatinib, compared with chemotherapy, leads to significantly increased progression-free survival (PFS) for patients with EGFR-mutant NSCLC^{48,287} and increased

overall survival (OS) for patients with EGFR exon 19 alterations specifically²⁸⁸. A Phase 3 trial comparing afatinib with erlotinib as second-line therapies for advanced lung squamous cell carcinoma reported significantly higher OS (7.9 months vs. 6.8 months) and disease control rate (DCR) (51% vs. 40%) for patients treated with afatinib²⁸⁹. A Phase 2b trial comparing afatinib with gefitinib in patients with EGFR-mutant NSCLC reported similar OS (27.9 months vs. 24.5 months), but significantly higher time-to-treatment failure (13.7 months vs. 11.5 months), and overall response rate (73% vs. 56%)²⁹⁰. Phase 2/3 studies of afatinib treatment for patients with erlotinib- or gefitinib-resistant NSCLC have generally reported partial responses (PRs) of only 7-9%²⁹¹⁻²⁹⁶ and DCRs of more than 50%²⁹⁵; in particular, disease control was achieved for 2/2 patients with EGFR-amplified NSCLC²⁹⁵ and 9/14 patients with T790M-positive NSCLC²⁹⁶. The T790M mutation has been implicated in reduced response to afatinib^{294,297-299}, with a secondary T790M mutation reported in 48% (20/42) of patients with afatinib-resistant lung adenocarcinoma²⁹⁷. The combination of afatinib with cetuximab resulted in a higher response rate (29%) for patients with erlotinib- or gefitinib-resistant disease³⁰⁰, including T790M-positive cases³⁰⁰⁻³⁰¹, although adverse reactions may be a concern with this combination³⁰². Upon progression on afatinib, further benefit has been reported from combination treatment with afatinib and paclitaxel³⁰³ and with afatinib and cetuximab³⁰⁴.

Crizotinib

Assay findings association

MET
T263M

APPROVED INDICATIONS

Crizotinib is an inhibitor of the kinases MET, ALK, ROS1, and RON. It is FDA approved to treat patients with metastatic non-small cell lung cancer (NSCLC) whose tumors are positive for ALK rearrangements or ROS1 rearrangements.

GENE ASSOCIATION

Sensitivity of MET alterations to crizotinib is suggested by extensive clinical data in patients with MET-amplified cancers, including non-small cell lung cancer (NSCLC)^{97-98,99-101}, gastric cancer¹⁰², gastroesophageal cancer¹⁰³, glioblastoma¹⁰⁴, and carcinoma of unknown primary¹⁰⁵, as well as in patients with MET-mutated cancers, including NSCLC¹⁰⁶⁻¹¹⁰, renal cell carcinoma (RCC)¹¹¹ and histiocytic sarcoma¹⁰⁶. However, it is not known whether this therapeutic approach would be relevant in the context of alterations that have not been fully characterized, as seen here.

SUPPORTING DATA

Clinical data indicate activity of crizotinib in MET-activated NSCLC, including lung adenocarcinoma and lung squamous cell carcinoma. In two ongoing studies of crizotinib (AcSé, NCT02034981 and PROFILE 1001, NCT00585195), 4 partial responses (PRs) and 4 stable diseases (SDs) were reported in 14 evaluable patients⁹⁷, and 4 PRs were reported in 12 evaluable patients³⁰⁵ with

MET-amplified NSCLC. Additional patients with MET-amplified NSCLC have been reported to experience clinical benefit from crizotinib in several case studies^{99-101,306-308}. A patient with lung adenocarcinoma harboring K860I and L858R EGFR mutations, who acquired both EGFR T790M and MET-amplification upon various treatments, experienced clinical benefit from subsequent combination treatment of osimertinib and crizotinib; this combination regimen was well tolerated³⁰⁹. A case report of a patient with chemotherapy-refractory, pulmonary sacomatoid carcinoma with a MET exon 14 splice site alteration and amplification experienced a partial response to crizotinib treatment¹¹³. In an ongoing expansion cohort of the PROFILE 1001 study, patients with NSCLC harboring MET exon 14 alterations experienced an overall response rate of 44% to crizotinib, including 8 confirmed PRs and 9 SDs out of 18 patients; treatment duration ranged from 0.5 to 12.2+ months, with 76% (16/21) of patients still on study³¹⁰. Several case studies have also reported response to crizotinib in NSCLC with MET exon 14 alterations, with or without concomitant MET amplification^{107-110,112,306,311-313}. In patients with NSCLC and MET overexpression with or without gene amplification, crizotinib elicited 11 PRs and 3 SDs in 19 evaluable patients⁹⁸. However, as the mutation reported here has not been characterized, it is not known if this therapeutic approach would be relevant.

Erlotinib

Assay findings association

EGFR
amplification, exon 19 deletion
(T751_I759>S)

APPROVED INDICATIONS

Erlotinib is a small molecule inhibitor of EGFR. It is FDA approved both as first-line and maintenance therapy, as well as second- or greater line of treatment after chemotherapy failure, for patients with metastatic non-small cell lung cancer (NSCLC) harboring EGFR exon 19 deletions or exon 21 (L858R) mutations. Erlotinib is also FDA approved in combination with gemcitabine as a first-line treatment for advanced pancreatic cancer.

GENE ASSOCIATION

Amplification or activation of EGFR may predict sensitivity to therapies such as erlotinib. In a prospective study of advanced NSCLC treated with gefitinib (n=102), EGFR copy gain was significantly associated with improved survival [hazard ratio (HR)=0.44]³¹⁴. Several meta-analyses spanning 14 to 20 studies of patients with advanced NSCLC receiving single-agent erlotinib or gefitinib (n=1725 to 1854) reported the association of increased EGFR copy number with improved overall survival (HR=0.72 to 0.77), although the survival benefit was not observed for East Asian populations (HR=0.79 to 1.11)³¹⁵⁻³¹⁷.

SUPPORTING DATA

The initial approval of erlotinib in NSCLC was based on the BR.21 Phase 3 randomized trial demonstrating prolonged overall survival for unselected patients with NSCLC treated with erlotinib compared with standard chemotherapy³¹⁸. Furthermore, several randomized Phase 3 trials have shown a significant improvement in response and progression-free survival for erlotinib compared with combination chemotherapy in patients with known EGFR mutations. This includes the EURTAC trial of erlotinib versus platinum-based chemotherapy as first-line treatments⁴⁶ and the SATURN trial of erlotinib as maintenance therapy following first-line platinum-based chemotherapy³¹⁹. On the other hand, the efficacy of erlotinib for patients lacking the common EGFR activating alterations (exon 19 deletion or L858R mutation) may be regimen-dependent. For patients with NSCLC and wild-type EGFR, chemotherapy was found to be more effective than erlotinib as first-, second-, or third-line treatment³²⁰⁻³²². However, as maintenance therapy, erlotinib reduced risk for progression compared with placebo by 19% (hazard ratio = 0.81)³²². The single-arm, Phase IV TRUST trial for genomically unselected patients with advanced NSCLC who failed on, or were unsuitable for, chemotherapy or who were ineligible for erlotinib clinical trials reported a disease control rate of 69%³²³.

THERAPIES WITH CLINICAL BENEFIT

IN PATIENT'S TUMOR TYPE

Gefitinib

Assay findings association

EGFR

amplification, exon 19 deletion (T751_I759>S)

APPROVED INDICATIONS

Gefitinib targets the tyrosine kinase EGFR and is FDA approved to treat non-small cell lung cancer (NSCLC) harboring exon 19 deletions or exon 21 (L858R) substitution mutations in EGFR.

GENE ASSOCIATION

Amplification or activation of EGFR may predict sensitivity to therapies such as gefitinib. In a prospective study of advanced NSCLC treated with gefitinib (n=102), EGFR copy gain was significantly associated with improved survival [hazard ratio (HR)=0.44]³¹⁴. Several meta-analyses spanning 14 to 20 studies of patients with advanced NSCLC receiving single-agent erlotinib or gefitinib (n=1725 to 1854) reported the association of increased EGFR copy number with improved overall

survival (HR=0.72 to 0.77), although the survival benefit was not observed for East Asian populations (HR=0.79 to 1.11)³¹⁵⁻³¹⁷.

SUPPORTING DATA

Gefitinib achieved an objective response rate of 69.8% and an overall survival of 19.2 months as first-line treatment of Caucasian patients with NSCLC and EGFR sensitizing mutations, which were mostly EGFR exon 19 deletions and EGFR L858R⁴⁷. In the retrospective analysis of a Phase 3 study in Asia, gefitinib increased progression-free survival in a subgroup of patients with EGFR mutation-positive NSCLC as compared with carboplatin/paclitaxel doublet chemotherapy (hazard ratio for progression 0.48)³²⁴⁻³²⁵.

Osimertinib

Assay findings association

EGFR

amplification, exon 19 deletion (T751_I759>S)

APPROVED INDICATIONS

Osimertinib is an irreversible EGFR tyrosine kinase inhibitor (TKI) that is selective for EGFR TKI-sensitizing mutations and the EGFR T790M mutation. It is FDA approved to treat patients with metastatic EGFR T790M-positive non-small cell lung cancer (NSCLC) and disease progression on or after EGFR TKI therapy.

GENE ASSOCIATION

EGFR TKI-sensitizing mutations and/or the EGFR T790M mutation may predict sensitivity to osimertinib^{49,326}. T790M-positive patients showed higher response rates than T790M-negative cases in a Phase 1 study for patients with acquired EGFR TKI resistance (61% vs. 21%)⁴⁹. Although tumors with EGFR amplification may not be sensitive to osimertinib, which selectively targets mutated EGFR, preclinical data indicate sensitivity of various activating EGFR alterations to osimertinib³²⁶.

SUPPORTING DATA

Osimertinib has been studied primarily for the treatment of EGFR-mutated NSCLC. In a Phase 3 study for patients with EGFR T790M-positive advanced NSCLC who progressed on EGFR tyrosine kinase inhibitor (TKI) therapy, osimertinib compared with combination platinum therapy led to longer median progression-free

survival (PFS; 10.1 months vs. 4.4 months), including for patients with metastases to the central nervous system (CNS; 8.5 months vs. 4.2 months). An objective response rate (ORR) of 71% was achieved with osimertinib compared to 31% with combination platinum therapy³²⁷. Earlier phase studies confirmed the efficacy of osimertinib in this setting^{49,328-329}. In contrast, T790M-negative patients with acquired EGFR TKI resistance experienced an ORR of 21% and median PFS of 2.8 months⁴⁹. As first-line therapy for advanced NSCLC with activating, sensitizing EGFR mutations, osimertinib compared with erlotinib or gefitinib significantly increased median PFS (18.9 vs. 10.2 months, hazard ratio of 0.46), including for patients with CNS metastases, and showed a superior toxicity profile³³⁰. First-line efficacy of osimertinib is supported by Phase 1 data (ORR of 77% and median PFS of 20.5 months across doses)³³¹. A Phase 1b study combined osimertinib with the immunotherapy durvalumab, MEK inhibitor selumetinib, or MET inhibitor savolitinib and observed partial responses (PRs) for each of the combinations (9/14 PRs with durvalumab, 9/23 PRs with selumetinib, 6/11 PRs with savolitinib)³³². Case studies report that 2 patients with T790M-mutated NSCLC achieved durable PRs to osimertinib rechallenge after the adverse events induced by initial osimertinib treatment had been resolved³³³⁻³³⁴.

Ado-trastuzumab emtansine

Assay findings association

ERBB3
P1212S

APPROVED INDICATIONS

Ado-trastuzumab emtansine (T-DM1) is an antibody-drug conjugate that targets the protein ERBB2/HER2 on the cell surface, inhibiting HER2 signaling³³⁵⁻³³⁶; it also releases the cytotoxic therapy DM1 into cells, leading to cell death³³⁶⁻³³⁷. T-DM1 is FDA approved for the treatment of HER2-positive (HER2+), metastatic breast cancer.

GENE ASSOCIATION

ERBB3/HER3 possesses a low-activity kinase domain and requires other ERBB family members for efficient signaling, ERBB2/HER2 in particular^{86-88,286}; therefore, ERBB3 amplification or activating mutations may indicate sensitivity to therapies such as T-DM1. However, it is not known whether this therapeutic approach would be relevant in the context of alterations that have not been fully characterized, as seen here.

SUPPORTING DATA

A patient with non-small cell lung cancer, disease progression on two prior lines of chemotherapy, and an activating ERBB2 alteration (exon 20 insertion)

experienced a rapid and durable response to T-DM1³³⁸⁻³³⁹. The vast majority of data on the therapeutic use of T-DM1 has been collected in the context of breast cancer, although clinical trials investigating T-DM1 are underway in several tumor types, primarily in HER2+ cancers. A Phase 3 trial in 602 patients with HER2+ breast cancer reported that those who received T-DM1 showed an improved progression-free survival (PFS) and a lower rate of adverse events than patients who received the physician's choice of therapy³⁴⁰. A second Phase 3 trial in 991 patients with HER2+ breast cancer reported that T-DM1 brought about significantly longer overall survival (OS) and PFS, as compared with lapatinib plus capecitabine, in patients previously treated with trastuzumab plus a taxane³⁴¹⁻³⁴². Two separate Phase 2 trials reported robust activity for single-agent T-DM1 as a treatment for HER2+ metastatic breast cancer in patients previously treated with standard HER2-directed therapies or HER2-directed therapies plus chemotherapy, with objective response rates of 34.5% and 25.9%, respectively, and PFS of 6.9 months and 4.9 months, respectively³⁴³⁻³⁴⁴.

Cabozantinib

Assay findings association

MET
T263M

APPROVED INDICATIONS

Cabozantinib inhibits multiple tyrosine kinases, including MET, RET, VEGFRs, and ROS1. It is FDA approved to treat advanced renal cell carcinoma (RCC), after prior anti-angiogenic therapy, and progressive, metastatic medullary thyroid cancer (MTC).

GENE ASSOCIATION

Sensitivity of MET alterations to cabozantinib is suggested by clinical responses in patients with non-small cell lung cancer (NSCLC) harboring MET mutations associated with MET exon 14 skipping, with or without concurrent MET amplification^{107,345}, as well as by extensive preclinical data¹¹⁴⁻¹²⁰. However, it is not known whether this therapeutic approach would be relevant in the context of alterations that have not been fully characterized, as seen here.

SUPPORTING DATA

Early clinical data report promising activity of cabozantinib in non-small cell lung carcinoma (NSCLC) harboring RET fusions or MET alterations. A Phase 2 study of cabozantinib in RET-rearranged lung adenocarcinoma reported an objective response rate (ORR) of 28%, with 7 partial responses (PRs) in 25 evaluable patients, median progression-free survival (PFS) of 6 months and overall survival of 10 months³⁴⁶⁻³⁴⁷. In

a retrospective analysis of patients with RET-rearranged NSCLC treated with various RET inhibitors, of the 14 patients treated with cabozantinib, 1 complete response (CR), 3 PRs, and 4 stable diseases (SDs) were reported, for an ORR of 31% and a disease control rate of 62%³⁴⁸. Additional studies of single-agent cabozantinib have reported 2 PRs and 1 SD in a series of 5 patients with RET-rearranged lung adenocarcinoma³⁴⁹, a CR in a patient with lung adenocarcinoma harboring MET amplification and a mutation associated with MET exon 14 skipping¹⁰⁷, and intracranial activity of cabozantinib in a patient with MET-mutated NSCLC without co-occurring MET amplification who had previously progressed on crizotinib³⁴⁵. In genomically unselected patients with metastatic NSCLC, a Phase 2 randomized discontinuation trial of cabozantinib in a heavily pretreated cohort reported PRs in 10% (6/60) of patients, tumor regression in 65% (31/48) of patients, a median PFS of 4.2 months, and a safety profile similar to that of other tyrosine kinase inhibitors³⁵⁰. In patients with EGFR wild-type non-squamous NSCLC who have progressed after previous treatment, patients treated with cabozantinib alone or in combination with erlotinib experienced a longer median PFS (4.3 months and 4.7 months, respectively) compared to single agent erlotinib (1.8 months)³⁵¹.

Cetuximab

Assay findings association

EGFR

amplification, exon 19 deletion (T751_I759>S)

APPROVED INDICATIONS

Cetuximab is a monoclonal antibody that targets EGFR. It is FDA approved for the treatment of head and neck squamous cell carcinoma (HNSCC) and KRAS wild-type metastatic colorectal cancer (CRC).

GENE ASSOCIATION

EGFR activating mutations or amplification may confer sensitivity to EGFR inhibitory antibodies such as cetuximab. For patients with metastatic CRC receiving cetuximab or panitumumab as mono- or combination therapy, increased EGFR copy number associated with improved overall survival (hazard ratio = 0.62) in a meta-analysis, although increased survival was not seen in populations that received first-line treatment with EGFR antibodies³⁵². In HNSCC, however, EGFR copy number did not associate with the efficacy of cetuximab plus chemotherapy³⁵³.

SUPPORTING DATA

In previously untreated patients with non-small cell lung cancer (NSCLC), the FLEX study demonstrated that in NSCLC tumors with high expression of EGFR, treatment with cetuximab plus chemotherapy resulted in longer overall survival compared to chemotherapy alone⁵⁰. There was no clear association between cetuximab

response and EGFR mutations in the FLEX trial⁵⁰. In a Phase 2 study of 31 patients with Stage 3 NSCLC, addition of cetuximab to radiotherapy and chemotherapy produced an overall response rate of 67%; EGFR gene copy number was not predictive of efficacy outcome³⁵⁴. A Phase 3 study of 938 patients with progressive NSCLC after platinum-based therapy concluded that, in unselected patients, the addition of cetuximab to chemotherapy was not recommended in this second-line setting³⁵⁵. Cetuximab is also being studied as part of a therapeutic regimen for patients with EGFR mutations who develop secondary resistance to erlotinib or gefitinib. A Phase 1b study combining afatinib and the anti-EGFR antibody cetuximab in patients with advanced EGFR-mutant lung cancer with acquired resistance to erlotinib/ gefitinib observed an overall objective response rate of 29%, and comparable response rates in both T790M-positive and T790M-negative tumors (32% vs. 25%)³⁰⁰. A Phase 1 study of combination erlotinib and cetuximab treatment in patients with NSCLC, including those with squamous tumors, inhibitor-resistant EGFR mutations, and wild-type EGFR, as well as those who had progressed on prior erlotinib treatment, reported partial responses in two of 20 patients and stable disease lasting at least 6 months in three of 20 patients³⁵⁶.

Lapatinib

Assay findings association

EGFR

ERBB3

P1212S

APPROVED INDICATIONS

Lapatinib is a tyrosine kinase inhibitor that targets EGFR, ERBB2/HER2, and to a lesser degree, ERBB4. It is FDA approved in combination with capecitabine or letrozole for the treatment of HER2-overexpressing (HER2+) metastatic breast cancer.

GENE ASSOCIATION

Activation or amplification of EGFR may predict sensitivity to lapatinib. However, a Phase 2 study of lapatinib in non-small cell lung cancer (NSCLC) did not observe any responses for five patients with EGFR amplification³⁵⁷. ERBB3/HER3 possesses a low-activity kinase domain and requires other ERBB family members for efficient signaling, ERBB2/HER2 in particular^{86-88,286}; therefore, ERBB3 amplification or activating mutations may indicate sensitivity to therapies such as lapatinib⁸⁶. A patient with HER2-negative breast

cancer harboring an activating ERBB3 mutation had a partial response to the combination of trastuzumab and lapatinib⁹⁰.

SUPPORTING DATA

Investigations into the efficacy of lapatinib have primarily been in the context of breast cancer. In preclinical assays, lapatinib reduced cell proliferation in vitro and reduced the number and size of tumors in mouse xenograft models of EGFR- and ERBB2-amplified NSCLC cells³⁵⁸. A Phase 1 study of single-agent lapatinib included 9 unselected patients with lung cancer and reported 1 case of prolonged stable disease³⁵⁹. In a Phase 2 trial in patients with advanced or metastatic NSCLC, lapatinib monotherapy did not result in significant tumor reduction, but further investigation of lapatinib in combination with other therapies may be warranted³⁵⁷.

THERAPIES WITH CLINICAL BENEFIT

IN OTHER TUMOR TYPE

Panitumumab

Assay findings association

EGFR

amplification, exon 19 deletion (T751_I759>S)

APPROVED INDICATIONS

Panitumumab is a monoclonal antibody that targets EGFR. It is FDA approved for the treatment of KRAS wild-type metastatic colorectal cancer (CRC).

GENE ASSOCIATION

EGFR activating mutations or amplification may confer sensitivity to EGFR inhibitory antibodies such as panitumumab. For patients with metastatic CRC receiving cetuximab or panitumumab as mono- or combination therapy, increased EGFR copy number associated with improved overall survival (hazard ratio = 0.62) in a meta-analysis, although increased survival was not seen in

populations that received first-line treatment with EGFR antibodies³⁵².

SUPPORTING DATA

In a Phase 2 trial of advanced non-small cell lung cancer (NSCLC), the addition of panitumumab to paclitaxel/carboplatin did not result in improved clinical benefit³⁶⁰, and subsequent studies investigating the addition of panitumumab to pemetrexed/cisplatin reported no benefit for patients with wild-type KRAS lung adenocarcinoma³⁶¹. The combination of afatinib and panitumumab has been explored for 2 patients with EGFR T790M NSCLC, with 1 partial response reported³⁶².

Pertuzumab

Assay findings association

ERBB3

P1212S

APPROVED INDICATIONS

Pertuzumab is a monoclonal antibody that interferes with the interaction between HER2 and ERBB3. It is FDA approved in combination with trastuzumab and docetaxel to treat patients with HER2-positive (HER2+) metastatic breast cancer who have not received prior chemotherapy or HER2-targeted therapy. It is also approved in combination with trastuzumab and chemotherapy as neoadjuvant treatment for HER2+, locally advanced, inflammatory, or early stage breast cancer and as adjuvant treatment for patients with HER2+ early breast cancer at high risk of recurrence.

GENE ASSOCIATION

ERBB3 amplification or activating mutations may predict sensitivity to pertuzumab. However, it is not known whether this therapeutic approach would be relevant in the context of alterations that have not been fully characterized, as seen here.

SUPPORTING DATA

Pertuzumab received FDA approval based on a Phase 3 randomized study that demonstrated significant improvement in progression-free survival (PFS), with a trend toward improvement in overall survival (OS), for the

combination of pertuzumab, trastuzumab, and docetaxel, as compared to treatment with trastuzumab and docetaxel alone, for patients with HER2-positive breast cancer³⁶²⁻³⁶³. In a Phase 1 study of pertuzumab in advanced cancer, 2/19 patients reported partial responses and 6/19 patients reported stable disease after two cycles, including one patient with lung cancer³⁶⁴. In another Phase 1 study in Japanese patients with solid tumors, no responses were observed and stable disease was reported in 1 of 7 patients with NSCLC³⁶⁵. In a Phase 2 study of pertuzumab in NSCLC, no responses were observed and the progression-free survival was 6.1 weeks³⁶⁶. Phase 1 and 2 trials of pertuzumab in combination with erlotinib in NSCLC have reported a response rate of 20% (3/15, 2 of the responders had mutant EGFR)³⁶⁷; a reduction in circulating tumor cells was noted and correlated with reduction in tumor size³⁶⁸. In a Phase 2 study of pertuzumab plus erlotinib in relapsed patients with NSCLC, PET-CT imaging showed that the primary endpoint of response rate (RR) was met in 19.5% of all patients (n = 41) and in 8.7% of patients with wild-type EGFR NSCLC (n = 23); however, 68.3% (28/41) of patients showed treatment-related grade 3 (or higher) adverse events³⁶⁹.

THERAPIES WITH CLINICAL BENEFIT

IN OTHER TUMOR TYPE

Trastuzumab

Assay findings association

ERBB3
P1212S

APPROVED INDICATIONS

Trastuzumab is a monoclonal antibody that targets the protein ERBB2/HER2. It is FDA approved as monotherapy and in combination with other therapies for HER2-positive (HER2+) metastatic and early breast carcinoma and in combination with chemotherapy for HER2+ metastatic gastric or gastroesophageal adenocarcinoma.

GENE ASSOCIATION

ERBB3/HER3 possesses a low-activity kinase domain and requires other ERBB family members for efficient signaling, ERBB2/HER2 in particular^{86-88,286}; therefore, ERBB3 amplification or activating mutations may indicate sensitivity to therapies such as trastuzumab⁸⁶. A patient with HER2-negative breast cancer harboring an activating ERBB3 mutation had a partial response to the combination of trastuzumab and lapatinib⁹⁰. However, it is not known whether this therapeutic approach would be relevant in the context of alterations that have not been fully characterized, as seen here.

SUPPORTING DATA

A Phase 2 clinical trial of docetaxel with trastuzumab in non-small cell lung cancer (NSCLC) reported partial responses in 8% of patients; response did not correlate with HER2 status as assessed by immunohistochemistry³⁷⁰. Another Phase 2 study of 169 patients with NSCLC reported an objective response rate of 23% (7/30 patients) in the patients treated with a combination therapy of docetaxel and trastuzumab, and 32% (11/34) in patients treated with paclitaxel and trastuzumab³⁷¹. HER2 expression did not impact the results of this study³⁷¹. In a Phase 2a umbrella basket study, trastuzumab plus pertuzumab elicited objective responses for 2/7 patients with NSCLC and ERBB2 amplification or overexpression³⁷². A patient with lung adenocarcinoma that was HER-positive by FISH and harbored an ERBB2 G776L mutation experienced a partial response on trastuzumab and paclitaxel³⁷³. In a retrospective analysis of patients with NSCLC harboring ERBB2 exon 20 insertion mutations, disease control was reported in 93% of patients (13/14) treated with trastuzumab in combination with chemotherapy³⁷⁴.

Trastuzumab-dkst

Assay findings association

ERBB3
P1212S

APPROVED INDICATIONS

Trastuzumab-dkst is FDA approved as a biosimilar therapy to trastuzumab. Trastuzumab-dkst is a monoclonal antibody that targets the protein ERBB2/HER2, and is FDA approved as monotherapy and in combination with chemotherapy for HER2-positive (HER2+) metastatic and early breast carcinoma and in combination with chemotherapy for HER2+ metastatic gastric or gastroesophageal junction adenocarcinoma.

GENE ASSOCIATION

ERBB3 possesses a low-activity kinase domain and requires other ERBB family members for efficient signaling, HER2 in particular^{86-88,286}; therefore, ERBB3 amplification or activating mutations may indicate sensitivity to anti-HER2 therapies such as trastuzumab-dkst⁸⁶. A patient with HER2-negative breast cancer

harboring an activating ERBB3 mutation had a partial response to the combination of trastuzumab and lapatinib⁹⁰. However, it is not known if this therapeutic approach would be relevant in the context of alterations that have not been fully characterized, as seen here.

SUPPORTING DATA

The Phase 3 Heritage study demonstrated comparable 24-week objective response rates (69.6% vs. 64.0%) and progression-free survival for patients with treatment-naïve HER2+ metastatic breast cancer treated with either trastuzumab-dkst or trastuzumab in combination with taxane³⁷⁵. In both patients with HER2+ breast cancer and in healthy adults, trastuzumab-dkst demonstrated comparable pharmacokinetic, safety, and immunomodulation profiles to trastuzumab^{375-376 377}.

Note: Genomic alterations detected may be associated with activity of certain FDA approved drugs, however the agents listed in this report may have little or no evidence in the patient's tumor type.

IMPORTANT Clinical trials are ordered by gene and prioritized by: age range inclusion criteria for pediatric patients, proximity to ordering medical facility, later trial phase, and verification of trial information within the last two months. While every effort is made to ensure the accuracy of the information contained below, the information available in the public domain

is continually updated and should be investigated by the physician or research staff. This is not a comprehensive list of all available clinical trials. Foundation Medicine displays a subset of trial options and ranks them in this order of descending priority: Qualification for pediatric trial → Geographical proximity → Later trial phase. Clinical trials listed here

may have additional enrollment criteria that may require medical screening to determine final eligibility. For additional information about listed clinical trials or to conduct a search for additional trials, please see [clinicaltrials.gov](https://www.clinicaltrials.gov). Or, visit <https://www.foundationmedicine.com/genomic-testing#support-services>.

GENE

EGFR

ALTERATION

**amplification, exon 19 deletion
(T751_I759>S)**

RATIONALE

EGFR amplification or activating mutations may predict sensitivity to EGFR-targeted therapies. Several strategies to circumvent resistance are under investigation, including irreversible EGFR tyrosine kinase inhibitors and the use of HSP90 inhibitors. Examples of clinical trials that may be appropriate for this patient are listed below. These trials were identified through a search of the trial

website [clinicaltrials.gov](https://www.clinicaltrials.gov) using keyword terms such as "EGFR", "cetuximab", "panitumumab", "erlotinib", "gefitinib", "lapatinib", "afatinib", "osimertinib", "BIBW 2992", "CO-1686", "AZD9291", "PF-00299804", "HSP90", "reolysin", "NSCLC", "lung", "solid tumor", and/or "advanced cancer".

NCT02193282

PHASE 3

Randomized Double Blind Placebo Controlled Study of Erlotinib or Placebo in Patients With Completely Resected Epidermal Growth Factor Receptor (EGFR) Mutant Non-Small Cell Lung Cancer (NSCLC)

TARGETS
EGFR

LOCATIONS: Kentucky, Tennessee, New Jersey, Alaska, Delaware, North Dakota, Montana, Ohio, Rhode Island, Maine, New Hampshire, Arizona, Wyoming, Hawaii, Georgia, South Carolina, West Virginia, Connecticut, Vermont, California, Michigan, Alabama, Nebraska, Texas, Arkansas, Nevada, Idaho, Washington, Iowa, Mississippi, Missouri, Wisconsin, Colorado, Louisiana, Virginia, South Dakota, New York, Oklahoma, Utah, Massachusetts, Florida, Maryland, Oregon, Indiana, Kansas, District of Columbia, Illinois, Minnesota, North Carolina, Pennsylvania, New Mexico

NCT02511106

PHASE 3

A Phase III, Double-blind, Randomized, Placebo-controlled Multi-centre, Study to Assess the Efficacy and Safety of AZD9291 Versus Placebo, in Patients With Epidermal Growth Factor Receptor Mutation Positive Stage IB-IIIA Non-small Cell Lung Carcinoma, Following Complete Tumour Resection With or Without Adjuvant Chemotherapy (ADAURA).

TARGETS
EGFR

LOCATIONS: Maryland, Guangzhou (China), Rio Grande do Sul (Brazil), Beer-Sheva (Israel), Hoofddorp (Netherlands), Vinnytsia (Ukraine), Nanjing (China), Taichung (Taiwan), Hamburg (Germany), Poznan (Poland), Kurralta Park (Australia), Iasi (Romania), Beijing (China), Saint-Petersburg (Russian Federation), Ankara (Turkey), Virginia, Milano (Italy), Seoul (Korea, Republic of), Muang (Thailand), Dnipropetrovsk (Ukraine), Fortaleza (Brazil), Zaporizhzhya (Ukraine), Limoges (France), Brussels (Belgium), Gent (Belgium), St. Petersburg (Russian Federation), Halle (Germany), Budapest (Hungary), New Jersey, Bron (France), Lugo (Spain), Changchun (China), Saint Petersburg (Russian Federation), Lyon (France), Curitiba (Brazil), Hawaii, Hangzhou (China), Zaragoza (Spain), Camperdown (Australia), Bergamo (Italy), Yonago-shi (Japan), Großhansdorf (Germany), Chiayi (Taiwan), Roma (Italy), Tainan (Taiwan), Łódź (Poland), Paris (France), Illinois, Otwock (Poland), Colorado, Immenhausen (Germany), Bedford Park (Australia), São José do Rio Preto (Brazil), Aachen (Germany), Arnhem (Netherlands), Washington, Kashiwa-shi (Japan), Taipei (Taiwan), Parma (Italy), Racibórz (Poland), San Sebastian (Spain), Shinjuku-ku (Japan), Liou Ying Township (Taiwan), Ho Chi Minh (Vietnam), Sagami-hara-shi (Japan), Bunkyo-ku (Japan), Málaga (Spain), Sunto-gun (Japan), Kanazawa (Japan), Kogarah (Australia), Timisoara (Romania), Sankt-Peterburg (Russian Federation), Ube-shi (Japan), Suwon-si (Korea, Republic of), Suwon (Korea, Republic of), Yangzhou (China), California, Livorno (Italy), Changhua (Taiwan), Xi'an (China), Székesfehérvár (Hungary), Sakai-shi (Japan), Osakasayama-shi (Japan), Khonkaen (Thailand), Izmir (Turkey), Meldola (Italy), Urumqi (China), Sendai-shi (Japan), Matsuyama-shi (Japan), Yokohama-shi (Japan), Sasebo-shi (Japan), Bruxelles (Belgium), Phitsanulok (Thailand), Lucca (Italy), Gauting (Germany), Homburg (Germany), Ryazan (Russian Federation), Gerlingen (Germany), Suzhou (China), Linköping (Sweden), Lübeck (Germany), Arkhangelsk (Russian Federation), Chom Thon (Thailand), Tennessee, Hanoi (Vietnam), São Paulo (Brazil), Hirakata-shi (Japan), Hiroshima-shi (Japan), Focsani (Romania), Bangkok (Thailand), Tianjin (China), Coswig (Germany), Salvador (Brazil), Ottignies (Belgium), Urumchi (China), Brussels (Anderlecht) (Belgium), Torokbalint (Hungary), Bologna (Italy), Pyatigorsk (Russian Federation), Rhode Island, Majadahonda (Spain), Uzhgorod (Ukraine), Valencia (Spain), Porto Alegre (Brazil), Florida, Hong Kong (Hong Kong), Georgia, Cheongju-si (Korea, Republic of), Tel Hashomer (Israel), Las Palmas de Gran Canaria (Spain), A Coruña (Spain), Köln (Germany), Eindhoven (Netherlands), Kassel (Germany), Shanghai (China), Santa Catarina (Brazil), Madrid (Spain), Florianópolis (Brazil), Cachoeira De Itapemirim (Brazil), Lille cedex (France), Bursa (Turkey), Bari (Italy), Nagoya-shi (Japan), Palermo (Italy), Heidelberg (Australia), Lviv (Ukraine), Songkla (Thailand), Padova (Italy), Haifa (Israel), Kunming (China), Istanbul (Turkey), Zwolle (Netherlands), Barcelona (Spain), Kortrijk (Belgium), Barretos (Brazil), Zhengzhou (China), Moscow (Russian Federation), Trier (Germany), Bucuresti (Romania), Itajai (Brazil), Craiova (Romania), Berlin (Germany), Dalian (China), Oslo (Norway), Baudour (Belgium), Novara (Italy), Tel Aviv (Israel), Connecticut, Kfar Saba (Israel), Cremona (Italy), Petah Tikva (Israel), Nanning (China), Kitakyushu-shi (Japan), Elche (Spain), Woolloongabba (Australia), Kazan (Russian Federation), Darlinghurst (Australia), Xiamen (China), Sumy (Ukraine), Kobe-shi (Japan), Bucharest (Romania), Frankston (Australia)

NCT02411448
PHASE 3

A Multicenter, Randomized, Double-Blind Study of Erlotinib in Combination With Ramucirumab or Placebo in Previously Untreated Patients With EGFR Mutation-Positive Metastatic Non-Small Cell Lung Cancer

TARGETS
EGFR, VEGFR2

LOCATIONS: Pordenone (Italy), Jinju (Korea, Republic of), Pok Fu Lam (Hong Kong), Texas, Chemnitz (Germany), Kaohsiung City (Taiwan), Poitiers (France), Hyogo (Japan), Dongjak-gu (Korea, Republic of), Taichung (Taiwan), Osaka (Japan), Cluj-Napoca (Romania), Pennsylvania, Kobe (Japan), Fukuoka (Japan), Grenoble (France), Gerlingen (Germany), Pamplona (Spain), Málaga (Spain), Okayama (Japan), Suwon (Korea, Republic of), Bucuresti (Romania), Torino (Italy), Milano (Italy), Montpellier (France), Taipei (Taiwan), Taoyuan (Taiwan), Hong Kong (Hong Kong), Akashi (Japan), Lille (France), New York, Berlin (Germany), Merseyside (United Kingdom), Wolverhampton (United Kingdom), Nagoya (Japan), Massachusetts, Nottingham (United Kingdom), Miyagi (Japan), Barcelona (Spain), Ravenna (Italy), Himeji (Japan), Hirakata (Japan), Paris (France), Ube (Japan), Hawaii, North Carolina, Edmonton (Canada), California, Ulm (Germany), Matsuyama (Japan), Kyoto (Japan), Hwasun (Korea, Republic of), Yau Ma Tei (Hong Kong), Bron (France), Kurume (Japan), London (United Kingdom), Nagasaki (Japan), Wakayama (Japan), Kansas, London (Canada), Lai Chi Kok (Hong Kong), Tokyo (Japan), Seowon-gu (Korea, Republic of), Kashiwa (Japan), Saitama (Japan), Preston (United Kingdom), Kishiwada (Japan), Florida, Großhansdorf (Germany), Sevilla (Spain), Otopeni (Romania), Valencia (Spain), Habikino (Japan), Bunkyo-Ku (Japan), Yokohama (Japan), Chelsea (United Kingdom), Palma de Mallorca (Spain), Tainan (Taiwan), Bologna (Italy), Osakasayama (Japan), Kanazawa (Japan), Niigata (Japan), Seoul (Korea, Republic of), Asahikawa (Japan), Athens (Greece), Seongnam-Si (Korea, Republic of), Shizuoka (Japan), Patras (Greece), Padova (Italy), Oregon, Sakai (Japan), Köln (Germany), Chiba (Japan), Ulsan-si (Korea, Republic of), Madrid (Spain), Heidelberg (Germany)

NCT02438722
PHASE 2 / PHASE 3

A Randomized Phase II/III Trial of Afatinib Plus Cetuximab Versus Afatinib Alone in Treatment-Naïve Patients With Advanced, EGFR Mutation Positive Non-small Cell Lung Cancer (NSCLC)

TARGETS
EGFR, ERBB2, ERBB4

LOCATIONS: Vermont, Kentucky, New York, Mississippi, Idaho, Iowa, New Jersey, Massachusetts, Florida, Indiana, Wisconsin, Oregon, North Dakota, Montana, Ohio, Tennessee, Maine, South Dakota, New Hampshire, Oklahoma, West Virginia, Hawaii, Pennsylvania, Colorado, South Carolina, Wyoming, California, Michigan, Kansas, Nebraska, Illinois, Missouri, Minnesota, North Carolina, Georgia, Connecticut, Texas, Washington, New Mexico, Arkansas

NCT01822496
PHASE 2

A Randomized Phase II Study of Individualized Combined Modality Therapy for Stage III Non-Small Cell Lung Cancer (NSCLC)

TARGETS
MET, EGFR, ALK, TOP2, ROS1, AXL, TRKC, TRKA

LOCATIONS: Connecticut, Idaho, Iowa, New Jersey, Massachusetts, Indiana, Oregon, Missouri, New Hampshire, Maine, Illinois, Colorado, Maryland, Oklahoma, Arizona, Wisconsin, South Carolina, Nebraska, Florida, Minnesota, Texas, Pennsylvania, California, West Virginia, Michigan, Ohio, Delaware, Georgia, North Carolina, New York

NCT01857271
PHASE 2

Evaluation of Erlotinib as a Neoadjuvant Therapy in Stage III NSCLC Patients With EGFR Mutations (EVENT Trial)

TARGETS
EGFR

LOCATIONS: New York

NCT02108964
PHASE 1 / PHASE 2

A Phase I/II, Multicenter, Open-label Study of EGFRmut-TKI EGF816, Administered Orally in Adult Patients With EGFRmut Solid Malignancies

TARGETS
EGFR

LOCATIONS: Madrid (Spain), Taipei (Taiwan), Nagoya (Japan), Fukuoka (Japan), Massachusetts, Singapore (Singapore), Seoul (Korea, Republic of), New York, Berlin (Germany), Koeln (Germany), Milano (Italy), Amsterdam (Netherlands), Barcelona (Spain), Toronto (Canada)

NCT02716116
PHASE 1 / PHASE 2

A Phase 1/2 Study of the Safety, Pharmacokinetics, and Anti-Tumor Activity of the Oral EGFR/HER2 Inhibitor AP32788 in Non-Small Cell Lung Cancer

TARGETS
EGFR, ERBB2

LOCATIONS: New York, California, Tennessee, Massachusetts, Colorado, Virginia

NCT02574078
PHASE 1 / PHASE 2

A Master Protocol of Phase 1/2 Studies of Nivolumab in Advanced NSCLC Using Nivolumab as Maintenance After Induction Chemotherapy or as First-line Treatment Alone or in Combination With Standard of Care Therapies (CheckMate 370: CHECKpoint Pathway and nivoluMab Clinical Trial Evaluation 370)

TARGETS
EGFR, PD-1

LOCATIONS: Mississippi, Nebraska, North Carolina, Ohio, Colorado, California, Tennessee, Oklahoma, Pennsylvania, New Mexico, Indiana, Alabama, Virginia, Kansas, Illinois, North Dakota, Minnesota, Oregon, New York, Michigan, Missouri, Louisiana, South Carolina, Maine, Texas, Nevada, Kentucky, Georgia, New Jersey, Washington, West Virginia, Montana, Arizona, Connecticut, South Dakota, Maryland, Florida

NCT02349633
PHASE 1 / PHASE 2

Phase 1/2 Open-Label Study Of PF-06747775 (Epidermal Growth Factor Receptor T790m Inhibitor) In Patients With Advanced Epidermal Growth Factor Receptor Mutant (Del 19 Or L858R ± T790M) Non-Small Cell Lung Cancer

TARGETS
EGFR

LOCATIONS: Seoul (Korea, Republic of), Washington, California, Connecticut, Pennsylvania

SAMPLE

GENE
ERBB3

ALTERATION
P1212S
RATIONALE

ERBB3 amplification or activating mutations may be associated with response to therapies targeting ERBB3. Additionally, the oncogenic effects of ERBB3 mutations have been shown to depend on ERBB2/HER2, and tumors with ERBB3 oncogenic mutations may therefore be sensitive to therapies that target ERBB2/HER2. However, because this mutation has not been functionally characterized, it is not known whether these approaches are beneficial here. Examples of

clinical trials that may be appropriate for this patient are listed below. These trials were identified through a search of the trial website clinicaltrials.gov using keyword terms such as "ERBB3", "MM-121", "U3-1287", "AV-203", "afatinib", "pertuzumab", "lapatinib", "trastuzumab", "ado-trastuzumab emtansine", "lung", "NSCLC", "solid tumor", and/or "advanced cancer".

NCT01920061

A Phase 1b Open-label Three-arm Multi-center Study To Assess The Safety And Tolerability Of Pf-05212384 (pi3k/Mtor Inhibitor) In Combination With Other Anti-tumor Agents

PHASE 1
TARGETS

EGFR, mTORC1, ERBB2, PI3K-gamma, mTORC2, PI3K-alpha, ERBB4

LOCATIONS: London (United Kingdom), Massachusetts, South Carolina, California, Milan (Italy), Vancouver (Canada), Madrid (Spain), Pennsylvania, Toronto (Canada), Barcelona (Spain), Michigan, Roma (Italy)

NCT01306045

Pilot Trial of Molecular Profiling and Targeted Therapy for Advanced Non-Small Cell Lung Cancer, Small Cell Lung Cancer, and Thymic Malignancies

PHASE 2
TARGETS

AKTs, EGFR, RET, ERBB2, MEK, FLT3, VEGFRs, CSF1R, PDGFRs, KIT

LOCATIONS: Maryland

NCT02506517

Molecular Basket Trial In Multiple Malignancies With Common Target Pathway Aberrancies

PHASE 2
TARGETS

EGFR, ERBB2, ERBB4

LOCATIONS: Toronto (Canada)

NCT02451553

Phase I/IB Multi-center Study of Irreversible EGFR/HER2 Tyrosine Kinase Inhibitor Afatinib (BIBW 2992) in Combination With Capecitabine for Advanced Solid Tumors and Pancretico-Biliary Cancers

PHASE 1
TARGETS

EGFR, ERBB2, ERBB4

LOCATIONS: Indiana, Washington

NCT02912949

A Phase I/II Study of MCLA-128, a Full Length IgG1 Bispecific Antibody Targeting HER2 and HER3, in Patients With Solid Tumors

PHASE 1 / PHASE 2
TARGETS

ERBB2, ERBB3

LOCATIONS: Milan (Italy), Madrid (Spain), Paris (France), Amsterdam (Netherlands), Barcelona (Spain)

GENE
MET

ALTERATION
T263M

RATIONALE
Activation of MET may lead to increased MET expression and activation and may therefore confer sensitivity to MET inhibitors. However, it is not known whether these therapeutic approaches would be relevant in the context of alterations that have not been fully characterized, as seen here. Examples of clinical trials that may

be appropriate for this patient are listed below. These trials were identified through a search of the trial website clinicaltrials.gov using keyword terms such as "c-MET", "crizotinib", "cabozantinib", "INC280", "MGCD265", "NSCLC", and/or "solid tumor".

NCT01639508

PHASE 2

A Phase II Study of Cabozantinib in Patients With RET Fusion-Positive Advanced Non-Small Cell Lung Cancer and Those With Other Genotypes: ROS1 or NTRK Fusions or Increased MET or AXL Activity

TARGETS
MET, ROS1, RET, VEGFRs

LOCATIONS: New York, New Jersey

NCT02414139

PHASE 2

A Phase II, Multicenter Study of Oral cMET Inhibitor INC280 in Adult Patients With EGFR Wild-type (wt), Advanced Non-small Cell Lung Cancer (NSCLC)

TARGETS
MET

LOCATIONS: Arkansas, California, Connecticut, Florida, Georgia, Illinois, Iowa, Massachusetts, Michigan, Minnesota, Nebraska, New Hampshire, New York, Oregon, Pennsylvania, South Carolina, Tennessee, Texas, Utah, Virginia, multiple ex-US locations

NCT02954991

PHASE 2

A Parallel Phase 2 Study of Glesatinib, Sitravatinib or Mocetinostat in Combination With Nivolumab in Advanced or Metastatic Non-Small Cell Lung Cancer

TARGETS
MET, HDAC, PDGFRA, RET, PD-1, KIT, DDR2, VEGFRs, FLT3, AXL, TRKB, TRKA

LOCATIONS: Virginia, Tennessee, Wisconsin, Pennsylvania, Alabama, Ohio, Michigan, California

NCT02693535

PHASE 2

Targeted Agent and Profiling Utilization Registry (TAPUR) Study

TARGETS
ABL, CDK4, PARP, EGFR, DDR2, VEGFRs, PDGFRs, ROS1, CSF1R, ERBB2, PD-1, ERBB3, MEK, RAF1, KIT, SMO, AXL, TRKC, mTOR, TRKA, MET, ALK, BRAF, RET, SRC, FLT3, CDK6

LOCATIONS: North Dakota, Pennsylvania, Washington, Illinois, Georgia, Arizona, Utah, North Carolina, Oklahoma, South Dakota, Michigan, Oregon, Nebraska

NCT02584634

PHASE 2

A Phase 1b/2, Open Label, Dose Finding Study To Evaluate Safety, Efficacy, Pharmacokinetics And Pharmacodynamics Of Avelumab (msb0010718c) In Combination With Either Crizotinib Or Pf 06463922 In Patients With Advanced Or Metastatic Non Small Cell Lung Cancer Javelin Lung 101

TARGETS
MET, ALK, PD-L1, ROS1, AXL, TRKC, TRKA

LOCATIONS: Badalona (Spain), Camperdown (Australia), Barcelona (Spain), Georgia, Tennessee, Goyang-Si (Korea, Republic of), Goyang-si (Korea, Republic of), Melbourne (Australia), Massachusetts, Parkville (Australia), Fukuoka (Japan), Seoul (Korea, Republic of), Chermshire (Australia)

NCT00585195
PHASE 1

Phase 1 Safety, Pharmacokinetic And Pharmacodynamic Study Of Pf-02341066, A C-met/Hgfr Selective Tyrosine Kinase Inhibitor, Administered Orally To Patients With Advanced Cancer

TARGETS

MET, ALK, ROS1, AXL, TRKC, TRKA

LOCATIONS: New York, Michigan, California, Colorado, Pennsylvania, Akashi (Japan), Massachusetts, Melbourne (Australia), North Carolina, Ohio, Seoul (Korea, Republic of), Vermont, Sapporo (Japan), Osakasayama (Japan)

NCT00697632
PHASE 1

Open-Label Dose-Escalation Trial to Evaluate the Safety, Pharmacokinetics, and Pharmacodynamics of Daily Oral MGCD265 Administered Without Interruption to Subjects With Advanced Malignancies

TARGETS

MET, AXL

LOCATIONS: Texas, North Carolina, Pennsylvania, New York, Gyeonggi-do (Korea, Republic of), Edmonton (Canada), Montreal (Canada), Utah, Illinois, Massachusetts, Missouri, Vancouver (Canada), Seoul (Korea, Republic of), Washington, California

NCT02219711
PHASE 1

A Phase 1/1b Study of MGCD516 in Patients With Advanced Solid Tumor Malignancies

TARGETS

MET, PDGFRA, RET, KIT, DDR2, VEGFRs, FLT3, AXL, TRKB, TRKA

LOCATIONS: Texas, Nebraska, Tennessee, Wisconsin, Florida, Pennsylvania, Seoul (Korea, Republic of), Alabama, Virginia, South Carolina, Illinois, Utah, California, Missouri, New York, Washington, Massachusetts, Michigan, New Mexico

NCT02099058
PHASE 1

A Multicenter, Phase 1/1b, Open-Label, Dose-Escalation Study of ABBV-399, an Antibody Drug Conjugate, in Subjects With Advanced Solid Tumors

TARGETS

MET, EGFR, VEGFA, PD-1

LOCATIONS: Villejuif, Cedex (France), Colorado, Illinois, Massachusetts, Tennessee, Michigan, Missouri, Marseille (France), California, North Carolina, Virginia, Texas

NCT02132598
PHASE 2

A Single-Arm Phase II Clinical Trial of Cabozantinib (XL184) in Patients With Previously Treated Non-Small Cell Lung Cancer (NSCLC) With Brain Metastases With and Without C Met Amplification

TARGETS

MET, ROS1, RET, VEGFRs

LOCATIONS: Pennsylvania

APPENDIX
Variants of Unknown Significance

NOTE One or more variants of unknown significance (VUS) were detected in this patient's tumor. These variants may not have been adequately characterized in the scientific literature at the time this report was issued, and/or the genomic context of these alterations makes their significance unclear. We choose to include them here in the event that they become clinically meaningful in the future.

ATR
A2165T

EPHB1
I332V

IDH1
I102T

IRF2
E232G

IRF4
amplification

NOTCH3
V508L

ROS1
S1463N

SDHA
amplification

TEK
loss

SAMPLE

INTENDED USE

FoundationOne CDx™ (F1CDx) is a next generation sequencing based in vitro diagnostic device for detection of substitutions, insertion and deletion alterations (indels), and copy number alterations (CNAs) in 324 genes and select gene rearrangements, as well as genomic signatures including microsatellite instability (MSI) and tumor mutational burden (TMB) using DNA isolated from formalin-fixed paraffin embedded (FFPE) tumor tissue specimens. The test is intended as a companion diagnostic to identify patients who may benefit from treatment with the targeted therapies listed in Table 1 in accordance with the approved therapeutic product labeling. Additionally, F1CDx is intended to provide tumor mutation profiling to be used by qualified health care professionals in accordance with professional guidelines in oncology for patients with solid malignant neoplasms. The F1CDx assay is a single-site assay performed at Foundation Medicine, Inc.

INDICATION	GENOMIC FINDINGS	THERAPY
Non-small cell lung cancer (NSCLC)	<i>EGFR</i> exon 19 deletions and <i>EGFR</i> exon 21 L858R alterations	Gilotrif® (Afatinib), Iressa® (Gefitinib), or Tarceva® (Erlotinib)
	<i>EGFR</i> exon 20 T790M alterations	Tagrisso® (Osimertinib)
	<i>ALK</i> rearrangements	Alecensa® (Alectinib), Xalkori® (Crizotinib), or Zykadia® (Ceritinib)
	<i>BRAF</i> V600E	Tafinlar® (Dabrafenib) in combination with Mekinist® (Trametinib)
Melanoma	<i>BRAF</i> V600E	Tafinlar® (Dabrafenib) or Zelboraf® (Vemurafenib)
	<i>BRAF</i> V600E or V600K	Mekinist® (Trametinib) or Cotellic® (Cobimetinib), in combination with Zelboraf® (Vemurafenib)
Breast cancer	<i>ERBB2</i> (HER2) amplification	Herceptin® (Trastuzumab), Kadcyla® (Ado-trastuzumab emtansine), or Perjeta® (Pertuzumab)
Colorectal cancer	<i>KRAS</i> wild-type (absence of mutations in codons 12 and 13)	Erbix® (Cetuximab)
	<i>KRAS</i> wild-type (absence of mutations in exons 2, 3, and 4) and <i>NRAS</i> wild type (absence of mutations in exons 2, 3, and 4)	Vectibix® (Panitumumab)
Ovarian cancer	<i>BRCA1/2</i> alterations	Rubraca® (Rucaparib)

TABLE 1

The median exon coverage for this sample is 869x

TEST PRINCIPLE

FoundationOne CDx will be performed exclusively as a laboratory service using DNA extracted from formalin-fixed, paraffin-embedded (FFPE) tumor samples. The proposed assay will employ a single DNA extraction method from routine FFPE biopsy or surgical resection specimens, 50-1000 ng of which will undergo whole-genome shotgun library construction and hybridization-based capture of all coding exons from 309 cancer-related genes, one promoter region, one non-coding (ncRNA), and select intronic regions from 34 commonly rearranged genes, 21 of which also include the coding exons. The assay therefore includes detection of alterations in a total of 324 genes. Using the Illumina® HiSeq 4000 platform, hybrid capture-selected libraries will be sequenced to high uniform depth (targeting >500X median coverage with >99% of exons at coverage >100X). Sequence data will be processed using a customized analysis pipeline designed to accurately detect all classes of genomic alterations, including base substitutions, indels, focal copy number amplifications, homozygous gene deletions, and selected genomic rearrangements (e.g., gene fusions). Additionally, genomic signatures including microsatellite instability (MS) and tumor mutational burden (TMB) will be reported.

PERFORMANCE CHARACTERISTICS

Please refer to product label:
foundationmedicine.com/f1cdx

LIMITATIONS

1. For *in vitro* diagnostic use.
2. For prescription use only. This test must be ordered by a qualified medical professional in accordance with clinical laboratory regulations.
3. Genomic findings other than those listed in Table 1 of the intended use are not prescriptive or conclusive for labeled use of any specific therapeutic product.
4. A negative result does not rule out the presence of a mutation below the limits of detection of the assay.
5. Samples with <25% tumor may have decreased sensitivity for the detection of CNAs including *ERBB2*.
6. Clinical performance of Tagrisso® (osimertinib) in patients with an *EGFR* exon 20 T790M mutation detected with an allele fraction <5% is ongoing and has not been established.
7. Concordance with other validated methods for CNA (with the exception of *ERBB2*) and gene rearrangement (with the exception of *ALK*) detection has not been demonstrated and will be provided in the post-market setting. Confirmatory testing using a clinically validated assay should be performed for all CNAs and rearrangements not associated with CDx claims noted in Table 1 of the Intended Use, but used for clinical decision making.
8. The MSI-H/MSS designation by FMI F1CDx test is based on genome wide analysis of 95 microsatellite loci and not based on the 5 or 7 MSI loci described in current clinical practice guidelines. Refer to the Summary of Safety of Effectiveness Data (SSED) for additional details on methodology. The threshold for MSI-H/MSS was determined by analytical concordance to comparator assays (IHC and PCR) using uterine, cecum and colorectal cancer FFPE tissue. The clinical validity of the qualitative MSI designation has not been established.
9. TMB by F1CDx is defined based on counting the total number of all synonymous and non-synonymous variants present at 5% allele frequency or greater (after filtering) and reported as mutations per megabase (mut/Mb) unit. The clinical validity of TMB defined by this panel has not been established.
10. Decisions on patient care and treatment must be based on the independent medical judgment of the treating physician, taking into consideration all applicable information concerning the patient's condition, such as patient and family history, physical examinations, information from other diagnostic tests, and patient preferences, in accordance with the standard of care in a given community.
11. The test is intended to be performed on specific serial number-controlled instruments by Foundation Medicine, Inc.

APPENDIX

Genes assayed in FoundationOne CDx

FoundationOne CDx™ is designed to include all genes known to be somatically altered in human solid tumors that are validated targets for therapy, either approved or in clinical trials, and/or that are unambiguous drivers of oncogenesis based on current knowledge. The current assay interrogates 324 genes as well as introns of 28 genes involved in rearrangements. The assay will be updated periodically to reflect new knowledge about cancer biology.

DNA GENE LIST: ENTIRE CODING SEQUENCE FOR THE DETECTION OF BASE SUBSTITUTIONS, INSERTION/DELETIONS, AND COPY NUMBER ALTERATIONS

ABL1	ACVR1B	AKT1	AKT2	AKT3	ALK	ALOX12B	AMER1 (FAM123B)	APC
AR	ARAF	ARFRP1	ARID1A	ASXL1	ATM	ATR	ATRAX	AURKA
AURKB	AXIN1	AXL	BAP1	BARD1	BCL2	BCL2L1	BCL2L2	BCL6
BCOR	BCORL1	BRAF	BRCA1	BRCA2	BRD4	BRIP1	BTG1	BTG2
BTK	C11orf30 (EMSY)	C17orf39 (GID4)	CALR	CARD11	CASP8	CBFB	CBL	CCND1
CCND2	CCND3	CCNE1	CD22	CD274 (PD-L1)	CD70	CD79A	CD79B	CDC73
CDH1	CDK12	CDK4	CDK6	CDK8	CDKN1A	CDKN1B	CDKN2A	CDKN2B
CDKN2C	CEBPA	CHEK1	CHEK2	CIC	CREBBP	CRKL	CSF1R	CSF3R
CTCF	CTNNA1	CTNNB1	CUL3	CUL4A	CXCR4	CYP17A1	DAXX	DDR1
DDR2	DIS3	DNMT3A	DOT1L	EED	EGFR	EP300	EPHA3	EPHB1
EPHB4	ERBB2	ERBB3	ERBB4	ERCC4	ERG	ERRF1	ESR1	EZH2
FAM46C	FANCA	FANCC	FANGC	FANCL	FAS	FBXW7	FGF10	FGF12
FGF14	FGF19	FGF23	FGF3	FGF4	FGF6	FGFR1	FGFR2	FGFR3
FGFR4	FH	FLCN	FLT1	FLT3	FOXL2	FUBP1	GABRA6	GATA3
GATA4	GATA6	GNA11	GNA13	GNAQ	GNAS	GRM3	GSK3B	H3F3A
HDAC1	HGF	HNFI1A	HRAS	HSD3B1	ID3	IDH1	IDH2	IGF1R
IKBKE	IKZF1	INPP4B	IRF2	IRF4	IRS2	JAK1	JAK2	JAK3
JUN	KDMSA	KDMS5C	KDM6A	KDR	KEAP1	KEL	KIT	KLHL6
KMT2A (MLL)	KMT2D (MLL2)	KRAS	LTK	LYN	MAF	MAP2K1 (MEK1)	MAP2K2 (MEK2)	MAP2K4
MAP3K1	MAP3K13	MAPK1	MCL1	MDM2	MDM4	MED12	MEF2B	MEN1
MERTK	MET	MITF	MKNK1	MLH1	MPL	MRE11A	MSH2	MSH3
MSH6	MST1R	MTAP	MTOR	MUTYH	MYC	MYCL (MYCL1)	MYCN	MYD88
NBN	NF1	NF2	NFE2L2	NFKBIA	NKX2-1	NOTCH1	NOTCH2	NOTCH3
NPM1	NRAS	NT5C2	NTRK1	NTRK2	NTRK3	P2RY8	PALB2	PARK2
PARP1	PARP2	PARP3	PAX5	PBRM1	PDCD1 (PD-1)	PDCD1LG2 (PD-L2)	PDGFRA	PDGFRB
PDK1	PIK3C2B	PIK3C2G	PIK3CA	PIK3CB	PIK3R1	PIM1	PMS2	POLD1
POLE	PPARG	PPP2R1A	PPP2R2A	PRDM1	PRKAR1A	PRKCI	PTCH1	PTEN
PTPN11	PTPRO	QKI	RAC1	RAD21	RAD51	RAD51B	RAD51C	RAD51D
RADS2	RADS4L	RAF1	RARA	RB1	RBM10	REL	RET	RICTOR
RNF43	ROS1	RPTOR	SDHA	SDHB	SDHC	SDHD	SETD2	SF3B1
SGK1	SMAD2	SMAD4	SMARCA4	SMARCB1	SMO	SNCAIP	SOC3	SOX2
SOX9	SPEN	SPOP	SRC	STAG2	STAT3	STK11	SUFU	SYK
TBX3	TEK	TET2	TGFBR2	TIPARP	TNFAIP3	TNFRSF14	TP53	TSC1
TSC2	TYRO3	U2AF1	VEGFA	VHL	WHSC1	WHSC1L1	WT1	XPO1
XRCC2	ZNF217	ZNF703						

DNA GENE LIST: FOR THE DETECTION OF SELECT REARRANGEMENTS

ALK	BCL2	BCR	BRAF	BRCA1	BRCA2	CD74	EGFR	ETV4
ETV5	ETV6	EWSR1	EZR	FGFR1	FGFR2	FGFR3	KIT	KMT2A (MLL)
MSH2	MYB	MYC	NOTCH2	NTRK1	NTRK2	NUTM1	PDGFRA	RAF1
RARA	RET	ROS1	RSPO2	SDC4	SLC34A2	TERC*	TERT**	TMPPRSS2

*TERC IS A NCRNA

**THE PROMOTER REGION OF TERT INTERROGATED

ADDITIONAL ASSAYS: FOR THE DETECTION OF SELECT CANCER BIOMARKERS

Microsatellite Status (MS)
Tumor Mutation Burden (TMB)

APPENDIX

Information Provided as a Professional Service

QUALIFIED ALTERATION CALLS (EQUIVOCAL AND SUBCLONAL)

An alteration denoted as “amplification –equivocal” implies that the FoundationOne CDx™ assay data provide some, but not unambiguous, evidence that the copy number of a gene exceeds the threshold for identifying copy number amplification. The threshold used in FoundationOne CDx™ for identifying a copy number amplification is four (4) for *ERBB2* and six (6) for all other genes. Conversely, an alteration denoted as “loss – equivocal” implies that the FoundationOne CDx™ assay data provide some, but not unambiguous, evidence for homozygous deletion of the gene in question. An alteration denoted as “subclonal” is one that the FoundationOne CDx™ analytical methodology has identified as being present in <10% of the assayed tumor DNA.

PROFESSIONAL SERVICES FINDINGS

Incorporates analyses of peer-reviewed studies and other publicly available information identified by Foundation Medicine; these analyses and information may include associations between a molecular alteration (or lack of alteration) and one or more drugs with potential clinical benefit (or potential lack of clinical benefit), including drug candidates that are being studied in clinical research. NOTE: A finding of biomarker alteration does not necessarily indicate pharmacologic effectiveness (or lack thereof) of any drug or treatment regimen; a finding of no biomarker alteration does not necessarily indicate lack of pharmacologic effectiveness (or effectiveness) of any drug or treatment regimen.

RANKING OF ALTERATIONS AND DRUGS

Biomarker Findings

Appear at the top of the report, but are not ranked higher than Genomic Findings.

Genomic Findings

Therapies with Clinical Benefit In Patient's Tumor Type → Therapies with Clinical Benefit in Other Tumor Type → Clinical Trial Options → No Known Options (if multiple findings exist within any of these categories, the results are listed alphabetically by gene name).

Therapies

Sensitizing therapies → Resistant therapies (if multiple therapies exist within any of these categories, they are listed in no particular order).

Clinical Trials

Pediatric trial qualification → Geographical Proximity → Later trial phase.

LEVEL OF EVIDENCE NOT PROVIDED

Drugs with potential clinical benefit (or potential lack of clinical benefit) are not evaluated for source or level of published evidence.

NO GUARANTEE OF CLINICAL BENEFIT

Foundation Medicine makes no promises or guarantees that a particular drug will be effective in the treatment of disease of any patient. This report also makes no promises or guarantees that a drug with potential lack of clinical benefit will in fact provide no clinical benefit.

NO GUARANTEE OF REIMBURSEMENT

Foundation Medicine makes no promises or guarantees that a healthcare provider, insurer or other third party payor, whether private or governmental, will reimburse a patient for the cost of FoundationOne CDx.

TREATMENT DECISIONS ARE RESPONSIBILITY OF PHYSICIAN

Drugs referenced may not be suitable for a particular patient. The selection of any, all or none of the drugs associated with potential clinical benefit (or potential lack of clinical benefit) resides with the physician. Indeed, the information in this Report must be considered in conjunction with all other relevant information regarding a particular patient, before the patient's treating physician recommends a course of treatment. Decisions on patient care and treatment must be based on the independent medical judgment of the treating physician, taking into consideration all applicable information concerning the patient's condition, such as patient and family history, physical examinations, information from other diagnostic tests, and patient preferences, in accordance with the standard of care in a given community. A treating physician's decisions should not be based on a single test, such as this Test, or the information contained in this Report.

Genomic Findings with Evidence of Clinical Significance

Genomic findings listed at Level 2 are associated with clinical significance. Clinical significance may be indicated by evidence of therapeutic sensitivity or resistance and/or diagnostic, prognostic or other clinically relevant implications. Included in this category will be findings associated with clinical validity as supported by professional guidelines and/or peer-reviewed publications.

Genomic Findings with Potential Clinical Significance

Genomic findings listed at Level 3 are cancer-related mutations and biomarkers with potential clinical significance. These include findings in genes known to be associated with cancer and are supported by evidence from publicly available databases, and/or peer-reviewed publications.

A Fluid Approach to Reporting Levels

As additional information becomes available, as recognized by the clinical community (professional guidelines and/or peer-reviewed publications), findings may move between Levels 2 and 3 in accordance with the above descriptions.

SELECT ABBREVIATIONS

ABBREVIATION	DEFINITION
CR	Complete response
DCR	Disease control rate
DNMT	DNA methyltransferase
HR	Hazard ratio
ITD	Internal tandem duplication
MMR	Mismatch repair
muts/Mb	Mutations per megabase
NOS	Not otherwise specified
ORR	Objective response rate
OS	Overall survival
PD	Progressive disease
PFS	Progression-free survival
PR	Partial response
SD	Stable disease
TKI	Tyrosine kinase inhibitor

APPENDIX

References Associated with Professional Services Content

- 1 Gatalica Z, Snyder C, Maney T, et al. (2014) Programmed cell death 1 (PD-1) and its ligand (PD-L1) in common cancers and their correlation with molecular cancer type. *Cancer Epidemiol Biomarkers Prev* 23(12):2965-70
- 2 Kroemer G, Galluzzi L, Zitvogel L, et al. (2015) Colorectal cancer: the first neoplasia found to be under immunosurveillance and the last one to respond to immunotherapy? *Oncoimmunology* 4(7):e1058597
- 3 Lal N, Beggs AD, Willcox BE, et al. (2015) An immunogenomic stratification of colorectal cancer: Implications for development of targeted immunotherapy. *Oncoimmunology* 4(3):e976052
- 4 Overman et al., 2016; ASCO Abstract 3501
- 5 Le DT, Uram JN, Wang H, et al. (2015) PD-1 Blockade in Tumors with Mismatch-Repair Deficiency. *N Engl J Med* ePub May 2015
- 6 ASCO-SITC 2016; Abstract P60
- 7 Warth A, Körner S, Penzel R, et al. (2016) Microsatellite instability in pulmonary adenocarcinomas: a comprehensive study of 480 cases. *Virchows Arch* 468(3):313-9
- 8 Ninomiya H, Nomura K, Satoh Y, et al. (2006) Genetic instability in lung cancer: concurrent analysis of chromosomal, mini- and microsatellite instability and loss of heterozygosity. *Br J Cancer* 94(10):1485-91
- 9 Woenckhaus M, Stoehr R, Dietmaier W, et al. (2003) Microsatellite instability at chromosome 8p in non-small cell lung cancer is associated with lymph node metastasis and squamous differentiation. *Int J Oncol* 23(5):1357-63
- 10 Chang JW, Chen YC, Chen CY, et al. (2000) Correlation of genetic instability with mismatch repair protein expression and p53 mutations in non-small cell lung cancer. *Clin Cancer Res* 6(5):1639-46
- 11 Fong KM, Zimmerman PV, Smith PJ (1995) Microsatellite instability and other molecular abnormalities in non-small cell lung cancer. *Cancer Res* 55(1):28-30
- 12 Hansen LT, Thykjaer T, Ørntoft TF, et al. (2003) The role of mismatch repair in small-cell lung cancer cells. *Eur J Cancer* 39(10):1456-67
- 13 Kocarnik JM, Shiovitz S, Phipps AI (2015) Molecular phenotypes of colorectal cancer and potential clinical applications. *Gastroenterol Rep (Oxf)* 3(4):269-76
- 14 You JF, Buhard O, Ligtenberg MJ, et al. (2010) Tumours with loss of MSH6 expression are MSI-H when screened with a pentaplex of five mononucleotide repeats. *Br J Cancer* 103(12):1840-5
- 15 Bairwa NK, Saha A, Gochhait S, et al. (2014) Microsatellite instability: an indirect assay to detect defects in the cellular mismatch repair machinery. *Methods Mol Biol* 1105:497-509
- 16 Boland CR, Thibodeau SN, Hamilton SR, et al. (1998) A National Cancer Institute Workshop on Microsatellite Instability for cancer detection and familial predisposition: development of international criteria for the determination of microsatellite instability in colorectal cancer. *Cancer Res* 58(22):5248-57
- 17 Pawlik TM, Raut CP, Rodriguez-Bigas MA (2004) Colorectal carcinogenesis: MSI-H versus MSI-L. *Dis Markers* 20(4-5):199-206
- 18 Boland CR, Goel A (2010) Microsatellite instability in colorectal cancer. *Gastroenterology* 138(6):2073-2087.e3
- 19 Snyder A, Makarov V, Merghoub T, et al. (2014) Genetic basis for clinical response to CTLA-4 blockade in melanoma. *N Engl J Med* 371(23):2189-99
- 20 Rosenberg JE, Hoffman-Censits J, Powles T, et al. (2016) Atezolizumab in patients with locally advanced and metastatic urothelial carcinoma who have progressed following treatment with platinum-based chemotherapy: a single-arm, multicentre, phase 2 trial. *Lancet* 387(10031):1909-20
- 21 Johnson DB, Frampton GM, Rioth MJ, et al. (2016) Targeted next generation sequencing identifies markers of response to PD-1 blockade. *Cancer Immunol Res* ePub Sep 2016
- 22 Balar AV, Galsky MD, Rosenberg JE, et al. (2017) Atezolizumab as first-line treatment in cisplatin-ineligible patients with locally advanced and metastatic urothelial carcinoma: a single-arm, multicentre, phase 2 trial. *Lancet* 389(10064):67-76
- 23 Dong ZY, Zhong WZ, Zhang XC, et al. (2016) Potential Predictive Value of TP53 and KRAS Mutation Status for Response to PD-1 Blockade Immunotherapy in Lung Adenocarcinoma. *Clin Cancer Res*
- 24 Rizvi NA, Hellmann MD, Snyder A, et al. (2015) Cancer immunology. Mutational landscape determines sensitivity to PD-1 blockade in non-small cell lung cancer. *Science* 348(6230):124-8
- 25 Mehnert JM, Panda A, Zhong H, et al. (2016) Immune activation and response to pembrolizumab in POLE-mutant endometrial cancer. *J Clin Invest* 126(6):2334-40
- 26 Santin AD, Bellone S, Buza N, et al. (2016) Regression of chemotherapy-resistant Polymerase epsilon (POLE) ultra-mutated and MSH6 hyper-mutated endometrial tumors with nivolumab. *Clin Cancer Res* ePub Aug 2016
- 27 Johanns TM, Miller CA, Dorward IG, et al. (2016) Immunogenomics of Hypermutated Glioblastoma: a Patient with Germline POLE Deficiency Treated with Checkpoint Blockade Immunotherapy. *Cancer Discov* ePub Sep 2016
- 28 Bouffet E, Larouche V, Campbell BB, et al. (2016) Immune Checkpoint Inhibition for Hypermutant Glioblastoma Multiforme Resulting From Germline Biallelic Mismatch Repair Deficiency. *J Clin Oncol* ePub Mar 2016
- 29 Van Allen EM, Miao D, Schilling B, et al. (2015) Genomic correlates of response to CTLA-4 blockade in metastatic melanoma. *Science* 350(6257):207-11
- 30 Spigel et al., 2016; ASCO Abstract 9017
- 31 Schwartz et al., 2016; ASCO Abstract 8533
- 32 Xiao D, Pan H, Li F, et al. (2016) Analysis of ultra-deep targeted sequencing reveals mutation burden is associated with gender and clinical outcome in lung adenocarcinoma. *Oncotarget* 7(16):22857-64
- 33 Shim HS, Kenudson M, Zheng Z, et al. (2015) Unique Genetic and Survival Characteristics of Invasive Mucinous Adenocarcinoma of the Lung. *J Thorac Oncol* 10(8):1156-62
- 34 Imielinski M, Berger AH, Hammerman PS, et al. (2012) Mapping the hallmarks of lung adenocarcinoma with massively parallel sequencing. *Cell* 150(6):1107-20
- 35 Govindan R, Ding L, Griffith M, et al. (2012) Genomic landscape of non-small cell lung cancer in smokers and never-smokers. *Cell* 150(6):1121-34
- 36 Ding L, Getz G, Wheeler DA, et al. (2008) Somatic mutations affect key pathways in lung adenocarcinoma. *Nature* 455(7216):1069-75
- 37 Kim Y, Hammerman PS, Kim J, et al. (2014) Integrative and comparative genomic analysis of lung squamous cell carcinomas in East Asian patients. *J Clin Oncol* 32(2):121-8
- 38 Pfeifer GP, You YH, Besaratinia A (2005) Mutations induced by ultraviolet light. *Mutat Res* 571(1-2):19-31
- 39 Hill VK, Gartner JJ, Samuels Y, et al. (2013) The genetics of melanoma: recent advances. *Annu Rev Genomics Hum Genet* 14:257-79
- 40 Pfeifer GP, Denissenko MF, Olivier M, et al. (2002) Tobacco smoke carcinogens, DNA damage and p53 mutations in smoking-associated cancers. *Oncogene* 21(48):7435-51
- 41 Cancer Genome Atlas Research Network, Kandoth C, Schultz N, et al. (2013) Integrated genomic characterization of endometrial carcinoma. *Nature* 497(7447):67-73
- 42 Briggs S, Tomlinson I (2013) Germline and somatic polymerase ε and δ mutations define a new class of hypermutated colorectal and endometrial cancers. *J Pathol* 230(2):148-53
- 43 Heitzer E, Tomlinson I (2014) Replicative DNA polymerase mutations in cancer. *Curr Opin Genet Dev* 24:107-13
- 44 Cancer Genome Atlas Network (2012) Comprehensive molecular characterization of human colon and rectal cancer. *Nature* 487(7407):330-7
- 45 Roberts SA, Gordenin DA (2014) Hypermutation in human cancer genomes: footprints and mechanisms. *Nat Rev Cancer* 14(12):786-800
- 46 Rosell R, Carcereny E, Gervais R, et al. (2012) Erlotinib versus standard chemotherapy as first-line treatment for European patients with advanced EGFR mutation-positive non-small-cell lung cancer (EURTAC): a multicentre, open-label, randomised phase 3 trial. *Lancet Oncol* 13(3):239-46
- 47 Douillard JY, Ostoros G, Cobo M, et al. (2014) First-line gefitinib in Caucasian EGFR mutation-positive NSCLC patients: a phase-IV, open-label, single-arm study. *Br J Cancer* 110(1):55-62
- 48 Sequist LV, Yang JC, Yamamoto N, et al. (2013) Phase III study of afatinib or cisplatin plus pemetrexed in patients with metastatic lung adenocarcinoma with EGFR mutations. *J Clin Oncol* 31(27):3327-34
- 49 Jänne PA, Yang JC, Kim DW, et al. (2015) AZD9291 in EGFR inhibitor-resistant non-small-cell lung cancer. *N Engl J Med* 372(18):1689-99
- 50 Pirker R, Pereira JR, von Pawel J, et al. (2012) EGFR expression as a predictor of survival for first-line chemotherapy plus cetuximab in patients with advanced non-small-cell lung cancer: analysis of data from the phase 3 FLEX study. *Lancet Oncol* 13(1):33-42
- 51 Jänne PA, Ou SH, Kim DW, et al. (2014) Dacomitinib as first-line treatment in patients with clinically or molecularly selected advanced non-small-cell lung cancer: a multicentre, open-label, phase 2 trial. *Lancet Oncol* 15(13):1433-41

APPENDIX

References Associated with Professional Services Content

- 52 Thatcher N, Hirsch FR, Luft AV, et al. (2015) Necitumumab plus gemcitabine and cisplatin versus gemcitabine and cisplatin alone as first-line therapy in patients with stage IV squamous non-small-cell lung cancer (SQUIRE): an open-label, randomised, controlled phase 3 trial. *Lancet Oncol* 16(7):763-74
- 53 Paz-Ares L, Mezger J, Ciuleanu TE, et al. (2015) Necitumumab plus pemetrexed and cisplatin as first-line therapy in patients with stage IV non-squamous non-small-cell lung cancer (INSPIRE): an open-label, randomised, controlled phase 3 study. *Lancet Oncol* 16(3):328-37
- 54 Garon et al., 2012; ASCO Abstract 7543
- 55 Sequist LV, Gettinger S, Senzer NN, et al. (2010) Activity of IPI-504, a novel heat-shock protein 90 inhibitor, in patients with molecularly defined non-small-cell lung cancer. *J Clin Oncol* 28(33):4953-60
- 56 Socinski MA, Goldman J, El-Hariri I, et al. (2013) A multicenter phase II study of ganetespib monotherapy in patients with genotypically defined advanced non-small cell lung cancer. *Clin Cancer Res* 19(11):3068-77
- 57 Isambert N, Delord JP, Soria JC, et al. (2015) Debio0932, a second-generation oral heat shock protein (HSP) inhibitor, in patients with advanced cancer—results of a first-in-man dose-escalation study with a fixed-dose extension phase. *Ann Oncol* ePub Feb 2015
- 58 Johnson ML, Yu HA, Hart EM, et al. (2015) Phase I/II Study of HSP90 Inhibitor AUY922 and Erlotinib for EGFR-Mutant Lung Cancer With Acquired Resistance to Epidermal Growth Factor Receptor Tyrosine Kinase Inhibitors. *J Clin Oncol* ePub Apr 2015
- 59 Piotrowska et al., 2015; ASCO Abstract 8015
- 60 Strong JE, Coffey MC, Tang D, et al. (1998) The molecular basis of viral oncolysis: usurpation of the Ras signaling pathway by reovirus. *EMBO J* 17(12):3351-62
- 61 Coffey MC, Strong JE, Forsyth PA, et al. (1998) Reovirus therapy of tumors with activated Ras pathway. *Science* 282(5392):1332-4
- 62 Gong J, Mita MM (2014) Activated ras signaling pathways and reovirus oncolysis: an update on the mechanism of preferential reovirus replication in cancer cells. *Front Oncol* 4:167
- 63 Forsyth P, Roldán G, George D, et al. (2008) A phase I trial of intratumoral administration of reovirus in patients with histologically confirmed recurrent malignant gliomas. *Mol Ther* 16(3):627-32
- 64 Vidal L, Pandha HS, Yap TA, et al. (2008) A phase I study of intravenous oncolytic reovirus type 3 Dearing in patients with advanced cancer. *Clin Cancer Res* 14(21):7127-37
- 65 Gollamudi R, Ghalib MH, Desai KK, et al. (2010) Intravenous administration of Reolysin, a live replication competent RNA virus is safe in patients with advanced solid tumors. *Invest New Drugs* 28(5):641-9
- 66 Harrington KJ, Karapanagiotou EM, Roulstone V, et al. (2010) Two-stage phase I dose-escalation study of intratumoral reovirus type 3 dearing and palliative radiotherapy in patients with advanced cancers. *Clin Cancer Res* 16(11):3067-77
- 67 Comins C, Spicer J, Protheroe A, et al. (2010) REO-10: a phase I study of intravenous reovirus and docetaxel in patients with advanced cancer. *Clin Cancer Res* 16(22):5564-72
- 68 Lolkema MP, Arkenau HT, Harrington K, et al. (2011) A phase I study of the combination of intravenous reovirus type 3 Dearing and gemcitabine in patients with advanced cancer. *Clin Cancer Res* 17(3):581-8
- 69 Galanis E, Markovic SN, Suman VJ, et al. (2012) Phase II trial of intravenous administration of Reolysin® (Reovirus Serotype-3-dearing Strain) in patients with metastatic melanoma. *Mol Ther* 20(10):1998-2003
- 70 Karapanagiotou EM, Roulstone V, Twigger K, et al. (2012) Phase I/II trial of carboplatin and paclitaxel chemotherapy in combination with intravenous oncolytic reovirus in patients with advanced malignancies. *Clin Cancer Res* 18(7):2080-9
- 71 Morris DG, Feng X, DiFrancesco LM, et al. (2013) REO-001: A phase I trial of percutaneous intralesional administration of reovirus type 3 dearing (Reolysin®) in patients with advanced solid tumors. *Invest New Drugs* 31(3):696-706
- 72 Liang Z, Zhang J, Zeng X, et al. (2010) Relationship between EGFR expression, copy number and mutation in lung adenocarcinomas. *BMC Cancer* 10:376
- 73 Cancer Genome Atlas Research Network (2014) Comprehensive molecular profiling of lung adenocarcinoma. *Nature* 511(7511):543-50
- 74 Park S, Choi YL, Sung CO, et al. (2012) High MET copy number and MET overexpression: poor outcome in non-small cell lung cancer patients. *Histol Histopathol* 27(2):197-207
- 75 Grob TJ, Hoenig T, Clauditz TS, et al. (2013) Frequent intratumoral heterogeneity of EGFR gene copy gain in non-small cell lung cancer. *Lung Cancer* 79(3):221-7
- 76 Vallee A, Sagan C, Le Loupp AG, et al. (2013) Detection of EGFR gene mutations in non-small cell lung cancer: lessons from a single-institution routine analysis of 1,403 tumor samples. *Int J Oncol* 43(4):1045-51
- 77 Watzka SB, Rauscher-Pötsch I, Nierlich P, et al. (2010) Concordance between epidermal growth factor receptor status in primary non-small-cell lung cancer and metastases: a post-mortem study. *Eur J Cardiothorac Surg* 38(1):34-7
- 78 Marks JL, Broderick S, Zhou Q, et al. (2008) Prognostic and therapeutic implications of EGFR and KRAS mutations in resected lung adenocarcinoma. *J Thorac Oncol* 3(2):111-6
- 79 Izar B, Sequist L, Lee M, et al. (2013) The impact of EGFR mutation status on outcomes in patients with resected stage I non-small cell lung cancers. *Ann Thorac Surg* 96(3):962-8
- 80 Ciardiello F, Tortora G (2008) EGFR antagonists in cancer treatment. *N Engl J Med* 358(11):1160-74
- 81 Bhargava R, Gerald WL, Li AR, et al. (2005) EGFR gene amplification in breast cancer: correlation with epidermal growth factor receptor mRNA and protein expression and HER-2 status and absence of EGFR-activating mutations. *Mod Pathol* 18(8):1027-33
- 82 Yang YL, Xu KL, Zhou Y, et al. (2012) Correlation of epidermal growth factor receptor overexpression with increased epidermal growth factor receptor gene copy number in esophageal squamous cell carcinomas. *Chin Med J* 125(3):450-4
- 83 Lynch TJ, Bell DW, Sordella R, et al. (2004) Activating mutations in the epidermal growth factor receptor underlying responsiveness of non-small-cell lung cancer to gefitinib. *N Engl J Med* 350(21):2129-39
- 84 Paez JG, Jänne PA, Lee JC, et al. (2004) EGFR mutations in lung cancer: correlation with clinical response to gefitinib therapy. *Science* 304(5676):1497-500
- 85 Pao W, Miller V, Zakowski M, et al. (2004) EGF receptor gene mutations are common in lung cancers from "never smokers" and are associated with sensitivity of tumors to gefitinib and erlotinib. *Proc Natl Acad Sci USA* 101(36):13306-11
- 86 Jaiswal BS, Kijavini NM, Stawiski EW, et al. (2013) Oncogenic ERBB3 mutations in human cancers. *Cancer Cell* 23(5):603-17
- 87 Baselga J, Swain SM (2009) Novel anticancer targets: revisiting ERBB2 and discovering ERBB3. *Nat Rev Cancer* 9(7):463-75
- 88 Jura N, Shan Y, Cao X, et al. (2009) Structural analysis of the catalytically inactive kinase domain of the human EGF receptor 3. *Proc Natl Acad Sci USA* 106(51):21608-13
- 89 Choudhury NJ, Campanile A, Antic T, et al. (2016) Afatinib Activity in Platinum-Refractory Metastatic Urothelial Carcinoma in Patients With ERBB Alterations. *J Clin Oncol* ePub Apr 2016
- 90 Bidard FC, Ng CK, Cottu P, et al. (2015) Response to dual HER2 blockade in a patient with HER3-mutant metastatic breast cancer. *Ann Oncol* ePub May 2015
- 91 Bellezza G, Del Sordo R, Colella R, et al. (2013) Co-expression of receptors of the HER family correlates with clinical outcome in non-small cell lung cancer (NSCLC). *Virchows Arch* 463(5):663-71
- 92 Müller-Tidow C, Diederichs S, Bulk E, et al. (2005) Identification of metastasis-associated receptor tyrosine kinases in non-small cell lung cancer. *Cancer Res* 65(5):1778-82
- 93 Sheng Q, Liu J (2011) The therapeutic potential of targeting the EGFR family in epithelial ovarian cancer. *Br J Cancer* 104(8):1241-5
- 94 Jung KH, Park BH, Hong SS (2012) Progress in cancer therapy targeting c-Met signaling pathway. *Arch Pharm Res* 35(4):595-604
- 95 Shaw AT, Kim DW, Nakagawa K, et al. (2013) Crizotinib versus chemotherapy in advanced ALK-positive lung cancer. *N Engl J Med* 368(25):2385-94
- 96 Hart CD, De Boer RH (2013) Profile of cabozantinib and its potential in the treatment of advanced medullary thyroid cancer. *Onco Targets Ther* 6:1-7
- 97 Vassal et al., 2015; ASCO Abstract 2595
- 98 Li et al., 2015; ASCO Abstract 8090
- 99 Ou SH, Kwak EL, Siwak-Tapp C, et al. (2011) Activity of crizotinib (PF02341066), a dual mesenchymal-epithelial transition (MET) and anaplastic lymphoma kinase (ALK) inhibitor, in a non-small cell lung cancer patient with de novo MET amplification. *J Thorac Oncol* 6(5):942-6

APPENDIX

References Associated with Professional Services Content

- 100 Schwab R, Petak I, Kollar M, et al. (2014) Major partial response to crizotinib, a dual MET/ALK inhibitor, in a squamous cell lung (SCC) carcinoma patient with de novo c-MET amplification in the absence of ALK rearrangement. *Lung Cancer* 83(1):109-11
- 101 Le X, Freed JA, VanderLaan PA, et al. (2015) Detection of Crizotinib-Sensitive Lung Adenocarcinomas With MET, ALK, and ROS1 Genomic Alterations via Comprehensive Genomic Profiling. *Clin Lung Cancer* ePub Mar 2015
- 102 Ali SM, Sanford EM, Klempner SJ, et al. (2015) Prospective comprehensive genomic profiling of advanced gastric carcinoma cases reveals frequent clinically relevant genomic alterations and new routes for targeted therapies. *Oncologist* 20(5):499-507
- 103 Lerner JK, Kwak EL, Ackerman A, et al. (2011) MET amplification identifies a small and aggressive subgroup of esophagogastric adenocarcinoma with evidence of responsiveness to crizotinib. *J Clin Oncol* 29(36):4803-10
- 104 Chi AS, Batchelor TT, Kwak EL, et al. (2012) Rapid radiographic and clinical improvement after treatment of a MET-amplified recurrent glioblastoma with a mesenchymal-epithelial transition inhibitor. *J Clin Oncol* 30(3):e30-3
- 105 Palma NA, Ali SM, O'Connor J, et al. (2014) Durable Response to Crizotinib in a MET-Amplified, KRAS-Mutated Carcinoma of Unknown Primary. *Case Rep Oncol* 7(2):503-8
- 106 Frampton GM, Ali SM, Rosenzweig M, et al. (2015) Activation of MET via diverse exon 14 splicing alterations occurs in multiple tumor types and confers clinical sensitivity to MET inhibitors. *Cancer Discov* ePub May 2015
- 107 Paik PK, Drilon A, Yu H, et al. (2015) Response to MET inhibitors in patients with stage IV lung adenocarcinomas harboring MET mutations causing exon 14 skipping. *Cancer Discov* ePub May 2015
- 108 Jenkins RW, Oxnard GR, Elkin S, et al. (2015) Response to Crizotinib in a Patient With Lung Adenocarcinoma Harboring a MET Splice Site Mutation. *Clin Lung Cancer* ePub Feb 2015
- 109 Waqar SN, Morgensztern D, Sehn J (2015) MET Mutation Associated with Responsiveness to Crizotinib. *J Thorac Oncol* 10(5):e29-31
- 110 Mendenhall MA, Goldman JW (2015) MET-Mutated NSCLC with Major Response to Crizotinib. *J Thorac Oncol* 10(5):e33-4
- 111 Stein MN, Hirshfield KM, Zhong H, et al. (2015) Response to Crizotinib in a Patient with MET-mutant Papillary Renal Cell Cancer After Progression on Tivantinib. *Eur Urol* 67(2):353-4
- 112 Awad MM, Oxnard GR, Jackman DM, et al. (2016) MET Exon 14 Mutations in Non-Small-Cell Lung Cancer Are Associated With Advanced Age and Stage-Dependent MET Genomic Amplification and c-Met Overexpression. *J Clin Oncol* ePub Jan 2016
- 113 Liu X, Jia Y, Stoopler MB, et al. (2015) Next-Generation Sequencing of Pulmonary Sarcomatoid Carcinoma Reveals High Frequency of Actionable MET Gene Mutations. *J Clin Oncol* ePub Jul 2015
- 114 Yakes FM, Chen J, Tan J, et al. (2011) Cabozantinib (XL184), a novel MET and VEGFR2 inhibitor, simultaneously suppresses metastasis, angiogenesis, and tumor growth. *Mol Cancer Ther* 10(12):2298-308
- 115 Weber H, Müller D, Müller M, et al. (2014) Cell Lines Expressing Recombinant Transmembrane Domain-Activated Receptor Kinases as Tools for Drug Discovery. *J Biomol Screen* ePub Sep 2014
- 116 Navis AC, Bourgonje A, Wesseling P, et al. (2013) Effects of dual targeting of tumor cells and stroma in human glioblastoma xenografts with a tyrosine kinase inhibitor against c-MET and VEGFR2. *PLoS ONE* 8(3):e58262
- 117 Yeh I, Botton T, Talevich E, et al. (2015) Activating MET kinase rearrangements in melanoma and Spitz tumours. *Nat Commun* 6:7174
- 118 Lee YH, Apolo AB, Agarwal PK, et al. (2014) Characterization of HGF/Met Signaling in Cell Lines Derived From Urothelial Carcinoma of the Bladder. *Cancers (Basel)* 6(4):2313-29
- 119 Torres KE, Zhu QS, Bill K, et al. (2011) Activated MET is a molecular prognosticator and potential therapeutic target for malignant peripheral nerve sheath tumors. *Clin Cancer Res* 17(12):3943-55
- 120 Sameni M, Tovar EA, Essenburg C, et al. (2015) Cabozantinib (XL184) inhibits growth and invasion of preclinical TNBC models. *Clin Cancer Res* ePub Oct 2015
- 121 Kwak et al., 2015; ASCO GI Abstract 01
- 122 Kwak EL, Ahnion LG, Srivastava G, et al. (2015) Molecular Heterogeneity and Receptor Coamplification Drive Resistance to Targeted Therapy in MET-Amplified Esophagogastric Cancer. *Cancer Discov* 5(12):1271-81
- 123 Gan et al., 2015; ASCO Abstract 487
- 124 Falchook et al., 2015; ASCO Abstract 2591
- 125 Kollmannsberger et al., 2015; ASCO Abstract 2589
- 126 Diamond JR, Salgia R, Varella-Garcia M, et al. (2013) Initial clinical sensitivity and acquired resistance to MET inhibition in MET-mutated papillary renal cell carcinoma. *J Clin Oncol* 31(16):e254-8
- 127 Choueiri TK, Vaishampayan U, Rosenberg JE, et al. (2013) Phase II and biomarker study of the dual MET/VEGFR2 inhibitor foretinib in patients with papillary renal cell carcinoma. *J Clin Oncol* 31(2):181-6
- 128 Spigel DR, Ervin TJ, Ramlau RA, et al. (2013) Randomized phase II trial of Onartuzumab in combination with erlotinib in patients with advanced non-small-cell lung cancer. *J Clin Oncol* 31(32):4105-14
- 129 Catenacci DV, Henderson L, Xiao SY, et al. (2011) Durable complete response of metastatic gastric cancer with anti-Met therapy followed by resistance at recurrence. *Cancer Discov* 1(7):573-9
- 130 Oliner et al., 2012; ASCO Abstract 4005
- 131 Cancer Genome Atlas Research Network (2012) Comprehensive genomic characterization of squamous cell lung cancers. *Nature* 489(7417):519-25
- 132 Cappuzzo F, Marchetti A, Skokan M, et al. (2009) Increased MET gene copy number negatively affects survival of surgically resected non-small-cell lung cancer patients. *J Clin Oncol* 27(10):1667-74
- 133 Zucali PA, Ruiz MG, Giovannetti E, et al. (2008) Role of cMET expression in non-small-cell lung cancer patients treated with EGFR tyrosine kinase inhibitors. *Ann Oncol* 19(9):1605-12
- 134 An SJ, Chen ZH, Su J, et al. (2012) Identification of enriched driver gene alterations in subgroups of non-small cell lung cancer patients based on histology and smoking status. *PLoS ONE* 7(6):e40109
- 135 Yang JJ, Chen HJ, Yan HH, et al. (2013) Clinical modes of EGFR tyrosine kinase inhibitor failure and subsequent management in advanced non-small cell lung cancer. *Lung Cancer* 79(1):33-9
- 136 Dziadziuszko R, Wynes MW, Singh S, et al. (2012) Correlation between MET gene copy number by silver in situ hybridization and protein expression by immunohistochemistry in non-small cell lung cancer. *J Thorac Oncol* 7(2):340-7
- 137 Chen YT, Chang JW, Liu HP, et al. (2011) Clinical implications of high MET gene dosage in non-small cell lung cancer patients without previous tyrosine kinase inhibitor treatment. *J Thorac Oncol* 6(12):2027-35
- 138 Sun W, Ai T, Gao Y, et al. (2013) Expression and prognostic relevance of MET and phospho-BAD in non-small cell lung cancer. *Onco Targets Ther* 6:1315-23
- 139 Appleman LJ (2011) MET signaling pathway: a rational target for cancer therapy. *J Clin Oncol* 29(36):4837-8
- 140 Konecny GE, Winterhoff B, Kolarova T, et al. (2011) Expression of p16 and retinoblastoma determines response to CDK4/6 inhibition in ovarian cancer. *Clin Cancer Res* 17(6):1591-602
- 141 Katsumi Y, Iehara T, Miyachi M, et al. (2011) Sensitivity of malignant rhabdoid tumor cell lines to PD 0332991 is inversely correlated with p16 expression. *Biochem Biophys Res Commun* 413(1):62-8
- 142 Cen L, Carlson BL, Schroeder MA, et al. (2012) p16-Cdk4-Rb axis controls sensitivity to a cyclin-dependent kinase inhibitor PD0332991 in glioblastoma xenograft cells. *Neuro-oncology* 14(7):870-81
- 143 Logan JE, Mostofizadeh N, Desai AJ, et al. (2013) PD-0332991, a potent and selective inhibitor of cyclin-dependent kinase 4/6, demonstrates inhibition of proliferation in renal cell carcinoma at nanomolar concentrations and molecular markers predict for sensitivity. *Anticancer Res* 33(8):2997-3004
- 144 Elvin JA, Gay LM, Ort R, et al. (2017) Clinical Benefit in Response to Palbociclib Treatment in Refractory Uterine Leiomyosarcomas with a Common CDKN2A Alteration. *Oncologist* ePub Mar 2017
- 145 Gao J, Adams RP, Swain SM (2015) Does CDKN2A loss predict palbociclib benefit? *Curr Oncol* 22(6):e498-501
- 146 Gopalan et al., 2014; ASCO Abstract 8077
- 147 Peguero et al., 2016; ASCO Abstract 2528
- 148 Konecny et al., 2016; ASCO Abstract 5557
- 149 DeMichele A, Clark A, Tan KS, et al. (2014) CDK 4/6 Inhibitor Palbociclib (PD0332991) in Rb+ Advanced Breast Cancer: Phase II Activity, Safety and Predictive Biomarker Assessment. *Clin Cancer Res* ePub Dec 2014

The content provided as a professional service by Foundation Medicine, Inc., has not been reviewed or approved by the FDA.

Electronically signed by Shakti Ramkissoon, M.D., Ph.D. | Jeffrey Ross, M.D., Medical Director
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

APPENDIX

References Associated with Professional Services Content

- 150 Finn RS, Crown JP, Lang I, et al. (2014) The cyclin-dependent kinase 4/6 inhibitor palbociclib in combination with letrozole versus letrozole alone as first-line treatment of oestrogen receptor-positive, HER2-negative, advanced breast cancer (PALOMA-1/TRIO-18): a randomised phase 2 study. *Lancet Oncol* ePub Dec 2014
- 151 Infante JR, Cassier PA, Gerecitano JF, et al. (2016) A Phase I Study of the Cyclin-Dependent Kinase 4/6 Inhibitor Ribociclib (LEE011) in Patients with Advanced Solid Tumors and Lymphomas. *Clin Cancer Res* 22(23):5696-5705
- 152 Johnson DB, Dahlman KH, Knol J, et al. (2014) Enabling a genetically informed approach to cancer medicine: a retrospective evaluation of the impact of comprehensive tumor profiling using a targeted next-generation sequencing panel. *Oncologist* 19(6):616-22
- 153 Van Maerken T, Rihani A, Dreidax D, et al. (2011) Functional analysis of the p53 pathway in neuroblastoma cells using the small-molecule MDM2 antagonist nutlin-3. *Mol Cancer Ther* 10(6):983-93
- 154 Gamble LD, Kees UR, Tweddle DA, et al. (2012) MYCN sensitizes neuroblastoma to the MDM2-p53 antagonists Nutlin-3 and MI-63. *Oncogene* 31(6):752-63
- 155 Gazzeri S, Gouyer V, Vour'ch C, et al. (1998) Mechanisms of p16INK4A inactivation in non-small-cell lung cancers. *Oncogene* 16(4):497-504
- 156 Kratzke RA, Greatens TM, Rubins JB, et al. (1996) Rb and p16INK4a expression in resected non-small cell lung tumors. *Cancer Res* 56(15):3415-20
- 157 Lee JU, Sul HJ, Son JW (2012) Promoter Methylation of CDKN2A, RARβ, and RASSF1A in Non-Small Cell Lung Carcinoma: Quantitative Evaluation Using Pyrosequencing. *Tuberc Respir Dis (Seoul)* 73(1):11-21
- 158 Cortot AB, Younes M, Martel-Planche G, et al. (2014) Mutation of TP53 and alteration of p14(arf) expression in EGFR- and KRAS-mutated lung adenocarcinomas. *Clin Lung Cancer* 15(2):124-30
- 159 Mounawar M, Mukeria A, Le Calvez F, et al. (2007) Patterns of EGFR, HER2, TP53, and KRAS mutations of p14arf expression in non-small cell lung cancers in relation to smoking history. *Cancer Res* 67(12):5667-72
- 160 Kawabuchi B, Moriyama S, Hironaka M, et al. (1999) p16 inactivation in small-sized lung adenocarcinoma: its association with poor prognosis. *Int J Cancer* 84(1):49-53
- 161 Xing XB, Cai WB, Luo L, et al. (2013) The Prognostic Value of p16 Hypermethylation in Cancer: A Meta-Analysis. *PLoS ONE* 8(6):e66587
- 162 Lou-Qian Z, Rong Y, Ming L, et al. (2013) The prognostic value of epigenetic silencing of p16 gene in NSCLC patients: a systematic review and meta-analysis. *PLoS ONE* 8(1):e54970
- 163 Quelle DE, Zindy F, Ashmun RA, et al. (1995) Alternative reading frames of the INK4a tumor suppressor gene encode two unrelated proteins capable of inducing cell cycle arrest. *Cell* 83(6):993-1000
- 164 Sharpless NE (2005) INK4a/ARF: a multifunctional tumor suppressor locus. *Mutat Res* 576(1-2):22-38
- 165 Roussel MF (1999) The INK4 family of cell cycle inhibitors in cancer. *Oncogene* 18(38):5311-7
- 166 Sherr CJ, Bertwistle D, DEN Besten W, et al. (2005) p53-Dependent and -independent functions of the Arf tumor suppressor. *Cold Spring Harb Symp Quant Biol* 70:129-37
- 167 Ozenne P, Eymin B, Brambilla E, et al. (2010) The ARF tumor suppressor: structure, functions and status in cancer. *Int J Cancer* 127(10):2239-47
- 168 Parry D, Peters G (1996) Temperature-sensitive mutants of p16CDKN2 associated with familial melanoma. *Mol Cell Biol* 16(7):3844-52
- 169 Yang R, Gombart AF, Serrano M, et al. (1995) Mutational effects on the p16INK4a tumor suppressor protein. *Cancer Res* 55(12):2503-6
- 170 Poi MJ, Yen T, Li J, et al. (2001) Somatic INK4a-ARF locus mutations: a significant mechanism of gene inactivation in squamous cell carcinomas of the head and neck. *Mol Carcinog* 30(1):26-36
- 171 Arap W, Knudsen ES, Wang JY, et al. (1997) Point mutations can inactivate in vitro and in vivo activities of p16(INK4a)/CDKN2A in human glioma. *Oncogene* 14(5):603-9
- 172 Jafri M, Wake NC, Ascher DB, et al. (2015) Germline Mutations in the CDKN2B Tumor Suppressor Gene Predispose to Renal Cell Carcinoma. *Cancer Discov* 5(7):723-9
- 173 Zhang Y, Xiong Y (1999) Mutations in human ARF exon 2 disrupt its nucleolar localization and impair its ability to block nuclear export of MDM2 and p53. *Mol Cell* 3(5):579-91
- 174 Zhang Y, Xiong Y, Yarbrough WG (1998) ARF promotes MDM2 degradation and stabilizes p53: ARF-INK4a locus deletion impairs both the Rb and p53 tumor suppression pathways. *Cell* 92(6):725-34
- 175 Munshi PN, Lubin M, Bertino JR (2014) 6-thioguanine: a drug with unrealized potential for cancer therapy. *Oncologist* 19(7):760-5
- 176 de Oliveira SF, Ganzinelli M, Chilà R, et al. (2016) Characterization of MTAP Gene Expression in Breast Cancer Patients and Cell Lines. *PLoS ONE* 11(1):e0145647
- 177 Lubin M, Lubin A (2009) Selective killing of tumors deficient in methylthioadenosine phosphorylase: a novel strategy. *PLoS ONE* 4(5):e5735
- 178 Tang B, Testa JR, Kruger WD (2012) Increasing the therapeutic index of 5-fluorouracil and 6-thioguanine by targeting loss of MTAP in tumor cells. *Cancer Biol Ther* 13(11):1082-90
- 179 Collins CC, Volik SV, Lapuk AV, et al. (2012) Next generation sequencing of prostate cancer from a patient identifies a deficiency of methylthioadenosine phosphorylase, an exploitable tumor target. *Mol Cancer Ther* 11(3):775-83
- 180 Bertino JR, Waud WR, Parker WB, et al. (2011) Targeting tumors that lack methylthioadenosine phosphorylase (MTAP) activity: current strategies. *Cancer Biol Ther* 11(7):627-32
- 181 Coulthard SA, Redfern CP, Vikingsson S, et al. (2011) Increased sensitivity to thiopurines in methylthioadenosine phosphorylase-deleted cancers. *Mol Cancer Ther* 10(3):495-504
- 182 Miyazaki S, Nishioka J, Shiraishi T, et al. (2007) Methylthioadenosine phosphorylase deficiency in Japanese osteosarcoma patients. *Int J Oncol* 31(5):1069-76
- 183 Efferth T, Miyachi H, Drexler HG, et al. Methylthioadenosine phosphorylase as target for chemoselective treatment of T-cell acute lymphoblastic leukemic cells. *Blood Cells Mol Dis* 28(1):47-56
- 184 Kindler HL, Burris HA, Sandler AB, et al. (2009) A phase II multicenter study of L-alanosine, a potent inhibitor of adenine biosynthesis, in patients with MTAP-deficient cancer. *Invest New Drugs* 27(1):75-81
- 185 Marjon K, Cameron MJ, Quang P, et al. (2016) MTAP Deletions in Cancer Create Vulnerability to Targeting of the MAT2A/PRMT5/RIOK1 Axis. *Cell Rep* 15(3):574-587
- 186 Mavrakis KJ, McDonald ER, Schlabach MR, et al. (2016) Disordered methionine metabolism in MTAP/CDKN2A-deleted cancers leads to dependence on PRMT5. *Science* 351(6278):1208-13
- 187 Kryukov GV, Wilson FH, Ruth JR, et al. (2016) MTAP deletion confers enhanced dependency on the PRMT5 arginine methyltransferase in cancer cells. *Science* 351(6278):1214-8
- 188 Wei R, Zhao M, Zheng CH, et al. (2016) Concordance between somatic copy number loss and down-regulated expression: A pan-cancer study of cancer predisposition genes. *Sci Rep* 6:37358
- 189 Zhao M, Zhao Z (2016) Concordance of copy number loss and down-regulation of tumor suppressor genes: a pan-cancer study. *BMC Genomics* 17 Suppl 7:532
- 190 Kirovski G, Stevens AP, Czech B, et al. (2011) Down-regulation of methylthioadenosine phosphorylase (MTAP) induces progression of hepatocellular carcinoma via accumulation of 5'-deoxy-5'-methylthioadenosine (MTA). *Am J Pathol* 178(3):1145-52
- 191 Huang HY, Li SH, Yu SC, et al. (2009) Homozygous deletion of MTAP gene as a poor prognosticator in gastrointestinal stromal tumors. *Clin Cancer Res* 15(22):6963-72
- 192 Marcé S, Balagué O, Colomo L, et al. (2006) Lack of methylthioadenosine phosphorylase expression in mantle cell lymphoma is associated with shorter survival: implications for a potential targeted therapy. *Clin Cancer Res* 12(12):3754-61
- 193 Meyer S, Wild PJ, Vogt T, et al. (2010) Methylthioadenosine phosphorylase represents a predictive marker for response to adjuvant interferon therapy in patients with malignant melanoma. *Exp Dermatol* 19(8):e251-7
- 194 Wild PJ, Meyer S, Bataille F, et al. (2006) Tissue microarray analysis of methylthioadenosine phosphorylase protein expression in melanocytic skin tumors. *Arch Dermatol* 142(4):471-6
- 195 Kim J, Kim MA, Min SY, et al. (2011) Downregulation of methylthioadenosine phosphorylase by homozygous deletion in gastric carcinoma. *Genes Chromosomes Cancer* 50(6):421-33
- 196 Li CF, Fang FM, Kung HJ, et al. (2014) Downregulated MTAP expression in myxofibrosarcoma: A characterization of inactivating mechanisms, tumor suppressive function, and therapeutic relevance. *Oncotarget* 5(22):11428-41
- 197 He HL, Lee YE, Shiue YL, et al. (2015) Characterization and Prognostic Significance of Methylthioadenosine Phosphorylase Deficiency in Nasopharyngeal Carcinoma. *Medicine (Baltimore)* 94(49):e2271

The content provided as a professional service by Foundation Medicine, Inc., has not been reviewed or approved by the FDA.

Electronically signed by Shakti Ramkissoon, M.D., Ph.D. | Jeffrey Ross, M.D., Medical Director
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

APPENDIX

References Associated with Professional Services Content

- 198 Su CY, Chang YC, Chan YC, et al. (2014) MTAP is an independent prognosis marker and the concordant loss of MTAP and p16 expression predicts short survival in non-small cell lung cancer patients. *Eur J Surg Oncol* 40(9):1143-50
- 199 Mirebeau D, Acquaviva C, Suci S, et al. (2006) The prognostic significance of CDKN2A, CDKN2B and MTAP inactivation in B-lineage acute lymphoblastic leukemia of childhood. Results of the EORTC studies 58881 and 58951. *Haematologica* 91(7):881-5
- 200 Becker AP, Scapulatempo-Neto C, Menezes WP, et al. (2015) Expression of Methylthioadenosine Phosphorylase (MTAP) in Pilocytic Astrocytomas. *Pathobiology* 82(2):84-9
- 201 Snezhkina AV, Kravnov GS, Lipatova AV, et al. (2016) The Dysregulation of Polyamine Metabolism in Colorectal Cancer Is Associated with Overexpression of c-Myc and C/EBPβ rather than Enterotoxigenic *Bacteroides fragilis* Infection. *Oxid Med Cell Longev* 2016:2353560
- 202 Bistulfi G, Affronti HC, Foster BA, et al. (2016) The essential role of methylthioadenosine phosphorylase in prostate cancer. *Oncotarget* 7(12):14380-93
- 203 Antonopoulou K, Stefanaki I, Lill CM, et al. (2015) Updated field synopsis and systematic meta-analyses of genetic association studies in cutaneous melanoma: the MelGene database. *J Invest Dermatol* 135(4):1074-1079
- 204 Maccioni L, Rachakonda PS, Bermejo JL, et al. (2013) Variants at the 9p21 locus and melanoma risk. *BMC Cancer* 13:325
- 205 Hyland PL, Zhang H, Yang Q, et al. (2016) Pathway, in silico and tissue-specific expression quantitative analyses of oesophageal squamous cell carcinoma genome-wide association studies data. *Int J Epidemiol* 45(1):206-20
- 206 Lin X, Yan C, Gao Y, et al. (2017) Genetic variants at 9p21.3 are associated with risk of esophageal squamous cell carcinoma in a Chinese population. *Cancer Sci* 108(2):250-255
- 207 Zhi L, Liu D, Wu SG, et al. (2016) Association of common variants in MTAP with susceptibility and overall survival of osteosarcoma: a two-stage population-based study in Han Chinese. *J Cancer* 7(15):2179-2186
- 208 Gu F, Pfeiffer RM, Bhattacharjee S, et al. (2013) Common genetic variants in the 9p21 region and their associations with multiple tumours. *Br J Cancer* 108(6):1378-86
- 209 Limm K, Dettmer K, Reinders J, et al. (2016) Characterization of the Methylthioadenosine Phosphorylase Polymorphism rs7023954 - Incidence and Effects on Enzymatic Function in Malignant Melanoma. *PLoS ONE* 11(8):e0160348
- 210 Tang B, Kadariya Y, Chen Y, et al. (2014) Expression of MTAP inhibits tumor-related phenotypes in HT1080 cells via a mechanism unrelated to its enzymatic function. *G3 (Bethesda)* 5(1):35-44
- 211 Limm K, Ott C, Wallner S, et al. (2013) Deregulation of protein methylation in melanoma. *Eur J Cancer* 49(6):1305-13
- 212 Stevens AP, Spangler B, Wallner S, et al. (2009) Direct and tumor microenvironment mediated influences of 5'-deoxy-5'-(methylthio)adenosine on tumor progression of malignant melanoma. *J Cell Biochem* 106(2):210-9
- 213 Limm K, Wallner S, Milenkovic VM, et al. (2014) The metabolite 5'-methylthioadenosine signals through the adenosine receptor A2B in melanoma. *Eur J Cancer* 50(15):2714-24
- 214 Bivona TG, Hieronymus H, Parker J, et al. (2011) FAS and NF-κB signalling modulate dependence of lung cancers on mutant EGFR. *Nature* 471(7339):523-6
- 215 Giannikopoulos et al., 2014; ASCO Abstract 8083
- 216 Zhao Z, Zhong X, Wu T, et al. (2014) Identification of a NFKBIA polymorphism associated with lower NFKBIA protein levels and poor survival outcomes in patients with glioblastoma multiforme. *Int J Mol Med* 34(5):1233-40
- 217 Geng P, Ou J, Li J, et al. (2015) Genetic Association Between NFKBIA -881A>G Polymorphism and Cancer Susceptibility. *Medicine (Baltimore)* 94(31):e1024
- 218 Zhang M, Huang J, Tan X, et al. (2015) Common Polymorphisms in the NFKBIA Gene and Cancer Susceptibility: A Meta-Analysis. *Med Sci Monit* 21:186-96
- 219 Weniger MA, Küppers R (2016) NF-κB deregulation in Hodgkin lymphoma. *Semin Cancer Biol* ePub May 2016
- 220 Liu X, Yu H, Yang W, et al. (2010) Mutations of NFKBIA in biopsy specimens from Hodgkin lymphoma. *Cancer Genet Cytogenet* 197(2):152-7
- 221 Lake A, Shield LA, Cordano P, et al. (2009) Mutations of NFKBIA, encoding IκappaB alpha, are a recurrent finding in classical Hodgkin lymphoma but are not a unifying feature of non-EBV-associated cases. *Int J Cancer* 125(6):1334-42
- 222 Birnstiel ML, Jacob J, Sirlin JL (1965) Analysis of nucleolar RNA synthesis in dipteran salivary glands. *Arch Biol (Liege)* 76(2):565-89
- 223 Cabannes E, Khan G, Aillet F, et al. (1999) Mutations in the IκBa gene in Hodgkin's disease suggest a tumour suppressor role for IκappaBα. *Oncogene* 18(20):3063-70
- 224 Bredel M, Scholtens DM, Yadav AK, et al. (2011) NFKBIA deletion in glioblastomas. *N Engl J Med* 364(7):627-37
- 225 Patané M, Poratti P, Bottega E, et al. (2013) Frequency of NFKBIA deletions is low in glioblastomas and skewed in glioblastoma neurospheres. *Mol Cancer* 12:160
- 226 Gao J, Aksoy BA, Dogrusoz U, et al. (2013) Integrative analysis of complex cancer genomics and clinical profiles using the cBioPortal. *Sci Signal* 6(269):p11
- 227 Zack TI, Schumacher SE, Carter SL, et al. (2013) Pan-cancer patterns of somatic copy number alteration. *Nat Genet* 45(10):1134-1140
- 228 Beroukhi R, Mermel CH, Porter D, et al. (2010) The landscape of somatic copy-number alteration across human cancers. *Nature* 463(7283):899-905
- 229 Hsu DS, Acharya CR, Balakumaran BS, et al. (2009) Characterizing the developmental pathways TTF-1, NKX2-8, and PAX9 in lung cancer. *Proc Natl Acad Sci USA* 106(13):5312-7
- 230 Yang L, Lin M, Ruan WJ, et al. (2012) Nkx2-1: a novel tumor biomarker of lung cancer. *J Zhejiang Univ Sci B* 13(11):855-66
- 231 Kumar A, Coleman I, Morrissey C, et al. (2016) Substantial interindividual and limited intraindividual genomic diversity among tumors from men with metastatic prostate cancer. *Nat Med* 22(4):369-78
- 232 Landa I, Ibrahimipasc T, Boucai L, et al. (2016) Genomic and transcriptomic hallmarks of poorly differentiated and anaplastic thyroid cancers. *J Clin Invest* ePub Feb 2016
- 233 Jiao Y, Yonescu R, Offerhaus GJ, et al. (2014) Whole-exome sequencing of pancreatic neoplasms with acinar differentiation. *J Pathol* 232(4):428-35
- 234 Jones S, Stransky N, McCord CL, et al. (2014) Genomic analyses of gynaecologic carcinosarcomas reveal frequent mutations in chromatin remodelling genes. *Nat Commun* 5:5006
- 235 Nakamura N, Miyagi E, Murata S, et al. (2002) Expression of thyroid transcription factor-1 in normal and neoplastic lung tissues. *Mod Pathol* 15(10):1058-67
- 236 Moldvay J, Jackel M, Bogos K, et al. (2004) The role of TTF-1 in differentiating primary and metastatic lung adenocarcinomas. *Pathol Oncol Res* 10(2):85-8
- 237 Gilbert-Sirieix M, Makouji J, Kimura S, et al. (2011) Wnt/β-catenin signaling pathway is a direct enhancer of thyroid transcription factor-1 in human papillary thyroid carcinoma cells. *PLoS ONE* 6(7):e22280
- 238 Robens J, Goldstein L, Gown AM, et al. (2010) Thyroid transcription factor-1 expression in breast carcinomas. *Am J Surg Pathol* 34(12):1881-5
- 239 Ni YB, Tsang JY, Shao MM, et al. (2014) TTF-1 expression in breast carcinoma: an unusual but real phenomenon. *Histopathology* 64(4):504-11
- 240 Tsai LH, Chen PM, Cheng YW, et al. (2014) LKB1 loss by alteration of the NKX2-1/p53 pathway promotes tumor malignancy and predicts poor survival and relapse in lung adenocarcinomas. *Oncogene* 33(29):3851-60
- 241 Tan D, Li Q, Deeb G, et al. (2003) Thyroid transcription factor-1 expression prevalence and its clinical implications in non-small cell lung cancer: a high-throughput tissue microarray and immunohistochemistry study. *Hum Pathol* 34(6):597-604
- 242 Haque AK, Syed S, Lele SM, et al. (2002) Immunohistochemical study of thyroid transcription factor-1 and HER2/neu in non-small cell lung cancer: strong thyroid transcription factor-1 expression predicts better survival. *Appl Immunohistochem Mol Morphol* 10(2):103-9
- 243 Pelosi G, Frassetto F, Pasini F, et al. (2001) Immunoreactivity for thyroid transcription factor-1 in stage I non-small cell carcinomas of the lung. *Am J Surg Pathol* 25(3):363-72
- 244 Yamaguchi T, Hosono Y, Yanagisawa K, et al. (2013) NKX2-1/TTF-1: an enigmatic oncogene that functions as a double-edged sword for cancer cell survival and progression. *Cancer Cell* 23(6):718-23
- 245 Mu D (2013) The complexity of thyroid transcription factor 1 with both pro- and anti-oncogenic activities. *J Biol Chem* 288(35):24992-5000
- 246 Hamdan H, Liu H, Li C, et al. (1998) Structure of the human Nkx2.1 gene. *Biochim Biophys Acta* 1396(3):336-48

The content provided as a professional service by Foundation Medicine, Inc., has not been reviewed or approved by the FDA.

Electronically signed by Shakti Ramkissoon, M.D., Ph.D. | Jeffrey Ross, M.D., Medical Director
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

APPENDIX

References Associated with Professional Services Content

- 247 Kwei KA, Kim YH, Girard L, et al. (2008) Genomic profiling identifies TITF1 as a lineage-specific oncogene amplified in lung cancer. *Oncogene* 27(25):3635-40
- 248 Hirai H, Arai T, Okada M, et al. (2010) MK-1775, a small molecule Wee1 inhibitor, enhances anti-tumor efficacy of various DNA-damaging agents, including 5-fluorouracil. *Cancer Biol Ther* 9(7):514-22
- 249 Bridges KA, Hirai H, Buser CA, et al. (2011) MK-1775, a novel Wee1 kinase inhibitor, radiosensitizes p53-defective human tumor cells. *Clin Cancer Res* 17(17):5638-48
- 250 Rajeshkumar NV, De Oliveira E, Ottenhof N, et al. (2011) MK-1775, a potent Wee1 inhibitor, synergizes with gemcitabine to achieve tumor regressions, selectively in p53-deficient pancreatic cancer xenografts. *Clin Cancer Res* 17(9):2799-806
- 251 Osman AA, Monroe MM, Ortega Alves MV, et al. (2015) Wee-1 kinase inhibition overcomes cisplatin resistance associated with high-risk TP53 mutations in head and neck cancer through mitotic arrest followed by senescence. *Mol Cancer Ther* 14(2):608-19
- 252 Gourley et al., 2016; ASCO Abstract 5571
- 253 Lehmann S, Bykov VJ, Ali D, et al. (2012) Targeting p53 in vivo: a first-in-human study with p53-targeting compound APR-246 in refractory hematologic malignancies and prostate cancer. *J Clin Oncol* 30(29):3633-9
- 254 Mohell N, Alfredsson J, Fransson Å, et al. (2015) APR-246 overcomes resistance to cisplatin and doxorubicin in ovarian cancer cells. *Cell Death Dis* 6:e1794
- 255 Fransson Å, Glaessgen D, Alfredsson J, et al. (2016) Strong synergy with APR-246 and DNA-damaging drugs in primary cancer cells from patients with TP53 mutant High-Grade Serous ovarian cancer. *J Ovarian Res* 9(1):27
- 256 Xu L, Huang CC, Huang W, et al. (2002) Systemic tumor-targeted gene delivery by anti-transferrin receptor scFv-immunoliposomes. *Mol Cancer Ther* 1(5):337-46
- 257 Xu L, Tang WH, Huang CC, et al. (2001) Systemic p53 gene therapy of cancer with immunoliposomes targeted by anti-transferrin receptor scFv. *Mol Med* 7(10):723-34
- 258 Camp ER, Wang C, Little EC, et al. (2013) Transferrin receptor targeting nanomedicine delivering wild-type p53 gene sensitizes pancreatic cancer to gemcitabine therapy. *Cancer Gene Ther* 20(4):222-8
- 259 Kim SS, Rait A, Kim E, et al. (2015) A tumor-targeting p53 nanodelivery system limits chemoresistance to temozolomide prolonging survival in a mouse model of glioblastoma multiforme. *Nanomedicine* 11(2):301-11
- 260 Pirolo KF, Nemunaitis J, Leung PK, et al. (2016) Safety and Efficacy in Advanced Solid Tumors of a Targeted Nanocomplex Carrying the p53 Gene Used in Combination with Docetaxel: A Phase 1b Study. *Mol Ther* 24(9):1697-706
- 261 Hajdenberg et al., 2012; ASCO Abstract e15010
- 262 Leijen S, van Geel RM, Pavlick AC, et al. (2016) Phase I Study Evaluating WEE1 Inhibitor AZD1775 As Monotherapy and in Combination With Gemcitabine, Cisplatin, or Carboplatin in Patients With Advanced Solid Tumors. *J Clin Oncol ePub Sep 2016*
- 263 Oza et al., 2015; ASCO Abstract 5506
- 264 Leijen et al., 2015; ASCO Abstract 2507
- 265 Ma CX, Cai S, Li S, et al. (2012) Targeting Chk1 in p53-deficient triple-negative breast cancer is therapeutically beneficial in human-in-mouse tumor models. *J Clin Invest* 122(4):1541-52
- 266 Kumar et al., 2012; AACR Abstract 2874
- 267 Mogi A, Kuwano H (2011) TP53 mutations in nonsmall cell lung cancer. *J Biomed Biotechnol* 2011:583929
- 268 Tekpli X, Landvik NE, Skaug V, et al. (2013) Functional effect of polymorphisms in 15q25 locus on CHRNA5 mRNA, bulky DNA adducts and TP53 mutations. *Int J Cancer* 132(8):1811-20
- 269 Vignot S, Frampton GM, Soria JC, et al. (2013) Next-generation sequencing reveals high concordance of recurrent somatic alterations between primary tumor and metastases from patients with non-small-cell lung cancer. *J Clin Oncol* 31(17):2167-72
- 270 Maeng CH, Lee HY, Kim YW, et al. (2013) High-throughput molecular genotyping for small biopsy samples in advanced non-small cell lung cancer patients. *Anticancer Res* 33(11):5127-33
- 271 Itakura M, Terashima Y, Shingyoji M, et al. (2013) High CC chemokine receptor 7 expression improves postoperative prognosis of lung adenocarcinoma patients. *Br J Cancer* 109(5):1100-8
- 272 Seo JS, Ju YS, Lee WC, et al. (2012) The transcriptional landscape and mutational profile of lung adenocarcinoma. *Genome Res* 22(11):2109-19
- 273 Brown CJ, Lain S, Verma CS, et al. (2009) Awakening guardian angels: drugging the p53 pathway. *Nat Rev Cancer* 9(12):862-73
- 274 Joerger AC, Fersht AR (2008) Structural biology of the tumor suppressor p53. *Annu Rev Biochem* 77:557-82
- 275 Kato S, Han SY, Liu W, et al. (2003) Understanding the function-structure and function-mutation relationships of p53 tumor suppressor protein by high-resolution missense mutation analysis. *Proc Natl Acad Sci USA* 100(14):8424-9
- 276 Kamada R, Nomura T, Anderson CW, et al. (2011) Cancer-associated p53 tetramerization domain mutants: quantitative analysis reveals a low threshold for tumor suppressor inactivation. *J Biol Chem* 286(1):252-8
- 277 Bougeard G, Renaux-Petel M, Flaman JM, et al. (2015) Revisiting Li-Fraumeni Syndrome From TP53 Mutation Carriers. *J Clin Oncol* 33(21):2345-52
- 278 Sorrell AD, Espenschied CR, Culver JO, et al. (2013) Tumor protein p53 (TP53) testing and Li-Fraumeni syndrome: current status of clinical applications and future directions. *Mol Diagn Ther* 17(1):31-47
- 279 Nichols KE, Malkin D, Garber JE, et al. (2001) Germ-line p53 mutations predispose to a wide spectrum of early-onset cancers. *Cancer Epidemiol Biomarkers Prev* 10(2):83-7
- 280 Taubert H, Meyer A, Würl P (1998) Soft tissue sarcomas and p53 mutations. *Mol Med* 4(6):365-72
- 281 Kleihues P, Schäuble B, zur Hausen A, et al. (1997) Tumors associated with p53 germline mutations: a synopsis of 91 families. *Am J Pathol* 150(1):1-13
- 282 Gonzalez KD, Noltner KA, Buzin CH, et al. (2009) Beyond Li Fraumeni Syndrome: clinical characteristics of families with p53 germline mutations. *J Clin Oncol* 27(8):1250-6
- 283 Laloo F, Varley J, Ellis D, et al. (2003) Prediction of pathogenic mutations in patients with early-onset breast cancer by family history. *Lancet* 361(9363):1101-2
- 284 Cappuzzo F, Finocchiaro G, Grossi F, et al. (2015) Phase II study of afatinib, an irreversible ErbB family blocker, in EGFR FISH-positive non-small-cell lung cancer. *J Thorac Oncol* 10(4):665-72
- 285 Kwak EL, Shapiro GI, Cohen SM, et al. (2013) Phase 2 trial of afatinib, an ErbB family blocker, in solid tumors genetically screened for target activation. *Cancer* 119(16):3043-51
- 286 Li M, Zhang Z, Li X, et al. (2014) Whole-exome and targeted gene sequencing of gallbladder carcinoma identifies recurrent mutations in the ErbB pathway. *Nat Genet* 46(8):872-6
- 287 Wu YL, Zhou C, Hu CP, et al. (2014) Afatinib versus cisplatin plus gemcitabine for first-line treatment of Asian patients with advanced non-small-cell lung cancer harbouring EGFR mutations (LUX-Lung 6): an open-label, randomised phase 3 trial. *Lancet Oncol* 15(2):213-22
- 288 Yang JC, Wu YL, Schuler M, et al. (2015) Afatinib versus cisplatin-based chemotherapy for EGFR mutation-positive lung adenocarcinoma (LUX-Lung 3 and LUX-Lung 6): analysis of overall survival data from two randomised, phase 3 trials. *Lancet Oncol ePub Jan 2015*
- 289 Soria JC, Felip E, Cobo M, et al. (2015) Afatinib versus erlotinib as second-line treatment of patients with advanced squamous cell carcinoma of the lung (LUX-Lung 8): an open-label randomised controlled phase 3 trial. *Lancet Oncol ePub Jul 2015*
- 290 Paz-Ares L, Tan EH, O'Byrne K, et al. (2017) Afatinib versus gefitinib in patients with EGFR mutation-positive advanced non-small-cell lung cancer: overall survival data from the phase IIb LUX-Lung 7 trial. *Ann Oncol* 28(2):270-277
- 291 Miller VA, Hirsh V, Cadranel J, et al. (2012) Afatinib versus placebo for patients with advanced, metastatic non-small-cell lung cancer after failure of erlotinib, gefitinib, or both, and one or two lines of chemotherapy (LUX-Lung 1): a phase 2b/3 randomised trial. *Lancet Oncol* 13(5):528-38
- 292 Chen X, Zhu Q, Zhu L, et al. (2013) Clinical perspective of afatinib in non-small cell lung cancer. *Lung Cancer* 81(2):155-61
- 293 Katakami N, Atagi S, Goto K, et al. (2013) LUX-Lung 4: a phase II trial of afatinib in patients with advanced non-small-cell lung cancer who progressed during prior treatment with erlotinib, gefitinib, or both. *J Clin Oncol* 31(27):3335-41
- 294 Landi L, Tiseo M, Chiari R, et al. (2014) Activity of the EGFR-HER2 Dual Inhibitor Afatinib in EGFR-Mutant Lung Cancer Patients With Acquired Resistance to Reversible EGFR Tyrosine Kinase Inhibitors. *Clin Lung Cancer* 15(6):411-417.e4

APPENDIX

References Associated with Professional Services Content

- 295 De Grève J, Moran T, Graas MP, et al. (2015) Phase II study of afatinib, an irreversible ErbB family blocker, in demographically and genotypically defined lung adenocarcinoma. *Lung Cancer* 88(1):63-9
- 296 Yang JC, Sequist LV, Geater SL, et al. (2015) Clinical activity of afatinib in patients with advanced non-small-cell lung cancer harbouring uncommon EGFR mutations: a combined post-hoc analysis of LUX-Lung 2, LUX-Lung 3, and LUX-Lung 6. *Lancet Oncol* ePub Jun 2015
- 297 Wu SG, Liu YN, Tsai MF, et al. (2016) The mechanism of acquired resistance to irreversible EGFR tyrosine kinase inhibitor-afatinib in lung adenocarcinoma patients. *Oncotarget* ePub Feb 2016
- 298 Kim Y, Ko J, Cui Z, et al. (2012) The EGFR T790M mutation in acquired resistance to an irreversible second-generation EGFR inhibitor. *Mol Cancer Ther* 11(3):784-91
- 299 Tanaka K, Nosaki K, Otsubo K, et al. (2017) Acquisition of the T790M resistance mutation during afatinib treatment in EGFR tyrosine kinase inhibitor-naïve patients with non-small cell lung cancer harboring EGFR mutations. *Oncotarget* 8(40):68123-68130
- 300 Janjigian YY, Smit EF, Groen HJ, et al. (2014) Dual Inhibition of EGFR with Afatinib and Cetuximab in Kinase Inhibitor-Resistant EGFR-Mutant Lung Cancer with and without T790M Mutations. *Cancer Discov* ePub Jul 2014
- 301 Ribeiro Gomes J, Cruz MR (2015) Combination of afatinib with cetuximab in patients with EGFR-mutant non-small-cell lung cancer resistant to EGFR inhibitors. *Onco Targets Ther* 8:1137-42
- 302 Castellanos EH, Rivera G, Wakelee H, et al. (2015) Overcoming Resistance Without the Risk of Reaction: Use of Afatinib and Panitumumab in Two Cases of Epidermal Growth Factor Receptor-Mutated Non-Small Cell Lung Cancer With T790M Mutations. *Clin Lung Cancer* ePub Mar 2015
- 303 Schuler M, Yang JC, Park K, et al. (2016) Afatinib beyond progression in patients with non-small-cell lung cancer following chemotherapy, erlotinib/gefitinib and afatinib: phase III randomized LUX-Lung 5 trial. *Ann Oncol* 27(3):417-23
- 304 Horn L, Gettinger S, Camidge DR, et al. (2017) Continued use of afatinib with the addition of cetuximab after progression on afatinib in patients with EGFR mutation-positive non-small-cell lung cancer and acquired resistance to gefitinib or erlotinib. *Lung Cancer* 113:51-58
- 305 Camidge et al., 2014; ASCO Abstract 8001
- 306 Jorge SE, Schulman S, Freed JA, et al. (2015) Responses to the multitargeted MET/ALK/ROS1 inhibitor crizotinib and co-occurring mutations in lung adenocarcinomas with MET amplification or MET exon 14 skipping mutation. *Lung Cancer* 90(3):369-74
- 307 Zhang Y, Wang W, Wang Y, et al. (2016) Response to Crizotinib Observed in Lung Adenocarcinoma with MET Copy Number Gain but without a High-Level MET/CEP7 Ratio, MET Overexpression, or Exon 14 Splicing Mutations. *J Thorac Oncol* 11(5):e59-62
- 308 Caparica R, Yen CT, Coudry R, et al. (2017) Responses to Crizotinib Can Occur in High-Level MET-Amplified Non-Small Cell Lung Cancer Independent of MET Exon 14 Alterations. *J Thorac Oncol* 12(1):141-144
- 309 York ER, Varella-Garcia M, Bang TJ, et al. (2017) Tolerable and Effective Combination of Full-Dose Crizotinib and Osimertinib Targeting MET Amplification Sequentially Emerging after T790M Positivity in EGFR-Mutant Non-Small Cell Lung Cancer. *J Thorac Oncol* ePub Mar 2017
- 310 Drilon et al., 2016; ASCO Abstract 108
- 311 Schrock AB, Frampton GM, Suh J, et al. (2016) Characterization of 298 Patients with Lung Cancer Harboring MET Exon 14 Skipping Alterations. *J Thorac Oncol* ePub Jun 2016
- 312 Mahjoubi L, Gazzah A, Besse B, et al. (2016) A never-smoker lung adenocarcinoma patient with a MET exon 14 mutation (D1028N) and a rapid partial response after crizotinib. *Invest New Drugs* 34(3):397-8
- 313 Benderra MA, Aspeslagh S, Postel-Vinay S, et al. (2016) Acquired EGFR Mutation as the Potential Resistance Driver to Crizotinib in a MET-Mutated Tumor. *J Thorac Oncol* 11(2):e21-3
- 314 Cappuzzo F, Hirsch FR, Rossi E, et al. (2005) Epidermal growth factor receptor gene and protein and gefitinib sensitivity in non-small-cell lung cancer. *J Natl Cancer Inst* 97(9):643-55
- 315 Zhang X, Zhang Y, Tang H, et al. (2017) EGFR gene copy number as a predictive/biomarker for patients with non-small-cell lung cancer receiving tyrosine kinase inhibitor treatment: a systematic review and meta-analysis. *J Investig Med* 65(1):72-81
- 316 Dahabreh IJ, Linardou H, Kosmidis P, et al. (2011) EGFR gene copy number as a predictive biomarker for patients receiving tyrosine kinase inhibitor treatment: a systematic review and meta-analysis in non-small-cell lung cancer. *Ann Oncol* 22(3):545-52
- 317 Dahabreh IJ, Linardou H, Sianni F, et al. (2010) Somatic EGFR mutation and gene copy gain as predictive biomarkers for response to tyrosine kinase inhibitors in non-small cell lung cancer. *Clin Cancer Res* 16(1):291-303
- 318 Shepherd FA, Rodrigues Pereira J, Ciuleanu T, et al. (2005) Erlotinib in previously treated non-small-cell lung cancer. *N Engl J Med* 353(2):123-32
- 319 Cappuzzo F, Ciuleanu T, Stelmakh L, et al. (2010) Erlotinib as maintenance treatment in advanced non-small-cell lung cancer: a multicentre, randomised, placebo-controlled phase 3 study. *Lancet Oncol* 11(6):521-9
- 320 Garassino MC, Martelli O, Brogginini M, et al. (2013) Erlotinib versus docetaxel as second-line treatment of patients with advanced non-small-cell lung cancer and wild-type EGFR tumours (TAILOR): a randomised controlled trial. *Lancet Oncol* 14(10):981-8
- 321 Kawaguchi T, Ando M, Asami K, et al. (2014) Randomized phase III trial of erlotinib versus docetaxel as second- or third-line therapy in patients with advanced non-small-cell lung cancer: Docetaxel and Erlotinib Lung Cancer Trial (DELTA). *J Clin Oncol* 32(18):1902-8
- 322 Liu J, Sheng Z, Zhang Y, et al. (2016) The Efficacy of Epidermal Growth Factor Receptor Tyrosine Kinase Inhibitors for Molecularly Selected Patients with Non-Small Cell Lung Cancer: A Meta-Analysis of 30 Randomized Controlled Trials. *Target Oncol* 11(1):49-58
- 323 Reck M, van Zandwijk N, Gridelli C, et al. (2010) Erlotinib in advanced non-small cell lung cancer: efficacy and safety findings of the global phase IV Tarceva Lung Cancer Survival Treatment study. *J Thorac Oncol* 5(10):1616-22
- 324 Fukuoka M, Wu YL, Thongprasert S, et al. (2011) Biomarker analyses and final overall survival results from a phase III, randomized, open-label, first-line study of gefitinib versus carboplatin/paclitaxel in clinically selected patients with advanced non-small-cell lung cancer in Asia (IPASS). *J Clin Oncol* 29(21):2866-74
- 325 Mok TS, Wu YL, Thongprasert S, et al. (2009) Gefitinib or carboplatin-paclitaxel in pulmonary adenocarcinoma. *N Engl J Med* 361(10):947-57
- 326 Cross DA, Ashton SE, Ghiorghiu S, et al. (2014) AZD9291, an Irreversible EGFR TKI, Overcomes T790M-Mediated Resistance to EGFR Inhibitors in Lung Cancer. *Cancer Discov* 4(9):1046-1061
- 327 Mok TS, Wu YL, Ahn MJ, et al. (2016) Osimertinib or Platinum-Pemetrexed in EGFR T790M-Positive Lung Cancer. *N Engl J Med* ePub Dec 2016
- 328 Yang et al., 2016; ELCC Abstract LBA2_PR
- 329 Goss G, Tsai CM, Shepherd FA, et al. (2016) Osimertinib for pretreated EGFR Thr790Met-positive advanced non-small-cell lung cancer (AURA2): a multicentre, open-label, single-arm, phase 2 study. *Lancet Oncol* 17(12):1643-1652
- 330 Ramalingam et al., 2017; ESMO Abstract LBA2_PR
- 331 Ramalingam SS, Yang JC, Lee CK, et al. (2017) Osimertinib As First-Line Treatment of EGFR Mutation-Positive Advanced Non-Small-Cell Lung Cancer. *J Clin Oncol* :JCO2017747576
- 332 Ramalingam et al., 2015; ASCO Abstract 2509
- 333 Miyauchi E, Ichinose M, Inoue A (2017) Successful Osimertinib Rechallenge in a Patient with T790M-Mutant Non-Small Cell Lung Cancer after Osimertinib-Induced Interstitial Lung Disease. *J Thorac Oncol* 12(5):e59-e61
- 334 Yoshida H, Kim YH (2017) Successful Osimertinib Rechallenge after Severe Osimertinib-Induced Hepatotoxicity. *J Thorac Oncol* 12(5):e61-e63
- 335 Junttila TT, Li G, Parsons K, et al. (2011) Trastuzumab-DM1 (T-DM1) retains all the mechanisms of action of trastuzumab and efficiently inhibits growth of lapatinib insensitive breast cancer. *Breast Cancer Res Treat* 128(2):347-56
- 336 Lewis Phillips GD, Li G, Dugger DL, et al. (2008) Targeting HER2-positive breast cancer with trastuzumab-DM1, an antibody-cytotoxic drug conjugate. *Cancer Res* 68(22):9280-90
- 337 Erickson HK, Park PU, Widdison WC, et al. (2006) Antibody-maytansinoid conjugates are activated in targeted cancer cells by lysosomal degradation and linker-dependent intracellular processing. *Cancer Res* 66(8):4426-33
- 338 Weiler D, Diebold J, Strobel K, et al. (2015) Rapid response to trastuzumab emtansine in a patient with HER2-driven lung cancer. *J Thorac Oncol* 10(4):e16-7
- 339 Mazières J, Barlesi F, Filleron T, et al. (2015) Lung cancer patients with HER2 mutations treated with chemotherapy and HER2-targeted drugs: Results from the European EUHER2 cohort. *Ann Oncol* ePub Nov 2015

APPENDIX

References Associated with Professional Services Content

- 340 Krop IE, Kim SB, González-Martín A, et al. (2014) Trastuzumab emtansine versus treatment of physician's choice for pretreated HER2-positive advanced breast cancer (TH3RESA): a randomised, open-label, phase 3 trial. *Lancet Oncol* 15(7):689-99
- 341 Verma S, Miles D, Gianni L, et al. (2012) Trastuzumab emtansine for HER2-positive advanced breast cancer. *N Engl J Med* 367(19):1783-91
- 342 Welslau M, Diéras V, Sohn JH, et al. (2014) Patient-reported outcomes from EMILIA, a randomized phase 3 study of trastuzumab emtansine (T-DM1) versus capecitabine and lapatinib in human epidermal growth factor receptor 2-positive locally advanced or metastatic breast cancer. *Cancer* 120(5):642-51
- 343 Krop IE, LoRusso P, Miller KD, et al. (2012) A phase II study of trastuzumab emtansine in patients with human epidermal growth factor receptor 2-positive metastatic breast cancer who were previously treated with trastuzumab, lapatinib, an anthracycline, a taxane, and capecitabine. *J Clin Oncol* 30(26):3234-41
- 344 Burris HA, Rugo HS, Vukelja SJ, et al. (2011) Phase II study of the antibody drug conjugate trastuzumab-DM1 for the treatment of human epidermal growth factor receptor 2 (HER2)-positive breast cancer after prior HER2-directed therapy. *J Clin Oncol* 29(4):398-405
- 345 Klempner SJ, Borghei A, Hakimian B, et al. (2016) Brief Report: Intracranial Activity of Cabozantinib in MET exon14 Positive NSCLC with Brain Metastases. *J Thorac Oncol* ePub Sep 2016
- 346 Drilon A, Rekhtman N, Arcila M, et al. (2016) Cabozantinib in patients with advanced RET-rearranged non-small-cell lung cancer: an open-label, single-centre, phase 2, single-arm trial. *Lancet Oncol* 17(12):1653-1660
- 347 Drilon A, Wang L, Hasanovic A, et al. (2013) Response to Cabozantinib in patients with RET fusion-positive lung adenocarcinomas. *Cancer Discov* 3(6):630-5
- 348 Gautschi et al., 2016; ASCO Abstract 9014
- 349 Mukhopadhyay S, Pennell NA, Ali SM, et al. (2014) RET-Rearranged Lung Adenocarcinomas with Lymphangitic Spread, Psammoma Bodies, and Clinical Responses to Cabozantinib. *J Thorac Oncol* 9(11):1714-1719
- 350 Hellerstedt et al., 2012; ASCO Abstract 7514
- 351 Neal JW, Dahlberg SE, Wakelee HA, et al. (2016) Erlotinib, cabozantinib, or erlotinib plus cabozantinib as second-line or third-line treatment of patients with EGFR wild-type advanced non-small-cell lung cancer (ECOG-ACRIN 1512): a randomised, controlled, open-label, multicentre, phase 2 trial. *Lancet Oncol* 17(12):1661-1671
- 352 Jiang Z, Li C, Li F, et al. (2013) EGFR gene copy number as a prognostic marker in colorectal cancer patients treated with cetuximab or panitumumab: a systematic review and meta analysis. *PLoS ONE* 8(2):e56205
- 353 Licitra L, Mesia R, Rivera F, et al. (2011) Evaluation of EGFR gene copy number as a predictive biomarker for the efficacy of cetuximab in combination with chemotherapy in the first-line treatment of recurrent and/or metastatic squamous cell carcinoma of the head and neck: EXTREME study. *Ann Oncol* 22(5):1078-87
- 354 Ramalingam SS, Kotsakis A, Tarhini AA, et al. (2013) A multicenter phase II study of cetuximab in combination with chest radiotherapy and consolidation chemotherapy in patients with stage III non-small cell lung cancer. *Lung Cancer* 81(3):416-21
- 355 Kim ES, Neubauer M, Cohn A, et al. (2013) Docetaxel or pemetrexed with or without cetuximab in recurrent or progressive non-small-cell lung cancer after platinum-based therapy: a phase 3, open-label, randomised trial. *Lancet Oncol* 14(13):1326-36
- 356 Wheler JJ, Tsimberidou AM, Falchook GS, et al. (2013) Combining erlotinib and cetuximab is associated with activity in patients with non-small cell lung cancer (including squamous cell carcinomas) and wild-type EGFR or resistant mutations. *Mol Cancer Ther* 12(10):2167-75
- 357 Ross HJ, Blumenschein GR, Aisner J, et al. (2010) Randomized phase II multicenter trial of two schedules of lapatinib as first- or second-line monotherapy in patients with advanced or metastatic non-small cell lung cancer. *Clin Cancer Res* 16(6):1938-49
- 358 Diaz R, Nguewa PA, Parrondo R, et al. (2010) Antitumor and antiangiogenic effect of the dual EGFR and HER-2 tyrosine kinase inhibitor lapatinib in a lung cancer model. *BMC Cancer* 10:188
- 359 Burris HA, Taylor CW, Jones SF, et al. (2009) A phase I and pharmacokinetic study of oral lapatinib administered once or twice daily in patients with solid malignancies. *Clin Cancer Res* 15(21):6702-8
- 360 Crawford J, Swanson P, Schwarzenberger P, et al. (2013) A phase 2 randomized trial of paclitaxel and carboplatin with or without panitumumab for first-line treatment of advanced non-small-cell lung cancer. *J Thorac Oncol* 8(12):1510-8
- 361 Schuette W, Behringer D, Stoecklacher J, et al. (2015) CHAMP: A Phase II Study of Panitumumab With Pemetrexed and Cisplatin Versus Pemetrexed and Cisplatin in the Treatment of Patients With Advanced-Stage Primary Nonsquamous Non-Small-Cell Lung Cancer With Particular Regard to the KRAS Status. *Clin Lung Cancer* ePub Jun 2015
- 362 Baselga J, Cortés J, Kim SB, et al. (2012) Pertuzumab plus trastuzumab plus docetaxel for metastatic breast cancer. *N Engl J Med* 366(2):109-19
- 363 Swain SM, Kim SB, Cortés J, et al. (2013) Pertuzumab, trastuzumab, and docetaxel for HER2-positive metastatic breast cancer (CLEOPATRA study): overall survival results from a randomised, double-blind, placebo-controlled, phase 3 study. *Lancet Oncol* 14(6):461-71
- 364 Agus DB, Gordon MS, Taylor C, et al. (2005) Phase I clinical study of pertuzumab, a novel HER dimerization inhibitor, in patients with advanced cancer. *J Clin Oncol* 23(11):2534-43
- 365 Yamamoto N, Yamada Y, Fujiwara Y, et al. (2009) Phase I and pharmacokinetic study of HER2-targeted rhuMab 2C4 (Pertuzumab, RO4368451) in Japanese patients with solid tumors. *Jpn J Clin Oncol* 39(4):260-6
- 366 Herbst RS, Davies AM, Natale RB, et al. (2007) Efficacy and safety of single-agent pertuzumab, a human epidermal receptor dimerization inhibitor, in patients with non small cell lung cancer. *Clin Cancer Res* 13(20):6175-81
- 367 Felip E, Ranson M, Cedrés S, et al. (2012) A phase Ib, dose-finding study of erlotinib in combination with a fixed dose of pertuzumab in patients with advanced non-small-cell lung cancer. *Clin Lung Cancer* 13(6):432-41
- 368 Punnoose EA, Atwal S, Liu W, et al. (2012) Evaluation of circulating tumor cells and circulating tumor DNA in non-small cell lung cancer: association with clinical endpoints in a phase II clinical trial of pertuzumab and erlotinib. *Clin Cancer Res* 18(8):2391-401
- 369 Hughes B, Mileskin L, Townley P, et al. (2014) Pertuzumab and Erlotinib in Patients With Relapsed Non-Small Cell Lung Cancer: A Phase II Study Using 18F-Fluorodeoxyglucose Positron Emission Tomography/Computed Tomography Imaging. *Oncologist* 19(2):175-6
- 370 Lara PN, Laptalo L, Longmate J, et al. (2004) Trastuzumab plus docetaxel in HER2/neu-positive non-small-cell lung cancer: a California Cancer Consortium screening and phase II trial. *Clin Lung Cancer* 5(4):231-6
- 371 Krug LM, Miller VA, Patel J, et al. (2005) Randomized phase II study of weekly docetaxel plus trastuzumab versus weekly paclitaxel plus trastuzumab in patients with previously untreated advanced nonsmall cell lung carcinoma. *Cancer* 104(10):2149-55
- 372 Hainsworth et al., 2016; ASCO Abstract LBA11511
- 373 Cappuzzo F, Bemis L, Varella-Garcia M (2006) HER2 mutation and response to trastuzumab therapy in non-small-cell lung cancer. *N Engl J Med* 354(24):2619-21
- 374 Mazières J, Peters S, Lepage B, et al. (2013) Lung cancer that harbors an HER2 mutation: epidemiologic characteristics and therapeutic perspectives. *J Clin Oncol* 31(16):1997-2003
- 375 Rugo et al., 2016; ASCO Abstract LBA503
- 376 Waller et al., 2016; ASCO Abstract 583
- 377 Audran et al., 2017; ASCO-SITC Clinical Immunology Symposium Abstract 10