

**MEDIA RELEASE • COMMUNIQUE AUX MEDIAS • MEDIENMITTEILUNG****Novartis announces JCO publication of Lutathera® NETTER-1 data showing significantly longer time to deterioration of key quality of life measures in patients with progressive midgut NETs**

- *Patients reported prolonged maintenance of global health status<sup>1</sup> (overall health)*
- *Longer time to deterioration (TTD) seen across multiple clinically relevant symptom categories including diarrhea, fatigue and pain<sup>1</sup>*
- *Functional health-related quality of life (HRQoL) categories, including those pertaining to basic and advanced activities of daily living, were sustained for a longer time<sup>1</sup>*

**Basel, June 7, 2018** – Novartis today announced that the *Journal of Clinical Oncology* has published results of an analysis of the impact of Lutathera® (lutetium Lu 177 dotatate\*) treatment on time to deterioration in HRQoL in the pivotal phase III NETTER-1 trial. The data demonstrate that treatment with Lutathera provides significantly longer time to deterioration of quality of life for patients with progressive midgut Neuroendocrine tumors (NETs) compared to octreotide LAR alone<sup>1</sup>. Lutathera is the first Peptide Receptor Radionuclide Therapy (PRRT) to receive regulatory registration, with approval by the European Commission in September 2017 and by the US Food and Drug Administration (FDA) in January 2018. Lutathera is an Advanced Accelerator Applications product.

“Neuroendocrine tumor progression is often associated with deterioration in quality of life due to tumor growth and production of hormones<sup>2,3</sup>,” said Jonathan Strosberg, MD, Associate Professor, Section Head, Neuroendocrine Tumor Program at Moffitt Cancer Center, and lead author of the publication. “In patients with advanced NETs, maintenance of an acceptable HRQoL is particularly important given the relatively long durations of treatment and overall survival compared to other advanced malignancies. These data provide hope for these patients and their families.”

TTD was significantly longer in the Lutathera arm versus the comparator arm for the following domains: global health status (self-assessment of overall health and quality of life), physical functioning, role functioning, fatigue, pain, diarrhea, disease related worries and body image<sup>1</sup>. Differences in median TTD were clinically significant in several domains: 28.8 months vs. 6.1 months for global health status, and 25.2 months vs. 11.5 months for physical functioning<sup>1</sup>. Domains where Lutathera treatment did not show a significant benefit in TTD include nausea and vomiting, dyspnea, insomnia, appetite loss, constipation, financial difficulties, endocrine scale (flushing, sweats), GI scale (bloating, flatulence), treatment scale, social functioning scale, muscle/bone pain, sexual function, and information/communication function. There were no domains in which TTD analysis showed significant benefit for the comparator arm<sup>1</sup>.

“The results demonstrated by Lutathera in the NETTER-1 study continue to be encouraging in a patient population with limited therapeutic options for control of progressive disease<sup>1</sup>,” said

Samit Hirawat, MD, Head of Novartis Oncology Global Drug Development. "We are focused on developing medicines that offer meaningful outcomes for patients, like Lutathera."

The NETTER-1 trial is an international phase III study in patients with progressive, somatostatin receptor-positive midgut neuroendocrine tumors. Patients were randomized to treatment with Lutathera and best supportive care (30 mg octreotide LAR), or 60 mg octreotide LAR alone<sup>4</sup>. European Organisation for Research and Treatment of Cancer (EORTC) quality of life questionnaires, a commonly used metric for analysis of HRQoL in cancer patients, were assessed during the trial to determine the impact of treatment on HRQoL<sup>5</sup>. Patients completed the questionnaires at baseline and every 12 weeks until tumor progression. TTD was defined as the time from randomization to the first QoL deterioration  $\geq 10$  points (on a 100-point scale) compared to baseline score for the same domain. Patients with no deterioration were censored at the last QoL assessment date. Patients with no baseline and/or no follow-up were censored at randomization. Analysis cut-off date was June 30, 2016. In total, 231 patients were randomized in the HRQoL study (117 in the Lutathera arm and 114 in the 60 mg octreotide LAR arm).

### **About Lutathera<sup>®</sup>**

Lutathera is a lutetium Lu 177-labeled somatostatin analog peptide. Lutathera belongs to a class of treatments called Peptide Receptor Radionuclide Therapy (PRRT). Lutathera is comprised of a targeting molecule which carries a radioactive component. Lutathera has received orphan drug designation from the FDA and the European Medicines Agency (EMA). In the US, Lutathera is indicated for the treatment of somatostatin receptor positive gastroenteropancreatic neuroendocrine tumors (GEP-NETs), including foregut, midgut, and hindgut neuroendocrine tumors in adults<sup>6</sup>. In Europe, Lutathera is indicated for the treatment of unresectable or metastatic, progressive, well differentiated (G1 and G2), somatostatin receptor positive gastroenteropancreatic neuroendocrine tumors (GEP-NETs) in adults<sup>7</sup>. Lutathera can cause serious side effects that may include bone marrow problems, kidney problems, liver problems, hormonal gland problems, fertility problems and problems arising from radiation exposure. Please see Important Safety Information and Full Prescribing Information at: [www.lutathera.com](http://www.lutathera.com).

\* USAN: lutetium Lu 177 dotatate / INN: lutetium (177Lu) oxodotretotide

### **Disclaimer**

This press release contains forward-looking statements within the meaning of the United States Private Securities Litigation Reform Act of 1995. Forward-looking statements can generally be identified by words such as "potential," "can," "will," "plan," "expect," "anticipate," "look forward," "believe," "committed," "investigational," "pipeline," "launch," or similar terms, or by express or implied discussions regarding potential marketing approvals, new indications or labeling for the investigational or approved products described in this press release, or regarding potential future revenues from such products. You should not place undue reliance on these statements. Such forward-looking statements are based on our current beliefs and expectations regarding future events, and are subject to significant known and unknown risks and uncertainties. Should one or more of these risks or uncertainties materialize, or should underlying assumptions prove incorrect, actual results may vary materially from those set forth in the forward-looking statements. There can be no guarantee that the investigational or approved products described in this press release will be submitted or approved for sale or for any additional indications or labeling in any market, or at any particular time. Nor can there be any guarantee that such products will be commercially successful in the future. In particular, our expectations regarding such products could be affected by, among other things, the uncertainties inherent in research and development, including clinical trial results and additional analysis of existing clinical data; regulatory actions or delays or government regulation generally; global trends toward health care cost containment, including government, payor and general public pricing and reimbursement pressures; our ability to obtain or maintain proprietary intellectual property protection; the particular prescribing

preferences of physicians and patients; general political and economic conditions; safety, quality or manufacturing issues; potential or actual data security and data privacy breaches, or disruptions of our information technology systems, and other risks and factors referred to in Novartis AG's current Form 20-F on file with the US Securities and Exchange Commission. Novartis is providing the information in this press release as of this date and does not undertake any obligation to update any forward-looking statements contained in this press release as a result of new information, future events or otherwise.

### **About Advanced Accelerator Applications**

Advanced Accelerator Applications, a Novartis company, is an innovative radiopharmaceutical company developing, producing and commercializing nuclear medicine theragnostics. AAA is also an established leader in radiopharmaceuticals for Positron Emission tomography (PET) and Single-Photon Emission Computed Tomography (SPECT) diagnostic imaging, mainly used in clinical oncology, cardiology and neurology. For more information, please visit: <https://www.adacap.com/>.

### **About Novartis**

Novartis provides innovative healthcare solutions that address the evolving needs of patients and societies. Headquartered in Basel, Switzerland, Novartis offers a diversified portfolio to best meet these needs: innovative medicines, cost-saving generic and biosimilar pharmaceuticals and eye care. Novartis has leading positions globally in each of these areas. In 2017, the Group achieved net sales of USD 49.1 billion, while R&D throughout the Group amounted to approximately USD 9.0 billion. Novartis Group companies employ approximately 124,000 full-time-equivalent associates. Novartis products are sold in approximately 155 countries around the world. For more information, please visit <http://www.novartis.com>.

Novartis is on Twitter. Sign up to follow @Novartis at <http://twitter.com/novartis>  
For Novartis multimedia content, please visit [www.novartis.com/news/media-library](http://www.novartis.com/news/media-library)  
For questions about the site or required registration, please contact [media.relations@novartis.com](mailto:media.relations@novartis.com)

### **References**

1. Strosberg J, Wolin E, Chasen B, et al. Health related quality of life in patients with progressive midgut neuroendocrine tumors treated with 177Lu-Dotatate in the phase III NETTER-1 Trial. JCO 2018.
2. Beaumont JL, Cella D, Phan AT, Choi S, Liu Z, Yao JC. Comparison of health-related quality of life in patients with neuroendocrine tumors with quality of life in the general US population. Pancreas 2012; 41(3): 461-6.
3. Singh S, Granberg D, Wolin E, et al. Patient-Reported Burden of a Neuroendocrine Tumor (NET) Diagnosis: Results From the First Global Survey of Patients With NETs. J Glob Oncol 2017; 3(1): 43-53.
4. Strosberg J, El-Haddad G, Wolin E, et al. Phase 3 Trial of 177Lu-Dotatate for Midgut Neuroendocrine Tumors. N Engl J Med 2017; 376:125-35.
5. Aaronson NK, Ahmedzai S, Bergman B, et al. The European Organization for Research and Treatment of Cancer QLQ-C30: a quality-of-life instrument for use in international clinical trials in oncology. Journal of the National Cancer Institute 1993; 85(5): 365-76.
6. LUTATHERA Prescribing Information
7. LUTATHERA Summary of Product Characteristics

# # #

### **Novartis Media Relations**

Central media line: +41 61 324 2200  
E-mail: [media.relations@novartis.com](mailto:media.relations@novartis.com)

Eric Althoff  
Novartis Global Media Relations  
+41 61 324 7999 (direct)  
+41 79 593 4202 (mobile)  
[eric.althoff@novartis.com](mailto:eric.althoff@novartis.com)

Rachel Levine  
Advanced Accelerator Applications  
+212 235 2395 (direct)  
+917 375 2935 (mobile)  
[rachel.levine@adacap.com](mailto:rachel.levine@adacap.com)

**Novartis Investor Relations**

Central investor relations line: +41 61 324 7944

E-mail: [investor.relations@novartis.com](mailto:investor.relations@novartis.com)

Central		North America	
Samir Shah	+41 61 324 7944	Richard Pulik	+1 212 830 2448
Pierre-Michel Bringer	+41 61 324 1065	Cory Twining	+1 212 830 2417
Thomas Hungerbuehler	+41 61 324 8425		
Isabella Zinck	+41 61 324 7188		