

Ready Reckoner 10 – SCI – Central Cord Syndrome

Table 23.94 Central Cord Syndrome C4: Tetraplegia, Sensory Impairment-Sacral Sparing

Upper Extremity:	26%		
MAS '2'			
Lower Extremity:			
MAS '1' (26%)			
Upper Extremity:	Sensory	56%	
MAS '2'	Impairment with		
Lower Extremity:	Sacral Sparing		
MAS '1' (26%)			
Upper Extremity:	35%		
MAS '3'			
Lower Extremity:			
MAS '1+' (35%)			
Upper Extremity:	Sensory	79%	
MAS '3'	Impairment with		
Lower Extremity:	Sacral Sparing		
MAS '1+' (35%)			
Upper Extremity:	Sensory	Retention of	89%
MAS '3'	Impairment with	Urine (50%)	
Lower Extremity:	Sacral Sparing		
MAS '1+' (35%)			

Upper Extremity:	Sensory	Retention of	Constipation	91%
MAS '3'	Impairment with	Urine (50%)	Class '3' (15%)	
Lower Extremity:	Sacral Sparing			
MAS '1+' (26%)				

Table 23.95 Central Cord Syndrome C5: Tetraplegia, Sensory Impairment-Sacral Sparing

Upper Extremity:	25%	
MAS '2'		
Lower Extremity:		
MAS '1' (25%)		
Upper Extremity:	Sensory	56%
MAS '2'	Impairment with	
Lower Extremity:	Sacral Sparing	
MAS '1' (29%)		
Upper Extremity:	33%	
MAS '3'		
Lower Extremity:		
MAS '1+' (33%)		
Upper Extremity:	Sensory	75%
MAS '3'	Impairment with	
Lower Extremity:	Sacral Sparing	
MAS '1+' (33%)		

Upper Extremity:	Sensory	Retention of	88%
MAS '3'	Impairment with	Urine (50%)	
Lower Extremity:	Sacral Sparing		
MAS '1+' (33%)			
Upper Extremity:	Sensory	Retention of	Constipation 89%
MAS '3'	Impairment with	Urine (50%)	Class '3' (15%)
Lower Extremity:	Sacral Sparing		
MAS '1+' (33%)			

Table 23.96 Central Cord Syndrome C6: Tetraplegia, Sensory Impairment-Sacral Sparing

Upper Extremity:	23%
MAS '2'	
Lower Extremity:	
MAS '1'	
(23%)	
Upper Extremity:	Sensory 48%
MAS '2'	Impairment with
Lower Extremity:	Sacral Sparing
MAS '1' (23%)	
Upper Extremity:	33%
MAS '3'	
Lower Extremity:	
MAS '1+' (33%)	

Upper Extremity:	Sensory	65%		
MAS '3'	Impairment with			
Lower Extremity:	Sacral Sparing			
MAS '1+' (33%)				
Upper Extremity:	Sensory	Retention of	83%	
MAS '3'	Impairment with	Urine (50%)		
Lower Extremity:	Sacral Sparing			
MAS '1+' (33%)				
Upper Extremity:	Sensory	Retention of	Constipation	85%
MAS '3'	Impairment with	Urine (50%)	Class '3' (15%)	
Lower Extremity:	Sacral Sparing			
MAS '1+' (33%)				

Table 23.97 Central Cord Syndrome C7: Tetraplegia, Sensory Impairment-Sacral Sparing

Upper Extremity:	24%		
MAS '2'			
Lower Extremity:			
MAS '1' (24%)			
Upper Extremity:	Sensory	44%	
MAS '2'	Impairment with		
Lower Extremity:	Sacral Sparing		
MAS '1' (24%)			
Upper Extremity:	31%		
MAS '3'			

Lower Extremity:

MAS '1+' (31%)

Upper Extremity: Sensory 59%

MAS '3' Impairment with

Lower Extremity: Sacral Sparing

MAS '1+' (31%)

Upper Extremity: Sensory Retention of 79%

MAS '3' Impairment with Urine (50%)

Lower Extremity: Sacral Sparing

MAS '1+' (31%)

Upper Extremity: Sensory Retention of Constipation 82%

MAS '3' Impairment with Urine (50%) Class '3' (15%)

Lower Extremity: Sacral Sparing

MAS '1+' (31%)
